

# COUNTING AND DIMENSIONS

EHUD HRUSHOVSKI<sup>1</sup> AND FRANK WAGNER<sup>2</sup>

ABSTRACT. We prove a theorem comparing a well-behaved dimension notion to a second, more rudimentary dimension. Specialising to a non-standard counting measure, this generalizes a theorem of Larsen and Pink on an asymptotic upper bound for the intersection of a variety with a general finite subgroup of an algebraic group. As a second application we apply this to bad fields of positive characteristic, to give an asymptotic estimate for the number of  $\mathbb{F}_q$ -rational points of a definable multiplicative subgroup similar to the Lang-Weil estimate for curves over finite fields.

## INTRODUCTION

In [1] Larsen and Pink show that if  $H$  is a “sufficiently general” finite subgroup of a connected almost simple algebraic group  $G$ , then for any subvariety  $X$  of  $G$

$$|H \cap X| \leq c \cdot |H|^{\dim(X)/\dim(G)},$$

where the constant  $c$  depends only on the form of  $G$  and  $X$ , but not on  $H$  (in other words,  $G$  and  $X$  are allowed to vary in a constructible family). This theorem was recast (in somewhat greater generality) in model-theoretic form by the first author of the present paper, and re-discovered by the second author in the context of bad fields. In the general form it allows to give an upper bound, for suitable minimal structures with a well-behaved dimension  $d$ , of a rudimentary dimension  $\delta$  (which may for instance be derived from counting measure in a quasi-finite subset) in terms of the original dimension  $d$ , typically giving Larsen-Pink like estimates for increasing families of finite subsets. We offer two proofs of the theorem: a more rapid one using types, and a more explicit construction using definable sets. The latter proof could in principle be used to get effective estimates on the constant  $c$ .

---

*Date:* July 14, 2005.

<sup>1</sup> Supported by Israel Science Foundation grant no. 244/03

<sup>2</sup> Membre junior de l’Institut universitaire de France.

## 1. THE MAIN THEOREM

**Definition 1.** Let  $\mathfrak{M}$  be an uncountably saturated structure. A *dimension theory* on  $\mathfrak{M}$  is an automorphism-invariant map  $d$  from the class of definable sets into  $\mathbb{Z}$ , together with a formal element  $-\infty$ , satisfying

- (1)  $d(\emptyset) = -\infty$  and  $d(\{x\}) = 0$  for any point  $x$ .
- (2)  $d(X \cup Y) = \max\{d(X), d(Y)\}$ .
- (3) Let  $f : X \rightarrow Y$  be a definable map.
  - (a) If  $d(f^{-1}(y)) = n$  for all  $y \in Y$ , then  $d(X) = d(Y) + n$ , for all  $n \in \mathbb{Z} \cup \{-\infty\}$ .
  - (b)  $\{y \in Y : d(f^{-1}(y)) = n\}$  is definable for all  $n \in \mathbb{Z} \cup \{-\infty\}$ .

It follows that  $d(X \times Y) = d(X) + d(Y)$ , and  $d(X) = d(Y)$  if  $X$  and  $Y$  are definably isomorphic. By uncountable saturation,  $d(f^{-1}(y))$  takes only finitely many values for  $y \in Y$ .

**Definition 2.** For a partial type  $\pi$  let  $d(\pi) := \min\{d(X) : X \in \pi\}$ ; note that the minimum is necessarily attained. If  $p = \text{tp}(x/A)$ , put  $d(x/A) := d(p)$ .

For two partial types  $\pi, \pi'$  over  $A$  let

$$\pi \otimes_A \pi' := (\pi \times \pi') \cup \{\neg X : X \text{ } A\text{-definable, } d(X) < d(\pi) + d(\pi')\}.$$

**Definition 3.** Let  $\mathfrak{M}$  be a structure with dimension  $d$ . A subset  $F \subset M^3$  is a *correspondence* on  $\mathfrak{M}$  if the projection to the first two coordinates is surjective with 0-dimensional fibres. We put

$$\begin{aligned} F(X) &:= \{y \in M : \models \exists(x, x') \in X \ F(x, x', y)\}, \text{ and} \\ F^{-1}(y) &:= \{(x, x') \in M^2 : \models F(x, x', y)\}. \end{aligned}$$

If  $\mathcal{F}$  is a set of correspondences,  $\mathfrak{M}$  is  $\mathcal{F}$ -*minimal* if for any  $A$  and partial 1-types  $\pi, \pi'$  over  $A$  with  $0 < d(\pi) \leq d(\pi') < d(M)$  and a partial type  $\rho$  over  $A$  extending  $\pi \otimes_A \pi'$ , there is  $F \in \mathcal{F}$  with  $d(F(\rho)) > d(\pi')$ .

**Lemma 1.** *The following are equivalent:*

- (1)  $\mathfrak{M}$  is  $\mathcal{F}$ -minimal.
- (2) For any  $x, x' \in M$  and parameters  $A$  with  $0 < d(x/A) \leq d(x'/A) < d(M)$  and  $d(xx'/A) = d(x/A) + d(x'/A)$  there is  $F \in \mathcal{F}$  and  $y \in F(xx')$  with  $d(F^{-1}(y) \cap \text{tp}(xx'/A)) < d(x/A)$ .
- (3) For any  $A$  and  $A$ -definable  $X, X'$  with  $0 < d(X) \leq d(X') < d(M)$  and  $(x, x') \in X \times X'$  there is  $A$ -definable  $W \subseteq X \times X'$  with  $(x, x') \in W$  such that either  $d(W) < d(X \times X')$  or  $d(F^{-1}(y) \cap W) < d(X)$  for some  $F \in \mathcal{F}$  and  $y \in F(xx')$ .

*Proof:* Suppose  $\mathfrak{M}$  is  $\mathcal{F}$ -minimal, and consider  $x, x', A$  as in (2). Put  $\pi = \text{tp}(x/A)$ ,  $\pi' = \text{tp}(x'/A)$  and  $\rho := \text{tp}(xx'/A)$ . Since  $d(xx'/A) =$

$d(x/A) + d(x'/A)$  we have  $\rho \subseteq \pi \otimes_A \pi'$ , so there is  $F \in \mathcal{F}$  with  $d(F(\rho)) > d(\pi')$ . In particular there is  $y \in F(xx')$  with  $d(y/A) > d(x'/A)$ . Choose  $A$ -definable  $W \in \text{tp}(xx'/A)$  with  $d(W) = d(xx'/A)$  and  $k = d(F^{-1}(y) \cap \text{tp}(xx'/A)) = d(F^{-1}(y) \cap W)$ , and  $A$ -definable  $Y \in \text{tp}(y/A)$  with  $d(F^{-1}(y') \cap W) = k$  for all  $y' \in Y$ . Then

$$\begin{aligned} d(x/A) + d(x'/A) &= d(xx'/A) = d(\rho) = d(W) \geq d(F \cap (W \times Y)) \\ &= d(Y) + k \geq d(y/A) + k > d(x'/A) + k, \end{aligned}$$

whence  $d(x/A) > k = d(F^{-1}(y) \cap \text{tp}(xx'/A))$ .

For the converse, consider partial types  $\pi, \pi'$  and  $\rho$  over  $A$  as in the definition of  $\mathcal{F}$ -minimality, and take  $xx' \models \rho$ . Since  $\rho \supseteq \pi \otimes_A \pi'$  we have  $d(\pi) = d(x/A), d(\pi') = d(x'/A), d(\rho) = d(xx'/A)$  and  $d(x/A) + d(x'/A) = d(xx'/A)$ . By (2) there is  $F \in \mathcal{F}$  and  $y \in F(xx')$  with  $d(F^{-1}(y) \cap \text{tp}(xx'/A)) < d(x/A)$ . Choose  $A$ -definable  $W \in \text{tp}(xx'/A)$  with  $k = d(F^{-1}(y) \cap \text{tp}(xx'/A)) = d(F^{-1}(y) \cap W)$ , and  $A$ -definable  $Y \in \text{tp}(y/A)$  with  $d(Y) = d(y/A)$  and  $d(F^{-1}(y') \cap W) = k$  for all  $y' \in Y$ . Then

$$\begin{aligned} d(x/A) + d(x'/A) &= d(xx'/A) = d(\rho) \leq d(F \cap (W \times Y)) \\ &= d(Y) + k = d(y/A) + k < d(y/A) + d(x/A), \end{aligned}$$

whence  $d(F(\rho)) \geq d(y/A) > d(x'/A) = d(\pi')$ .

The equivalence (2)  $\Leftrightarrow$  (3) follows from the fact that for any partial type  $\pi$  there is  $X \in \pi$  with  $d(\pi) = d(X)$ .  $\square$

**Example 1.** Let  $G$  be a simple algebraic group (or more generally, a simple group of finite Morley rank). Let  $\mathcal{F}$  be the collection of maps  $F_c(x, y) = cx^{-1}c^{-1}y$ , where  $c$  runs over a countable Zariski-dense subgroup  $\Gamma$  (respectively, subgroup  $\Gamma$  not contained in any proper definable subgroup of  $G$ ). Then  $G$  is  $\mathcal{F}$ -minimal.

*Proof:* In any group of finite Morley rank,  $d = \text{RM}$  is additive and definable. So consider  $A \supseteq \Gamma$  and  $x \downarrow_A x'$  with  $0 < \text{RM}(x/A) \leq \text{RM}(x'/A)$ , and suppose  $\text{RM}(x, x'/cx^{-1}c^{-1}x', A) \geq \text{RM}(x/A)$  for all  $c \in \Gamma$ . Then  $x \downarrow_A cx^{-1}c^{-1}x'$ , whence  $x^{-c^{-1}} \downarrow_A x^{-c^{-1}}x'$  for all  $c \in \Gamma$ . So for any two independent realizations  $x_0, x_1$  of  $\text{stp}(x/A, x')$  both  $x_0^{-c^{-1}}x'$  and  $x_1^{-c^{-1}}x'$  satisfy the unique non-forking extension of  $\text{stp}(x^{-c^{-1}}x'/A)$  to  $A, x_0, x_1$ , and  $(x_0x_1^{-1})^{c^{-1}}x' \models \text{stp}(x'/A)$ . Since  $x_0, x_1 \downarrow_A x'$  this means that  $(x_0x_1^{-1})^{c^{-1}} \in \text{stab}(x'/A)$  for any two independent realisations  $x_0, x_1$  of  $\text{stp}(x/A)$ , and any  $c \in \Gamma$ . So this stabilizer is an infinite definable subgroup, as is the intersection  $H$  of its  $\Gamma$ -conjugates. But then the normalizer of  $H$  contains  $\Gamma$ , whence  $G$  by our choice of  $\Gamma$ ; since  $H$

is infinite and  $G$  is simple, we get  $H = G = \text{stab}(x'/A)$ . Therefore  $\text{tp}(x'/A)$  is generic, and  $\text{RM}(x'/A) = \text{RM}(G)$ .  $\square$

**Example 2.** A field of finite Morley rank is  $\{+, \times\}$ -minimal.

*Proof:* Suppose  $0 < \text{RM}(x/A) \leq \text{RM}(x'/A)$  and  $x \downarrow_A x'$ . If both  $\text{RM}(x, x'/x + x', A) \geq \text{RM}(x/A)$  and  $\text{RM}(x, x'/xx', A) \geq \text{RM}(x/A)$ , then  $x \downarrow_A x + x'$  and  $x \downarrow_A xx'$ . Let  $x_0, x_1$  be independent realizations of  $\text{stp}(x/A, x')$ . Since  $x_0 + x'$  and  $x_1 + x'$  realize the same strong type over  $A$ , they realize the same non-forking extension to  $A, x_0, x_1$ , and  $x_0 - x_1 + x' \models \text{stp}(x'/A)$ . As  $x' \downarrow_A x_0 - x_1$ , we get  $x_0 - x_1 \in \text{stab}^+(x'/A)$ ; similarly  $x_0 x_1^{-1} \in \text{stab}^\times(x'/A)$ . As  $x_0, x_1$  are independent non-algebraic, both stabilizers are infinite; note that obviously  $\text{stab}^+(x'/A)$  is  $\text{stab}^\times(x'/A)$ -invariant. However, in a field  $K$  of finite Morley rank the only definable additive subgroup  $A$  invariant under an infinite multiplicative subgroup is  $K$  itself (otherwise  $\{c \in K : cA \leq A\}$  would define an infinite subring, and hence an infinite subfield, a contradiction). Thus  $\text{stab}^+(x'/A) = K$ , and  $\text{RM}(x'/A) = \text{RM}(K)$ .  $\square$

**Definition 4.** Let  $\mathfrak{M}$  be any structure. A *quasi-dimension* on  $\mathfrak{M}$  is a map  $\delta$  from the class of definable sets into an ordered abelian group  $G$ , together with a formal element  $-\infty$ , satisfying

- (1)  $\delta(\emptyset) = -\infty$ , and  $\delta(\{x\}) = 0$  for any point  $x$ .
- (2)  $\delta(X \cup Y) = \max\{\delta(X), \delta(Y)\}$ , and  $\delta(X \times Y) = \delta(X) + \delta(Y)$ .
- (3) For any definable  $X \subseteq M^k$  and projection  $\pi$  to some of the coordinates, if  $\delta(\pi^{-1}(\bar{x})) \leq g$  for all  $\bar{x} \in \pi(X)$ , then  $\delta(X) \leq \delta(\pi(X)) + g$ , for all  $g \in G \cup \{-\infty\}$ .

We can now state the main theorem.

**Theorem 2.** *Let  $\mathfrak{M}$  be an  $\mathcal{F}$ -minimal structure, where  $\mathcal{F}$  is a set of  $\emptyset$ -definable correspondences for some dimension  $d$ . Let  $\delta$  be a quasi-dimension on  $\mathfrak{M}$  such that*

- (0)  $d(X) = 0$  implies  $\delta(X) \leq 0$  for all definable  $X$ .
- (4) For any  $F \in \mathcal{F}$  and definable  $X \subseteq M^2$ ,  $Y \subseteq M$  we have  $\delta(F \cap (X \times Y)) \geq \delta(X)$ , provided for all  $xx' \in X$  there is  $y \in Y$  with  $F(xx'y)$ .

Then  $d(M)\delta(X) \leq d(X)\delta(M)$  for any definable set  $X \subseteq M$ .

**Remark 1.** (1)  $\delta(F \cap (X \times Y)) \leq \delta(X)$  follows from axiom (3) and the fact that the fibres of the projection  $F \cap (X \times Y) \rightarrow X$  have  $d$ -dimension zero, and hence  $\delta$ -dimension zero.

- (2) Requirement (4) holds in particular if  $\mathcal{F}$  consists of definable functions, and  $\delta$  is invariant under definable bijections.

*Proof:* Clearly we may assume  $d(M) > 0$ . We use induction on  $d(X)$ . For  $d(X) = 0$  the assertion follows from condition (1). So suppose the assertion holds for dimension less than  $k$ , and  $d(X) = k$ . Put  $\alpha = \delta(M)/d(M)$  and suppose  $\delta(X) \geq \alpha k$ .

**Lemma 3.** *Let  $X, Y \subseteq M$  be  $B$ -definable with  $0 < d(X) \leq d(Y)$ . Then there is a  $B$ -definable finite partition  $X \times Y = W_0 \cup \dots \cup W_n$ , correspondences  $F_i \in \mathcal{F}$  and sets  $Z_i$  with  $F_i^{-1}(Z_i) = W_i$  for  $i > 0$ , such that*

- $d(W_i) = d(X) + d(Y)$  for  $i > 0$ , and  $d(W_0) < d(X) + d(Y)$ .
- for all  $i > 0$  we have  $d(Z_i) > d(Y)$ , and  $d(F_i^{-1}(z) \cap W_i) = d(X) + d(Y) - d(Z_i)$  for all  $z \in Z_i$ .

*Proof:* For  $F \in \mathcal{F}$  and  $B$ -definable  $W \subseteq X \times Y$  put

$$W_F := \{(x, y) \in W : \exists z \in F(xy) \ d(F^{-1}(z) \cap W) < d(X)\}, \text{ and}$$

$$Z_F := \{z \in F(W_F) : d(F^{-1}(z) \cap W) < d(X)\}.$$

By  $\mathcal{F}$ -minimality the  $B$ -definable sets

$$\{V \subset X \times Y : d(V) < d(X) + d(Y)\} \cup \{W_F : F \in \mathcal{F}, W \text{ } B\text{-definable}\}$$

cover  $X \times Y$ . By compactness a finite subset covers  $X \times Y$ ; shrinking the sets if necessary, we may assume that the sets form a partition of  $X \times Y$ . For  $i = d(X) - 1, d(X) - 2, \dots, 0$  partition every  $Z_F$  involved into parts

$$Z_F^i := \{z \in Z_F : d(F^{-1}(z) \cap (W_F \setminus \bigcup_{j>i} W_F^j)) = i\},$$

and put  $W_F^i = F^{-1}(Z_F^i) \cap (W_F \setminus \bigcup_{j>i} W_F^j)$ . Let  $W_0$  be the union of those sets of dimension strictly less than  $d(X) + d(Y)$ , and enumerate the others as  $W_1, \dots, W_n$  and  $Z_1, \dots, Z_n$ , respectively, with correspondences  $F_1, \dots, F_n$ . This satisfies the conditions.  $\square$

We inductively choose a tree of subsets of  $M$  with  $Y_\emptyset := X$  and  $d(Y_{\eta'}) < d(Y_\eta)$  whenever  $\eta' < \eta$  is a proper initial segment. Suppose we have found  $Y_\eta$ . If  $d(Y_\eta) = d(M)$  this branch stops. Otherwise put  $Y = Y_\eta$  in Lemma 3 and let  $Y_{\eta_i} := Z_i$  for  $i > 0$ . Put  $F_{\eta_i} := F_i$ ,  $W_{\eta_i} := W_i$ , and  $n_{\eta_i} := n_i = d(X) + d(Y_\eta) - d(Y_{\eta_i})$ . As  $d(Y_{\eta_i}) > d(Y_\eta)$  for all  $\eta$ , the tree is finite. Let  $m$  be the maximal length of a branch, and put  $m_\eta = m - |\eta|$ , where  $0 \leq |\eta| \leq m$  is the length of  $\eta$ .

**Lemma 4.** *If  $W \subset X^{m_{\eta i}} \times Y_{\eta i}$  with  $d(W) < d(X^{m_{\eta i}} \times Y_{\eta i})$ , then  $d((id_{X^{m_{\eta-1}}} \times F_{\eta i})^{-1}(W) \cap (X^{m_{\eta-1}} \times W_{\eta i})) < d(X^{m_{\eta}} \times Y_{\eta})$ .*

*Proof:* Since the fibres have constant dimension  $n_{\eta i}$ , we have

$$\begin{aligned} d((id_{X^{m_{\eta-1}}} \times F_{\eta i})^{-1}(W) \cap (X^{m_{\eta-1}} \times W_{\eta i})) &= d(W) + n_{\eta i} \\ &< d(X^{m_{\eta i}} \times Y_{\eta i}) + d(X) + d(Y_{\eta}) - d(Y_{\eta i}) \\ &= d(X^{m_{\eta}} \times Y_{\eta}). \quad \square \end{aligned}$$

If  $d(Y_{\eta}) = d(M)$  put  $V_m = \emptyset$ , and if  $V_{\eta i}$  has been defined for all  $i > 0$  put

$$V_{\eta} := (X^{m_{\eta-1}} \times W_{\eta 0}) \cup \bigcup_{i>0} [(id_{X^{m_{\eta-1}}} \times F_{\eta i})^{-1}(V_{\eta i}) \cap (X^{m_{\eta-1}} \times W_{\eta i})].$$

Then inductively  $d(V_{\eta}) < d(X^{m_{\eta}} \times Y_{\eta})$ . In particular  $d(V_{\emptyset}) < d(X^{m+1})$ .

**Lemma 5.** *If  $W \subset X^n$  with  $d(W) < n d(X)$ , then  $\delta(W) < n \delta(X)$ .*

*Proof:* We use induction on  $n$ , the assertion being trivial for  $n = 0, 1$ . So assume it holds for  $n$ , and consider  $W \subseteq X^{n+1}$ . Let  $\pi$  be the projection of  $W$  to the first  $n$  coordinates, and put  $W_i = \{\bar{x} \in \pi(W) : d(\pi^{-1}(\bar{x})) = i\}$  for  $i \leq k$ . Since  $d(W) < d(X^{n+1})$ , we have  $d(W_k) < d(X^n)$ . So by inductive hypothesis

$$\delta(\pi^{-1}(W_k)) \leq \delta(W_k \times X) = \delta(W_k) + \delta(X) < \delta(X^n) + \delta(X) = (n+1) \delta(X).$$

On the other hand, for  $\bar{x} \in W_i$  with  $i < k$  we have

$$\delta(\pi^{-1}(\bar{x})) \leq \alpha d(\pi^{-1}(\bar{x})) = \alpha i$$

by our global inductive hypothesis. Hence by requirement (3)

$$\delta(\pi^{-1}(W_i)) \leq \delta(W_i) + \alpha i \leq \delta(X^n) + \alpha(k-1) < (n+1) \delta(X)$$

since we assume  $\delta(X) \geq \alpha k$ . Thus

$$\delta(W) = \max_{i \leq k} \delta(\pi^{-1}(W_i)) < (n+1) \delta(X). \quad \square$$

It follows that  $\delta(V_{\emptyset}) < \delta(X^{m+1})$ , and

$$(m+1) \delta(X) = \delta(X^{m+1}) = \delta((X^{m_0} \times Y_{\emptyset}) \setminus V_{\emptyset}).$$

For  $\bar{y} \in (X^{m_{\eta i}} \times Y_{\eta i}) \setminus V_{\eta i}$

$$d((id_{X^{m_{\eta-1}}} \times F_{\eta i})^{-1}(\bar{y}) \cap [(X^{m_{\eta-1}} \times W_{\eta i}) \setminus V_{\eta i}]) \leq n_{\eta i} < k,$$

so by inductive hypothesis

$$\delta((id_{X^{m_{\eta-1}}} \times F_{\eta i})^{-1}(\bar{y}) \cap [(X^{m_{\eta-1}} \times W_{\eta i}) \setminus V_{\eta i}]) \leq \alpha n_{\eta i}.$$

Hence

$$\begin{aligned} \delta((id_{X^{m_{\eta-1}}} \times F_{\eta i}) \cap ((X^{m_{\eta-1}} \times W_{\eta i}) \setminus V_{\eta}) \times [(X^{m_{\eta i}} \times Y_{\eta i}) \setminus V_{\eta i}])) \\ \leq \delta((X^{m_{\eta i}} \times Y_{\eta i}) \setminus V_{\eta i}) + \alpha n_{\eta i} \end{aligned}$$

by assumption (3), and

$$\begin{aligned} \delta((id_{X^{m_{\eta-1}}} \times F_{\eta i}) \cap ((X^{m_{\eta-1}} \times W_{\eta i}) \setminus V_{\eta}) \times [(X^{m_{\eta i}} \times Y_{\eta i}) \setminus V_{\eta i}])) \\ = \delta((X^{m_{\eta-1}} \times W_{\eta i}) \setminus V_{\eta}) \end{aligned}$$

by assumption (4). Since  $(X^{m_{\eta}} \times Y_{\eta}) \setminus V_{\eta} = \bigcup_{i>0} (X^{m_{\eta-1}} \times W_{\eta i}) \setminus V_{\eta}$ ,

$$\begin{aligned} \delta((X^{m_{\eta}} \times Y_{\eta}) \setminus V_{\eta}) &= \max_{i>0} \delta((X^{m_{\eta-1}} \times W_{\eta i}) \setminus V_{\eta}) \\ &\leq \max_{i>0} \delta((X^{m_{\eta i}} \times Y_{\eta i}) \setminus V_{\eta i}) + \alpha n_{\eta i}. \end{aligned}$$

On the other hand,  $d(X) + d(Y_{\eta}) = d(Y_{\eta i}) + n_{\eta i}$  for all  $\eta$  and  $i > 0$ . Let  $\eta$  be the branch which corresponds always to the maximum of the  $\delta$ -dimensions. Summing over the initial segments of  $\eta$  we obtain

$$\begin{aligned} (m+1) \delta(X) &= \delta((X^m \times Y_{\emptyset}) \setminus V_{\emptyset}) \leq \delta(X^{m_{\eta}} \times Y_{\eta}) + \alpha \sum_{\emptyset < \eta' \leq \eta} n_{\eta'} \\ &= m_{\eta} \delta(X) + \delta(Y_{\eta}) + \alpha \sum_{\emptyset < \eta' \leq \eta} n_{\eta'} \\ &\leq (m - |\eta|) \delta(X) + \delta(M) + \alpha \sum_{\emptyset < \eta' \leq \eta} n_{\eta'}, \end{aligned}$$

whereas

$$(|\eta| + 1) d(X) = d(Y_{\eta}) + \sum_{\emptyset < \eta' \leq \eta} n_{\eta'} = d(M) + \sum_{\emptyset < \eta' \leq \eta} n_{\eta'}.$$

Therefore

$$(|\eta| + 1) \delta(X) \leq \alpha (d(M) + \sum_{\emptyset < \eta' \leq \eta} n_{\eta'}) = \alpha (|\eta| + 1) d(X),$$

and  $\delta(X) \leq \alpha d(X)$ . This proves the theorem.  $\square$

We shall now give a second, type-based proof for Theorem 2.

*Proof:* For two partial types  $\pi, \pi'$  we put  $\delta(\pi) \leq \delta(\pi')$  if for every  $X' \in \pi'$  there is  $X \in \pi$  with  $\delta(X) \leq \delta(X')$ . Note that  $\leq$  is transitive.

**Claim.** *It is enough to prove the assertion for complete types.*

*Proof of Claim:* Let  $X$  be an  $A$ -definable set, and  $\mathfrak{X}$  the collection of  $A$ -definable  $X' \subseteq X$  such that  $d(M)\delta(X') \leq d(X')\delta(M)$ . Then  $\mathfrak{X}$  is closed under finite unions, so either  $d(M)\delta(X) \leq d(X)\delta(M)$ , or there is a type  $p \in S(A)$  completing the partial type  $\{X \setminus X' : X' \in \mathfrak{X}\}$ . By

assumption  $d(M)\delta(p) \leq d(p)\delta(M)$ . So there are  $A$ -definable  $X_1, X_2 \in p$  with  $d(M)\delta(X_1) \leq d(p)\delta(M)$  and  $d(X_2) = d(p)$ . But then  $X_1 \cap X_2 \in \mathfrak{X}$ , a contradiction.  $\square$

So let  $p \in S_1(A)$ . We shall use induction on  $d(p) =: k$ . Clearly we may assume that  $d(M)\delta(p) \geq d(p)\delta(M)$ . For ease of notation we also assume that the value group  $G$  of  $\delta$  is divisible.

**Claim.** *If  $p' \in S_1(A)$ , there is  $q \in S_2(A)$  extending  $p \otimes_A p'$  with  $\delta(p \times p') \leq \delta(q)$ .*

*Proof of Claim:* Suppose not, and consider

$$\mathfrak{X} := \{X \subseteq M^2 \text{ } A\text{-definable} : \delta(p \times p') \not\leq \delta((p \times p') \cup \{X\})\}.$$

Then  $\mathfrak{X}$  is closed under unions, and we can put  $\rho := (p \times p') \cup \{\neg X : X \in \mathfrak{X}\}$ , a consistent partial type. By assumption  $d(\rho) < d(p) + d(p')$ , so the projection to the first coordinate has fibres of dimension  $i < k$ . So there are definable sets  $X \in p$ ,  $X' \in p'$  and  $X \times X' \supset Y \in \rho$  with  $d(X) = d(p)$ ,  $d(X') = d(p')$ ,  $d(Y) = d(\rho)$  and  $d(Y \cap (X \times \{x'\})) = i$  for all  $x' \in X'$ ; we may assume in addition that  $\delta(X') + \frac{i}{k}\delta(p) < \delta(p \times p')$ . By inductive hypothesis  $d(M)\delta(Y \cap (X \times \{x'\})) \leq i\delta(M)$  for all  $x' \in X'$ , so

$$\delta(Y) \leq \delta(X') + i\delta(M)/d(M) \leq \delta(X') + i\delta(p)/d(p) < \delta(p \times p'),$$

a contradiction to the definition of  $\rho$ .  $\square$

By  $\mathcal{F}$ -minimality there is  $n < \omega$ , a sequence  $p = p_0, p_1, \dots, p_n$  of complete types over  $A$ , a complete  $A$ -type  $q_i \supseteq p \otimes_A p_i$  with  $\delta(p) + \delta(p_i) \leq \delta(q_i)$  for  $i < n$ , and correspondences  $(F_i : i < n)$  in  $\mathcal{F}$ , such that  $p_{i+1}$  is a completion of  $F_i(q_i)$  for all  $i < n$  with  $d(p_i) < d(p_{i+1})$ , and  $d(p_n) = d(M)$ . For  $i < n$  put  $R_i := F_i \cap (q_i \times p_{i+1})$ , and choose  $A$ -definable sets  $X \in q_i$ ,  $X' \in p_{i+1}$  and  $Y \in R_i$  with  $d(X) = d(q_i) = d(p) + d(p_i)$ ,  $d(X') = d(p_{i+1})$ ,  $Y \subseteq X \times X'$ , and such that the fibres of the projection  $\pi$  of  $Y$  to the first two coordinates have constant dimension  $j_i = d(\pi^{-1}(a))$ , where  $a \models p_{i+1}$ . Then

$$i_j = d(X) - d(X') = d(p) + d(p_i) - d(p_{i+1}) < d(p)$$

by axiom (3a). By inductive hypothesis  $\delta(\pi^{-1}(a)) \leq i_j \delta(M)/d(M)$  for all  $a \in X'$ , whence  $\delta(Y) \leq \delta(X') + i_j \delta(M)/d(M)$ . Letting  $X'$  converge to  $p_{i+1}$  and  $Y$  to  $R_i$ , we obtain  $\delta(R_i) \leq \delta(p_{i+1}) + i_j \delta(M)/d(M)$ .

Since condition (4) implies  $\delta(q_i) \leq \delta(F_i \cap (q_i \times p_{i+1})) = \delta(R_i)$ , we get

$$\delta(p) + \delta(p_i) \leq \delta(q_i) \leq \delta(R_i) \leq \delta(p_{i+1}) + i_j \delta(M)/d(M).$$



By induction we obtain for  $m < n$

$$m \delta(p) \leq \delta(p_m) + \frac{\delta(M)}{d(M)} \sum_{j < m} i_j.$$

In particular

$$n \delta(p) \leq \delta(M) + \frac{\delta(M)}{d(M)} \sum_{j < n} i_j.$$

On the other hand,

$$d(M) + \sum_{j < n} i_j = d(p_n) + \sum_{j < n} [d(p) + d(p_i) - d(p_{i+1})] = n d(p),$$

whence

$$n \delta(p) \leq \frac{\delta(M)}{d(M)} [d(M) + \sum_{j < n} i_j] = \frac{\delta(M)}{d(M)} n d(p),$$

which proves the theorem.  $\square$

**Remark 2.** The above proof of Theorem 2 defined the relation  $\delta(\pi) \leq \delta(\pi')$  without actually defining the quantities  $\delta(\pi)$ . Perhaps for other applications an invariant  $\delta(\pi)$  for types may be useful. We sketch now how this may be done.

Definition 4 requires  $\delta$  to be a function into the non-negative elements of a linearly ordered group  $G$  that can be assumed divisible. In place of this, let us gain generality by taking  $G = (G, +, 0, <)$  to be a divisible linearly ordered commutative semi-group. This means that (1)–(2) below hold; we may as well assume (3); we assume cancellation only in the limited form (4), with respect to a distinguished element  $\delta(M)$ .

- (1)  $(G, +, 0)$  is an additive semi-group, with every element uniquely divisible by any positive integer.
- (2)  $<$  is a linear ordering, and  $x \leq y$  implies  $x + z \leq y + z$ .
- (3) For any  $x \in G$  there is  $k < \omega$  with  $0 \leq x \leq k \delta(M)$ .
- (4)  $x + \delta(M) > x$  for any  $x$ .

It follows that  $x + \frac{1}{n} \delta(M) > x$  for any  $x$  and integer  $n > 0$ .

These more general assumptions have the advantage that the semi-group  $G$  can be completed by means of Dedekind cuts. The assumptions continue to hold; in particular (4) does, since if  $U$  is a Dedekind cut invariant under adding  $\delta(M)$ , then by (3) it must include all of  $\Gamma$ , but Dedekind cuts are assumed bounded.

Now for any partial type  $\pi = \bigwedge_{i \in I} X_i$  we can define  $\delta(\pi) = \inf_{i \in I} \delta(X_i)$ . The earlier definition of the inequality is now a consequence. Whether the greater generality has any additional use, we do not know.

**Corollary 6.** *Under the same hypotheses as Theorem 2, let  $X \subset M^n$  be definable. Then  $d(M) \delta(X) \leq d(X) \delta(M)$ .*

*Proof:* We use induction on  $n$ , the assertion being Theorem 2 for  $n = 1$ . For  $X \subseteq M^{n+1}$  let  $\pi$  be the projection to the first  $n$  coordinates, and partition  $Y := \pi(X)$  into sets

$$Y_i := \{\bar{x} \in Y : d(\pi^{-1}(\bar{x}) \cap X) = i\}.$$

Let  $X_i := \pi^{-1}(Y_i) \cap X$ , then  $(X_i : i \leq d(M))$  partitions  $X$ , and

$$d(X) = \max_{i \leq d(M)} d(X_i) = \max_{i \leq d(M)} d(Y_i) + i.$$

For every  $i \leq d(M)$  and  $\bar{x} \in Y_i$  Theorem 2 yields  $\delta(\pi^{-1}(\bar{x}) \cap X) \leq \alpha i$ , with  $\alpha = \delta(M)/d(M)$ . By inductive hypothesis  $\delta(Y_i) \leq \alpha d(Y_i)$ , so

$$\delta(X) = \max_{i \leq d(M)} \delta(X_i) \leq \max_{i \leq d(M)} \delta(Y_i) + \alpha i \leq \alpha \max_{i \leq d(M)} d(Y_i) + i = \alpha d(X).$$

□

**Remark 3.** If  $\mathfrak{M}$  is  $\mathcal{F}$ -minimal, then  $\mathfrak{M}^n$  can be shown to be minimal with respect to the induced set of correspondences; this yields an alternative proof of Corollary 6.

## 2. AN EXAMPLE THAT COUNTS

Let  $(\mathfrak{M}_n : n < \omega)$  be a family of  $\mathcal{L}$ -structures for some language  $\mathcal{L}$ , and  $\Gamma_n$  finite subsets of  $M_n$ . For some ultrafilter on  $\omega$  let  $\langle \mathfrak{M}, \Gamma \rangle$  be the ultraproduct of the structures  $\langle \mathfrak{M}_n, \Gamma_n \rangle$ . The ultraproduct of the counting measures on the  $\Gamma_n$  yields a finitely additive measure  $\mu$  on the definable subsets of  $\Gamma$  which takes values in some non-standard real closed field  $\mathbb{R}^*$ . Note that  $\langle \mathfrak{M}, \Gamma, \mathbb{R}^*, \mu, \log \rangle$  is  $\aleph_0$ -saturated (in fact, even  $\aleph_1$ -saturated).

Let  $I$  be the convex hull of  $\mathbb{Z}$  in  $\mathbb{R}^*$ , and  $\pi : \mathbb{R}^* \rightarrow \mathbb{R}^*/I$  the natural (additive) quotient map. Define

$$\delta(X) = \pi \log \mu(X),$$

and note that  $\delta(X) = 0$  if and only if  $\log \mu(X) \in I$ , that is  $\mu(X) \in I$ , in other words  $\mu_n(X_n) = O(1)$  in the factors, that is  $X$  is finite in the ultraproduct. For a definable subset of  $M$  we put  $\delta(M) := \delta(M \cap \Gamma)$ .

**Lemma 7.** *This  $\delta$  satisfies conditions (0)–(4) from Theorem 2, provided  $d(X) = 0$  implies  $X$  finite, and  $\Gamma$  is closed under the correspondences (i.e. for all  $xx' \in \Gamma^2$  and  $y \in M$  such that  $F(xx'y)$  holds,  $y \in \Gamma$  as well).*

*Proof:* (1) is obvious. For (2) note that

$$\mu(X \cup Y) \leq \mu(X) + \mu(Y) \leq 2 \max\{\mu(X), \mu(Y)\},$$

whence  $\log(\mu(X \cup Y)) \leq \log 2 + \max\{\log \mu(X), \log \mu(Y)\}$ . Since  $\log 2 \in I$ , we get  $\delta(X \cup Y) \leq \max\{\delta(X), \delta(Y)\}$ ; the other inequality follows from monotonicity.

We claim that for any definable map  $f : X \rightarrow Y$ , if  $\delta(f^{-1}(y)) \leq \alpha$  for all  $y \in Y$ , then there is  $r \in \mathbb{R}^*$  with  $\pi(r) = \alpha$  and  $\log \mu(f^{-1}(y)) \leq r$  for all  $y \in Y$ . Indeed, pick any  $r_0 \in \mathbb{R}^*$  with  $\pi(r_0) = \alpha$ . Put

$$Y_n := \{y \in Y : \log \mu(f^{-1}(y)) \leq r_0 + n\}.$$

Then  $Y_n \subset Y_{n+1}$  for all  $n < \omega$ , and  $Y = \bigcup_{n < \omega} Y_n$ ; by  $\aleph_0$ -saturation there is  $n_0$  with  $Y = Y_{n_0}$ . Then  $r := r_0 + n_0$  will do.

This shows (3). Finally, (4) is clear, since the fibres of the projection of any  $F \in \mathcal{F}$  to the first two coordinates must have  $d$ -dimension zero, hence be finite in the ultraproduct, and thus uniformly finite in the factors; they are non-empty by closedness of  $\Gamma$  under  $\mathcal{F}$ .  $\square$

Unwinding the definitions, for this choice of  $\delta$  (and suitable dimension  $d$ ) the inequality  $d(M)\delta(X) \leq \delta(M)d(X)$  becomes

$$|X_n \cap \Gamma_n| \leq O(|\Gamma_n|^{d(X)/d(M)}).$$

Possible choices for  $d$  include algebraic dimension, Morley rank, Shelah rank, Lascar rank, SU-rank or  $S_1$ -rank, whenever it is finite, additive and definable in the pure  $\mathcal{L}$ -structure  $\mathfrak{M}$ .

**Remark 4.** Uniformity in parameters of the constant intervening in the  $O$ -notation follows automatically from compactness.

**Remark 5.** Note that for any definable map  $f : X \rightarrow Y$ :

- (1) If  $\delta(f^{-1}(y)) \geq \alpha$  for all  $y \in Y$ , then  $\delta(Y) + \alpha \leq \delta(X)$ .
- (2) If  $\delta(f^{-1}(y)) \leq \alpha$  for all  $y \in Y$  and  $f(X \cap \Gamma) \subseteq \Gamma$ , then  $\delta(X) \leq \delta(Y) + \alpha$ .

In particular  $\delta$  is invariant under definable bijections  $f$  preserving  $\Gamma$  (i.e.  $x \in \Gamma$  if and only if  $f(x) \in \Gamma$ ).

## 3. AN APPLICATION

We shall now give the model-theoretic formulation of the theorem by Larsen and Pink alluded to in the introduction.

**Theorem 8.** [1] *Let  $G_n$  be a simple algebraic group varying in an algebraic family and  $\Gamma_n$  a finite subgroup such that in the ultraproduct  $G$  the subgroup  $\Gamma$  is Zariski-dense. Then for any subvariety  $V$  of  $G$*

$$|V_n \cap \Gamma_n| \leq O(|\Gamma_n|^{\dim(V)/\dim(G)}).$$

*Proof:* Since  $G_n$  varies in an algebraic family,  $G$  is a simple algebraic group, and  $d = \dim = \text{RM}$  is finite, additive and definable. Let  $\mathcal{F}$  be the collection of maps  $F_c(x, y) = cx^{-1}c^{-1}y$ , where  $c$  runs over a countable Zariski-dense subgroup  $\Gamma_0$  of  $\Gamma$ . Clearly  $\Gamma$  is  $\mathcal{F}$ -closed; moreover  $G$  is  $\mathcal{F}$ -minimal by Example 1. Theorem 2 and Lemma 7 yield the result.  $\square$

**Corollary 9.** [1] *In the setting of Theorem 8 consider  $a \in \Gamma$  with  $\text{RM}(C_G(a)) > 0$ ,  $\text{RM}(a^G) > 0$  and  $\delta(G) > 0$ . Then  $\Gamma$  meets both  $C_G(a)$  and  $a^G$  in infinite sets.*

*Proof:* Using the definable map  $x \mapsto a^x$  and translation maps between  $C_G(a)$  and its cosets, we see that

$$\begin{aligned} \text{RM}(C_G(a)) + \text{RM}(a^G) &= \text{RM}(G), \text{ and} \\ \delta(C_G(a)) + \delta(a^G) &= \delta(G). \end{aligned}$$

If  $\alpha = \delta(G)/\text{RM}(G)$ , then  $\delta(C_G(a)) \leq \alpha \text{RM}(C_G(a))$  and  $\delta(a^G) \leq \alpha \text{RM}(a^G)$  by Theorem 8, so equality must hold.  $\square$

## 4. BAD FIELDS

A *bad field* [2] is a structure  $\langle K, 0, 1, +, -, \cdot, T \rangle$  of finite Morley rank, where  $T$  is a predicate for a distinguished infinite proper connected multiplicative subgroup (or even a non-algebraic connected subgroup of  $(K^\times)^n$  for some  $n$ , but these shall not be considered here). Such an object appears naturally when considering a faithful action of an abelian group  $M$  on an  $M$ -minimal abelian group  $A$ , the whole of finite Morley rank: We obtain that there is an algebraically closed field  $K$  such that  $A \cong K^+$  and  $M \hookrightarrow K^\times$ ; one knows that the image of  $M$  generates  $K$  additively, but *a priori* it could be a proper subgroup. In particular, the possible existence of bad fields (and of bad groups) prevents us from proving an analogue of the Feit-Thompson theorem

for simple groups of finite Morley rank, namely that they contain an involution (or, indeed, any torsion element at all).

In [3] the second author showed that under the assumption that there are infinitely many prime numbers of the form  $(p^n - 1)/(p - 1)$  (called *p-Mersenne primes*), there is no bad field of characteristic  $p > 0$ . In [4] he obtained an asymptotic estimate for the number of  $\mathbb{F}_q$ -rational points of a multiplicative subgroup of rank 1; this shows the nonexistence of bad fields with  $\text{RM}(T)$  of rank 1 modulo a slightly weaker number-theoretic hypothesis. We can now obtain an analogous asymptotic estimate for multiplicative subgroups of arbitrary rank.

For two functions  $f$  and  $g$  on  $\mathbb{N}$  we put  $f \asymp g$  if there are positive constants  $c, c'$  with  $cf(n) \leq g(n) \leq c'f(n)$  for all  $n \in \mathbb{N}$ .

**Theorem 10.** *For any definable subset  $X$  of a bad field  $K$  of positive characteristic and any finite subfield  $\mathbb{F}_q \leq K$  we have  $|X \cap \mathbb{F}_q| \leq O(q^{\text{RM}(X)/\text{RM}(K)})$ . In particular  $|T \cap \mathbb{F}_{p^n}| \asymp p^{n \text{RM}(T)/\text{RM}(K)}$ .*

*Proof:* Let  $\langle K, T \rangle$  be a bad field of characteristic  $p > 0$ . We put  $\mathfrak{M}_n = \langle K, T \rangle$  for all  $n < \omega$ , and  $\Gamma_n = \mathbb{F}_{p^n}$ ; our correspondences  $\mathcal{F}$  will be addition and multiplication. Clearly  $\Gamma$  is closed under  $\mathcal{F}$ , and  $K$  is  $\mathcal{F}$ -minimal by Example 2. So Theorem 2 and Lemma 7 imply the first assertion.

By [3, Theorem 2] there is an  $\emptyset$ -definable partial function  $f : K \rightarrow T$  with generic domain and an integer  $\ell > 0$  such that  $f(ta) = t^\ell f(a)$  for all  $a \in \text{dom}(f)$  and all  $t \in T$  (in particular  $\text{dom}(f)$  is closed under multiplication by  $T$ ). Since  $T$  is  $\ell$ -divisible, all fibres have the same rank, namely  $\text{RM}(K) - \text{RM}(T)$ . Hence the number of  $\mathbb{F}_q$ -points on a fibre is bounded by  $O(q^{1-\alpha})$ , where  $\alpha = \text{RM}(T)/\text{RM}(K)$ . Moreover, the complement of the domain has rank at most  $\text{RM}(K) - 1$ , so its number of  $\mathbb{F}_q$ -points is bounded by  $O(q^{1-1/\text{RM}(K)})$ . Since  $\mathbb{F}_q$  is precisely the set of fixed points of the definable automorphism  $x \mapsto x^q$ , it is closed under all  $\mathbb{F}_q$ -definable functions. Hence the number of  $\mathbb{F}_q$ -points of  $T$  is at least  $(q - O(q^{1-1/\text{RM}(K)}))/O(q^{1-\alpha}) \geq cq^\alpha$  for some constant  $c$ .  $\square$

**Definition 5.** Let  $\pi$  be a set of prime numbers. For an integer  $n$  the  $\pi$ -part  $n_\pi$  is the biggest  $\pi$ -number (with all prime divisors in  $\pi$ ) dividing  $n$ .

**Corollary 11.** *Suppose  $\langle K, T \rangle$  is a bad field of characteristic  $p > 0$ , and let  $\pi$  be the set of prime orders of elements in  $T$ . Then*

$$(p^n - 1)_\pi \asymp p^{\alpha n},$$

with  $\alpha = \text{RM}(T)/\text{RM}(K)$ .

*Proof:* Since  $T$  is divisible, it is a direct sum of Prüfer groups. Hence if  $k$  is the subfield of  $K$  with  $p^n$  elements and  $q$  is a prime dividing  $|T \cap k^\times|$ , then  $T$  contains all of the  $q$ -part of  $k^\times$ . Thus  $|T \cap k| = (p^n - 1)_\pi$ .  $\square$

**Definition 6.** Let  $0 < \alpha < 1$ . A set  $\pi$  of primes is  $(p, \alpha)$ -balanced if  $((p^n - 1)_\pi) \asymp p^{\alpha n}$ . It is  $p$ -balanced if it is  $(p, \alpha)$ -balanced for some  $\alpha$  with  $0 < \alpha < 1$ .

Note that if  $\pi$  is  $(p, \alpha)$ -balanced, then the complement of  $\pi$  is  $(p, 1 - \alpha)$ -balanced.

**Corollary 12.** *If there is no  $p$ -balanced set, then there is no bad field of characteristic  $p$ .*

*Proof:* This follows immediately from Corollary 11.  $\square$

## REFERENCES

- [1] Michael J. Larsen and Richard Pink. Finite subgroups of algebraic groups. Preprint 1998.
- [2] Bruno Poizat. *Groupes Stables*. Nur al-Mantiq wal-Marifah, Villeurbanne, 1987. Translated as: *Stable Groups*, AMS, 2002.
- [3] Frank O. Wagner. Bad fields of positive characteristic. *Bulletin of the London Mathematical Society*, 35:499-502, 2003.
- [4] Olivier Roche and Frank O. Wagner. Bad fields with a torus of rank 1. *Proceedings of the Euro-conference on Model Theory and Applications, Ravello 2002*, Quaderni di Matematica, Seconda Università di Napoli, Caserta, 2005.

EHUD HRUSHOVSKI, EINSTEIN INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, JERUSALEM 91904, ISRAEL

*E-mail address:* ehud@math.huji.ac.il

FRANK O. WAGNER, INSTITUT CAMILLE JORDAN, UNIVERSITÉ CLAUDE BERNARD (LYON-1), 43 BOULEVARD DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CÉDEX, FRANCE

*E-mail address:* wagner@math.univ-lyon1.fr