

# Injectivity Radius of Lorentzian Manifolds

Bing-Long Chen\* and Philippe G. LeFloch†

December 29, 2006

## Abstract

Motivated by the application to spacetimes of general relativity we investigate the geometry and regularity of Lorentzian manifolds under certain curvature and volume bounds. We establish several injectivity radius estimates at a point or on the past null cone of a point. Our estimates are entirely local and geometric, and are formulated via a reference Riemannian metric that we canonically associate with a given observer  $(p, T)$  –where  $p$  is a point of the manifold and  $T$  is a future-oriented time-like unit vector prescribed at  $p$ . The proofs are based on a generalization of arguments from Riemannian geometry. We first establish estimates on the reference Riemannian metric, and then express them in term of the Lorentzian metric. In the context of general relativity, our estimates should be useful to investigate the regularity of spacetimes satisfying Einstein field equations.

## 1 Introduction

### Aims of this paper

The regularity and compactness of Riemannian manifolds under a priori bounds on geometric quantities such as curvature, volume, or diameter represent important issues in Riemannian geometry. In particular, the derivation of lower bounds on the injectivity radius of a Riemannian manifold, and the construction of local coordinate charts in which the metric has optimal regularity are now well-understood. Moreover, Cheeger-Gromov's theory provides geometric conditions for the strong compactness of sequences of manifolds and has become a central tool in Riemannian geometry. See for instance [1, 4, 5, 7, 8, 9, 16, 21, 22].

---

\*Department of Mathematics, Sun Yet Sen University, 510275 Guang-Zhou, P.R. of China. E-mail: [mcsobl@mail.sysu.edu.cn](mailto:mcsobl@mail.sysu.edu.cn).

†Laboratoire Jacques-Louis Lions & Centre National de la Recherche Scientifique, Université de Paris 6, 4 place Jussieu, 75252 Paris, France. E-mail : [lefloch@ann.jussieu.fr](mailto:lefloch@ann.jussieu.fr)  
2000 *AMS Subject Classification*: 53C50, 83C05, 53C12, 53C22.

*Key Words*: Lorentzian geometry, Riemannian geometry, general relativity, Einstein equation, injectivity radius, Jacobi field, null cone, comparison geometry.

Our objective in the present paper is to present some extension of these classical techniques and results to Lorentzian manifolds. Recall that a Lorentzian metric is not positive definite, but has signature  $(-, +, \dots, +)$ . Motivated by recent work by Anderson [2] and Klainerman and Rodnianski [19], we derive here several injectivity radius estimates for Lorentzian manifolds satisfying certain curvature and volume bounds. That is, we provide sharp lower bounds on the size of the geodesic ball around one point within which the exponential map is a global diffeomorphism and, therefore, we obtain sharp control of the manifold geometry. Our proofs rely on arguments that are known to be flexible and efficient in Riemannian geometry, and are here extended to the Lorentzian setting; we analyze the properties of Jacobi fields and rely on volume comparison and homotopy arguments.

In our presentation (see for instance our main result in Theorem 1.1 at the end of this introduction) we emphasize the importance of having assumptions and estimates that are stated locally and geometrically, and avoid direct use of coordinates. When necessary, coordinates should be constructed a posteriori, once uniform bounds on the injectivity radius have been established.

Our motivation comes from general relativity, where one of the most challenging problems is the formation and the structure of singularities in solutions to the Einstein field equations. Relating curvature and volume bounds to the regularity of the manifold, as we do in this paper, is necessary before tackling an investigation of the geometric properties of singular spacetimes satisfying Einstein equations. (See, for instance, [2, 3] for some background on this subject.)

Two preliminary observations should be made. First, since the Lorentzian norm of a non-zero tensor may vanish it is clear that only limited information would be gained from an assumption on the Lorentzian norm of the curvature tensor. This justifies that we endow the Lorentzian manifold with a “reference” Riemannian metric (denoted by  $g_T$  below); this metric is defined at a point  $p$  once we prescribe a future-oriented time-like unit vector  $T$ . We refer to the pair  $(p, T)$  as an observer located at the point  $p$ . This reference vector is necessary in order to define appropriate notions of conjugate and injectivity radii. (See Section 2 below for details.)

Secondly, we rely here on the elementary but essential observation that, in the flat Riemannian and Lorentzian spaces, geodesics (are straight lines and therefore) coincide. Under our assumptions, we will see that geodesics associated with the given Lorentzian metric are comparable to geodesics associated with the reference Riemannian metric. On the other hand, the curvature bound assumed on the Lorentzian metric implies, in general, no information on the curvature of the reference metric. As we show below, one of the main issues is to guarantee the regularity of a foliation of the manifold by spacelike hypersurfaces.

## Earlier work

Let us briefly review some classical results from Riemannian geometry. Let  $(M, g)$  be a differentiable  $n$ -manifold (possibly with boundary) endowed with a Riemannian metric  $g$ . (Throughout the present paper, the manifolds and metrics are always assumed to be smooth.) Denote by  $\mathcal{B}(p, r)$  the corresponding geodesic ball centered at  $p \in M$  and with radius  $r > 0$ . Suppose that at some point  $p \in M$  the unit ball  $\mathcal{B}(p, 1)$  is compactly contained in  $M$  and that the Riemann curvature bound and the lower volume bound,

$$\|\mathbf{Rm}_g\|_{\mathbf{L}^\infty(\mathcal{B}(p,1))} \leq K, \quad \mathbf{Vol}_g(\mathcal{B}(p,1)) \geq v_0, \quad (1.1)$$

hold for some constants  $K, v_0 > 0$ . (We use the standard notation  $\mathbf{L}^m$ ,  $1 \leq m \leq \infty$ , for the spaces of Lebesgue measurable functions.) Then, according to Cheeger, Gromov, and Taylor [10] the injectivity radius  $\mathbf{Inj}_g(M, p)$  at the point  $p$  is bounded below by a positive constant  $i_1 = i_1(K, v_0, n)$ ,

$$\mathbf{Inj}_g(M, p) \geq i_1. \quad (1.2)$$

It should be noticed that this is a local statement; for earlier work on the injectivity radius see [5, 11, 15].

Moreover, Jost and Karcher [16] relied on the regularity theory for elliptic operators and established the existence of coordinates in which the metric has optimal regularity and are defined in a ball with radius  $i_2 = i_2(K, v_0, n)$ . Precisely, given  $\varepsilon > 0$  and  $0 < \gamma < 1$  there exist a positive constant  $C(\varepsilon, \gamma)$  (depending also upon  $(K, v_0, n)$ ) and a system of harmonic coordinates defined in the geodesic ball  $\mathcal{B}(p, i_2)$  in which the metric  $g$  is close to the Euclidian metric  $g_E$  in these coordinates and has optimal regularity, in the following sense:

$$\begin{aligned} e^{-\varepsilon} g_E &\leq g \leq e^\varepsilon g_E, \\ r^{1+\gamma} \|\partial g\|_{\mathbf{C}^\gamma(\mathcal{B}(p,r))} &\leq C(\varepsilon, \gamma), \quad r \in (0, i_2]. \end{aligned} \quad (1.3)$$

Here,  $\mathbf{C}^0$  and  $\mathbf{C}^\gamma$  are the spaces of continuous and Hölder continuous functions, respectively. Harmonic coordinates are optimal [12] in the sense that if the metric is of class  $\mathbf{C}^{k,\gamma}$  in certain coordinates then it has at least the same regularity in harmonic coordinates.

The above results were later generalized by Anderson [1] and Petersen [22] who replaced the  $\mathbf{L}^\infty$  curvature bound by an  $\mathbf{L}^m$  curvature bound with  $m > n/2$ . For instance, one can take  $m = 2$  in dimension  $n = 3$  in the application to general relativity (since time-slices of Lorentzian 4-manifolds are Riemannian 3-manifolds).

It is only more recently that the same questions were tackled for Lorentzian  $(n + 1)$ -manifolds  $(M, g)$ . Anderson [2, 3] studied the long-time evolution of solutions to the Einstein field equations and formulate several conjecture. In particular, assuming the Riemann curvature bound in some domain  $\Omega$

$$\|\mathbf{Rm}_g\|_{\mathbf{L}^\infty(\Omega)} \leq K \quad (1.4)$$

and other regularity conditions, he investigated the existence of coordinates that are harmonic in each spacelike slice of a time foliation of  $M$ . This work by Anderson motivated our investigation in the present paper.

On the other hand, motivated by applications to general relativity and nonlinear wave equations using harmonic analysis tools, Klainerman and Rodnianski [19] considered asymptotically flat spacetimes endowed with a time foliation and satisfying the  $L^2$  curvature bound

$$\|\mathbf{Rm}_g\|_{L^2(\Sigma)} \leq K \tag{1.5}$$

for every spacelike hypersurface  $\Sigma$ . To control the injectivity radius of past null cones, they relied on their earlier work [17, 18] on the conjugate radius of null cones in terms of Bell-Robinson's energy and energy flux, and derived in [19] a new estimate for the null cut locus radius. We refer to these papers for further details and references on the Einstein equations. Section 6 of the present paper is a prolongation of the work [19].

## Outline of this paper

The present paper establishes four estimates on the radius of injectivity of Lorentzian manifolds, which hold either in a neighborhood of a point or on the past null cone at a point. Our assumptions are formulated within a geodesic ball (or within a null cone) and possibly apply in a ball with arbitrary size as long as our curvature and volume assumptions hold. All assumptions and statements are local and geometric.

An outline of the paper is as follows. In Section 2, we begin with basic material from Lorentzian geometry and we introduce the notions of reference metric and exponential map for Lorentzian manifolds. In Section 3, we state our first estimate (Theorem 3.1 below) for a class of manifolds that have bounded curvature and admit a time foliation by slices with bounded extrinsic curvature. In Section 4, we provide a proof of this first estimate and we introduce a technique that will be used (in variants) throughout this paper; we combine two main ingredients : sharp estimates for Jacobi fields along geodesics, and an homotopy argument based on contracting a possible loop to two linear segments. In Section 5, our second main result (Theorem 5.1) shows, under the same assumptions, the existence of convex functions (distance functions) and convex neighborhoods; this result leads us to a lower bound of the convexity radius.

In Section 6, our third estimate (Theorem 6.1) covers the case of null cones under the assumption that the manifold has  $L^2$  bounded curvature on every spacelike slice; this provides a generalization and an alternative proof to the result by Klainerman and Rodnianski in [19].

Next, in Section 7, we establish our principal and fourth result (stated in Theorem 1.1 below) which provides an injectivity radius bound under the mild assumption that the exponential map  $\exp_p$  is defined in some ball and the

curvature  $\mathbf{Rm}$  is bounded. Most importantly, this is a general result that does not require a time foliation of the manifold but solely a single reference (future-oriented time-like unit) vector  $T$  at the base point  $p$ . This is very natural in the context of general relativity and  $(p, T)$  is interpreted as an observer at the point  $p$ .

Given an observer  $(p, T)$ , we can define the ball  $B_T(0, r) \subset T_p M$  with radius  $r$ , determined by the reference Riemannian inner product at  $p$ , and we can also define the geodesic ball  $\mathcal{B}_T(p, r) := \exp_p(B_T(0, r))$ . In turn, the radius of injectivity  $\mathbf{Inj}_g(M, p, T)$  is defined as the largest radius  $r$  such that the exponential map is a diffeomorphism from  $B_T(0, r)$  onto  $\mathcal{B}_T(p, r)$ . Let us then consider an arbitrary geodesic  $\gamma = \gamma(s)$  initiating at  $p$  and let us  $g$ -parallel transport the vector  $T$  along this geodesic, defining therefore a vector field  $T_\gamma$  along this geodesic, only. At every point of  $\gamma$  we introduce the reference metric  $g_{T_\gamma}$  and compute the curvature norm  $|\mathbf{Rm}_g|_{T_\gamma}$ . This allows us to express the curvature bound. For the convenience of the reader we state here our main result and refer to Section 7 for further details.

**Theorem 1.1** (Injectivity radius of Lorentzian manifolds). *Let  $M$  be a time-orientable Lorentzian, differentiable  $(n + 1)$ -manifold. Consider an observer  $(p, T)$  consisting of a point  $p \in M$  and a reference (future-oriented time-like unit) vector  $T \in T_p M$ . Assume that the exponential map  $\exp_p$  is defined in the ball  $B_T(0, r_0) \subset T_p M$  and the Riemann curvature satisfies*

$$\sup_{\gamma} |\mathbf{Rm}_g|_{T_\gamma} \leq \frac{1}{r_0^2}, \quad (1.6)$$

where the supremum is over the domain of definition of  $\gamma$  and over every  $g$ -geodesic  $\gamma$  initiating from a vector in the Riemannian ball  $B_T(0, r_0) \subset T_p M$ . Then, there exists a constant  $c(n)$  depending only on the dimension  $n$  such that

$$\frac{\mathbf{Inj}_g(M, p, T)}{r_0} \geq c(n) \frac{\mathbf{Vol}_g(\mathcal{B}_T(p, c(n)r_0))}{r_0^{n+1}}. \quad (1.7)$$

This result should be compared with the injectivity radius estimate established by Cheeger, Gromov, and Taylor [10] in Riemannian geometry. Observe that the curvature assumption (1.6) can always be satisfied by suitably rescaling the metric tensor. It would be interesting to replace the volume term in the right-hand side of (1.7) by  $\mathbf{Vol}_g(\mathcal{B}(p, r_0))$ .

Finally, in the last two sections of this paper, we establish a volume comparison theorem for future cones and generalize our main theorem to the volume of a future cone (Section 8), and we briefly discuss the regularity of the Lorentzian metric in harmonic-like coordinates, and present a generalization to pseudo-Riemannian manifolds (Section 9).

## 2 Preliminaries on Lorentzian geometry

### Basic definitions

It is useful to discuss first some basic definitions from Lorentzian geometry, for which we can refer to the textbook by Penrose [20]. Throughout this paper,  $(M, g)$  is a connected and differentiable  $(n + 1)$ -manifold, endowed with a Lorentzian metric tensor  $g$  with signature  $(-, +, \dots, +)$ . To emphasize the role of the metric  $g$  or the point  $p$  we use any of the following notations

$$g_p(X, Y) = \langle X, Y \rangle_{g_p} = \langle X, Y \rangle_g = \langle X, Y \rangle_p$$

for the inner product of two vectors  $X, Y$  at a point  $p \in M$ ; we sometimes also write  $|X|_{g_p}^2$  instead of  $g_p(X, X)$ . Recall that the tangent vectors  $X \in T_p M$  are called time-like, null, or spacelike depending whether the norm  $g_p(X, X)$  is negative, zero, or positive, respectively. Vectors that are time-like or null are called causal.

The time-like vectors form a cone with two connected components. The manifold  $(M, g)$  is said to be time-orientable if we can select in a continuous way a half-cone of time-like vectors at every point  $p$ . The choice of a specific orientation allows us to decompose the cone of time-like vectors into future-oriented and past-oriented ones. The set of all future-oriented, time-like vectors at  $p$  and the corresponding bundle on  $M$  are denoted by  $T_p^+ M$  and  $T^+ M$ , respectively. We also introduce the bundle  $T_1^+ M$  consisting of elements of  $T^+ M$  with unit length.

By definition, a trip is a continuous curve  $\gamma : (a, b) \rightarrow M$  made of finitely many future-oriented, time-like geodesics. We write  $p \ll q$  if there exists a trip from  $p$  to  $q$ . A causal trip is defined similarly except that the geodesics may be causal instead of time-like, and we write  $p < q$  if there exists a causal trip from  $p$  to  $q$ .

The set  $\mathcal{J}^+(p) := \{q \in M / p \ll q\}$  is called the chronological future of the point  $p$ , and  $\mathcal{J}^-(p) := \{q \in M / q \ll p\}$  is called the chronological past. The causal future and past are defined similarly by replacing  $\ll$  by  $<$ . The future or past sets of a set  $S \subset M$  are defined by

$$\mathcal{J}^\pm(S) := \bigcup_{p \in S} \mathcal{J}^\pm(p), \quad \mathcal{I}^\pm(S) := \bigcup_{p \in S} \mathcal{I}^\pm(p),$$

and one easily checks that  $\mathcal{J}^\pm(S)$  are open, but that  $\mathcal{I}^\pm(S)$  need not be closed in general.

A future set  $F \subset M$  by definition has the form  $F = \mathcal{J}^+(S)$  for some set  $S \subset M$ . Similarly, a past set satisfies  $F = \mathcal{J}^-(S)$  for some  $S$ . A set is called achronal if no two points are connected by a time-like trip. Observe that a set can be spacelike at every point without being achronal and that an achronal set can be null at some (or even at every) point. A set  $B \subset M$  is called an achronal boundary if it is the boundary of a future set, that is  $B = \partial \mathcal{J}^+(S) = \overline{\mathcal{J}^+(S)} \setminus \mathcal{J}^+(S)$  for some  $S \subset M$ . One can check that given a non-empty achronal boundary the manifold can be

partitioned as  $M = P \cup B \cup F$ , where  $B$  is the boundary of both  $F$  and  $P$  and, moreover, any trip from  $p \in P$  to  $q \in F$  meets  $B$  at a unique point. Observe also that any achronal boundary is a Lipschitz continuous  $n$ -manifold. For instance, in Section 6 below, we will be interested in the geometry of past null cones, that is the sets  $\partial\mathcal{J}^-(p)$  for  $p \in M$ .

Given an arbitrary achronal and closed set  $S \subset M$ , we define the (future or past) domains of dependence of  $S$  in  $M$  by

$$\begin{aligned}\mathcal{D}^\pm(S) &:= \{p \in M / \text{every future (resp. past) endless trip containing } p \text{ meets } S\}, \\ \mathcal{D}(S) &:= \mathcal{D}^-(S) \cup \mathcal{D}^+(S).\end{aligned}$$

Observe that domains of dependence are closed sets. Next, define the (future or past) Cauchy horizons

$$\begin{aligned}\mathcal{H}^\pm(S) &:= \{p \in \mathcal{D}^\pm(S) / \mathcal{J}^\pm(p) \cap \mathcal{D}^\pm(S) = \emptyset\} = \mathcal{D}^\pm(S) \setminus \mathcal{J}^\pm(\mathcal{D}^\pm(S)), \\ \mathcal{H}(S) &:= \mathcal{H}^-(S) \cup \mathcal{H}^+(S).\end{aligned}$$

For instance, the future Cauchy horizon is the future boundary of the future domain of dependence of  $S$ . One can check that the Cauchy horizons are closed and achronal sets, with  $\partial\mathcal{D}^+(S) = \mathcal{H}^+(S) \cup S$  and  $\partial\mathcal{D}(S) = \mathcal{H}(S)$ .

Finally, a (future) Cauchy hypersurface for  $M$  is defined as an achronal (but not necessarily closed) set  $S$  satisfying  $\mathcal{D}^+(S) = M$ . For instance, it is sufficient for  $\bar{S}$  to be smooth, achronal, spacelike and such that every endless null geodesic meet  $M$ .

## Reference metric

As explained in the introduction one should not use the Lorentzian metric to compute the norm of a tensor since the Lorentzian norm may vanish even when the tensor does not. This motivates the introduction of a ‘‘reference’’ Riemannian metric associated with a time-like vector field, as follows.

Let  $T$  be a future-oriented, time-like, unit vector field, satisfying therefore  $g_p(T, T) = -1$  at every point  $p$ . We refer to  $T$  as the *reference vector field* prescribed on  $M$ . Introduce a moving frame  $E_\alpha$  ( $\alpha = 0, 1, \dots, n$ ) defined in  $M$ , that is,  $E_\alpha$  is an orthonormal basis of the tangent space at every point and consists of the vector  $e_0 = T$  supplemented with  $n$  spacelike unit vectors  $e_j$  ( $j = 1, \dots, n$ ). Denoting by  $E^\alpha$  the corresponding dual frame, the Lorentzian metric tensor takes the form

$$g = \eta_{\alpha\beta} E^\alpha \otimes E^\beta,$$

where  $\eta_{\alpha\beta}$  is the Minkowski ‘‘metric’’. This decomposition suggests to consider the Riemannian version obtained by switching the minus sign in  $\eta_{00} = -1$  into a plus sign, that is

$$g_T := \delta_{\alpha\beta} E^\alpha \otimes E^\beta,$$

where  $\delta_{\alpha\beta}$  is the Euclidian ‘‘metric’’. Clearly,  $g_T$  is a positive definite metric; it is referred to as the *reference Riemannian metric* associated with the frame  $E_\alpha$ .

For every  $p \in M$ , since  $T_p$  is time-like, the restriction of the metric  $g_p$  to the orthogonal complement  $\{T_p\}^\perp \subset T_p M$  is positive definite, and the reference metric can be computed as follows: if  $V = a T_p + V'$  and  $W = b T_p + W'$  with  $V', W' \in \{T_p\}^\perp$ , then

$$g_{T,p}(V, W) = a b + g_p(V', W').$$

In the following, we use the notation

$$g_{T,p}(V, W) = \langle V, W \rangle_{T,p}, \quad g_{T,p}(V, V) = |V|_{T,p}^2$$

for vectors; the norm of tensors is defined and denoted similarly.

In contrast with the Lorentzian norm, the Riemannian norm  $|A|_{T,p}$  of a tensor  $A$  at a point  $p \in M$  vanishes if and only if the tensor is zero at  $p$ . Moreover, as long as  $T$  remains in a compact subset of the bundle of half-cone  $T^+ M$ , the norms associated with different reference vectors are equivalent.

The reference Riemannian metric also allows one to define the functional norms for Lebesgue and Sobolev spaces of tensors defined on  $M$  (as well as on submanifolds of  $M$ ), allowing us for instance to view  $\mathbf{L}^2(M, g_T)$  as a Banach space. In particular, we will use later the  $\mathbf{L}^2$  norm of a tensor field  $T$  on  $M$  restricted to an hypersurface  $\Sigma$ :

$$\|\nabla h\|_{\mathbf{L}^2(\Sigma, g_T)} := \int_{\Sigma} |\nabla h|_T^2 dV_{\Sigma, g_T},$$

where  $dV_{\Sigma, g_T}$  is the volume form induced on  $\Sigma$  by the reference Riemannian metric. The functional norm above depends upon the choice of the vector field  $T$ , but another choice of  $T$  would give rise to an equivalent norm (provided  $T$  remains in a fixed compact subset). Observe in passing that the volume forms associated with the metrics  $g$  and  $g_T$  coincide, so that the spacetime integrals of *functions* in  $(M, g)$  or  $(M, g_T)$  coincide; for instance, the volume  $\mathbf{Vol}_g(A)$  and  $\mathbf{Vol}_{g_T}(A)$  of a set  $A \subset M$  coincide.

Furthermore, we observe that in order to define the reference metric  $g_T$  at a given point  $p$ , it suffices to prescribe a future-oriented time-like unit *vector*  $T$  at that point  $p$  only; it is not necessary to prescribe a vector field. In the situation where the reference metric need only be defined at the base point  $p$ , we refer to  $T$  as the *reference vector* (rather than vector field) and we refer to  $(p, T) \in T_1^+ M$  as the *observer at the point  $p$* . This will be the standpoint adopted for our main result in Section 7 below.

## Exponential map

On a complete Riemannian manifold the exponential map  $\mathbf{exp}_p : T_p M \rightarrow M$  at some point  $p \in M$  is defined on the whole tangent space  $T_p M$  and is smooth. For sufficiently small radius  $r$  the restriction of  $\mathbf{exp}_p$  to the ball  $B_{g_p}(0, r) \subset T_p M$  (determined by the metric  $g_p$  at the point  $p$ ) is a diffeomorphism on its image. The radius of injectivity at the point  $p$  is defined as the largest value of  $r$  such that the restriction  $\mathbf{exp}_p|_{B_{g_p}(0, r)}$  is a global diffeomorphism.



In the Lorentzian case, the exponential map is defined similarly but some care is needed in defining the notion of radius of injectivity. First of all, if the manifold is not geodesically complete (which is a rather generic situation, as illustrated by Penrose and Hawking's incompleteness theorems [14]), the map  $\mathbf{exp}_p$  need not be defined on the whole tangent space  $T_pM$  but only on a neighborhood of the origin in  $T_pM$ . More importantly, the Lorentzian norm of a non-zero vector may well vanish; consequently, the radius of injectivity should not be defined directly from the Lorentzian metric  $g$ . The definition below depends on the prescribed Riemannian metric  $g_{T,p}$  at the point  $p$ .

**Definition 2.1.** *The conjugate radius  $\mathbf{Conj}_g(M, p, T)$  of an observer  $(p, T) \in T_1^+M$  is the largest radius  $r$  such that the exponential map  $\mathbf{exp}_p$  is a local diffeomorphism from the Riemannian ball  $B_T(0, r) = B_{g_{T,p}}(0, r) \subset T_pM$  to a neighborhood of  $p$  in the manifold  $M$ . Similarly, the injectivity radius  $\mathbf{Inj}_g(M, p, T)$  of an observer  $(p, T) \in T_1^+M$  is the largest radius  $r$  such that the exponential map is a global diffeomorphism at every point of the ball  $B_T(0, r)$ .*

When a vector field  $T$  is prescribed on the manifold (rather than a vector at the point  $p$ ), we use the notation  $\mathbf{Inj}_g(M, p, T)$  instead of  $\mathbf{Inj}_g(M, p, T_p)$ . The radii  $\mathbf{Conj}_g(M, p, T)$  and  $\mathbf{Inj}_g(M, p, T)$  are essentially independent of the choice of the reference vector, as long as it remains in a fixed compact subset of  $T_p^+M$ .

We also need the notion of injectivity radius for null cones. Given a point  $p \in M$  and a reference vector  $T \in T_pM$ , we consider the *past null cone* at  $p$ ,

$$N_p^- := \{X \in T_pM \mid g_p(X, X) = 0, g_p(T, X) \geq 0\},$$

which is defined a subset of the tangent space at  $p$ . Denote by

$$B_T^N(0, r) = B_{g_{T,p}}^N(0, r) := B_{g_{T,p}}(0, r) \cap N_p^-$$

the intersection of the Riemannian  $g_{T,p}$ -ball with radius  $r$  and the past null cone, and by

$$\mathcal{N}^-(p) := \partial \mathcal{J}^-(p)$$

the past null cone at  $p$ .

Consider now the restriction of  $\mathbf{exp}_p$  to the past null cone, denoted by

$$\mathbf{exp}_p^N : B_T^N(0, r) \subset N_p^- \rightarrow \mathcal{N}^-(p) \subset M,$$

which we refer to as the *null exponential map*.

**Definition 2.2.** *The (past) null conjugate radius  $\mathbf{Null Conj}_g(M, p, T)$  of an observer  $(p, T) \in T_1^+M$  is the largest radius  $r$  such that the null exponential map  $\mathbf{exp}_p^N$  is a local diffeomorphism from the punctured Riemannian ball  $B_T^N(0, r) \setminus \{0\} \subset T_pM$  to a neighborhood of  $p$  in the past null cone. The null injectivity radius  $\mathbf{Null Inj}_g(M, p, T)$  of an observer  $(p, T) \in T_1^+M$  is defined similarly by requiring the map  $\mathbf{exp}_p^N$  to be a global diffeomorphism.*

### 3 Lorentzian manifold endowed with a reference vector field

#### A first injectivity radius estimate

From now on, we fix a reference vector field  $T$  which allows us to define the Riemannian metric  $g_T$  and compute the norms of tensors. We begin with a set of assumptions encompassing a large class of Lorentzian manifolds with  $L^\infty$  bounded curvature and we state our first injectivity estimate, in Theorem 3.1 below. The forthcoming sections will be devoted to further generalizations and variants of this result.

We fix a point  $p \in M$  and assume that a domain  $\Omega \subset M$  containing  $p$  is foliated by spacelike hypersurfaces  $\Sigma_t$  with future-oriented time-like unit normal  $T$ ,

$$\Omega = \bigcup_{t \in [-1,1]} \Sigma_t. \quad (3.1)$$

The positive coefficient  $n$  is defined by the relation  $\frac{\partial}{\partial t} = nT$ , or

$$n^2 := -g\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right).$$

In the context of general relativity,  $n$  is the proper time of an observer moving orthogonally to the hypersurfaces, and is called the *lapse function*. The geometry of the foliation is determined by this function  $n$  together with the Lie derivative  $\mathcal{L}_T g$ . The latter is nothing but the second fundamental form, or extrinsic curvature, of the slices  $\Sigma_t$  embedded in the manifold  $M$ .

We always assume that the geodesic ball  $\mathcal{B}_{\Sigma_0}(p, 1) \subset \Sigma_0$  (determined by the induced metric  $g|_{\Sigma_0}$ ) is compactly contained in  $\Sigma_0$ . We introduce the following assumptions:

$$e^{-K_0} \leq n \leq e^{K_0} \quad \text{in } \Omega, \quad (A1)$$

$$|\mathcal{L}_T g|_T \leq K_1 \quad \text{in } \Omega, \quad (A2)$$

$$|\mathbf{Rm}_g|_T \leq K_2 \quad \text{in } \Omega, \quad (A3)$$

$$\mathbf{Vol}_{g|_{\Sigma_0}}(\mathcal{B}_{\Sigma_0}(p, 1)) \geq v_0, \quad (A4)$$

where  $K_0, K_1, K_2$  and  $v_0$  are positive constants. Observe that Assumption (A4) is a condition on the initial slice only; together with the other assumptions it actually implies a lower volume bound for every slice of the foliation.

We will prove:

**Theorem 3.1** (Injectivity radius of foliated manifolds). *Let  $M$  be a differentiable manifold endowed with a Lorentzian metric  $g$  satisfying the regularity assumptions (A1)–(A4) at some point  $p$  and for some foliation (3.1). Then, there exists a positive constant  $i_0$  depending only upon the foliation bounds  $K_0, K_1$ , the curvature bound  $K_2$ , the volume bound  $v_0$ , and the dimension of the manifold such that the injectivity radius at  $p$  satisfies*

$$\mathbf{Inj}_g(M, p, T) \geq i_0.$$

The following section is devoted to the proof of this theorem. Observe that the conditions (A1)–(A4) are *local* about one point of the manifold and are stated in purely geometric terms, requiring no particular choice of coordinates. Of course, the conclusion of Theorem 3.1 hold globally in  $M$  if the assumptions (A1)–(A4) hold also globally at every point of the manifold. Our assumptions do depend on the choice of the time-like vector field  $T$ , but the dependence of the constants arising in (A1)–(A4) should not be essential; however, it is conceivable that, when applying this theorem in a specific situation a quantitatively sharper estimate would be obtained with a choice of an “almost Killing” field, that is a field  $T$  corresponding to a “small” Lie derivative  $\mathcal{L}_T g$ . Later in Section 7, a more general approach is presented in which the vector field  $T$  is constructed from a single vector prescribed at the point  $p$ .

### Basic estimates on the reference metric

To establish Theorem 3.1 it is convenient to introduce coordinates on  $\Omega$ , chosen as follows. Fix arbitrarily some coordinates  $(x^i)$  on the initial slice  $\Sigma_0$ . Then, transport these coordinates to the whole of  $\Omega$  along the integral curves of the vector field  $T$ . This construction generates coordinates  $(x^\alpha)$  on  $\Omega$  such that  $x^0 = t$  and the vector  $\partial/\partial t$  is orthogonal to  $\partial/\partial x^j$ , so that the Lorentzian metric takes the form

$$g = -n^2 dt^2 + g_{ij} dx^i dx^j, \quad (3.2)$$

where  $n$  is the lapse function and  $g_{ij}$  is the Riemannian metric induced on the slices  $\Sigma_t$ . The reference Riemannian metric in the domain  $\Omega$  then takes the form

$$g_T = n^2 dt^2 + g_{ij} dx^i dx^j, \quad (3.3)$$

and the Riemannian norm of a vector  $X$  has the explicit form:  $g_T(X, X) := n^2 X^0 X^0 + X^j X_j$ .

We want to control the discrepancy between the reference Riemannian metric  $g_T$  and the original Lorentzian metric  $g$ , as measured in the connections  $\nabla$  and  $\nabla_{g_T}$  and the curvature tensors  $\mathbf{Rm}$  and  $\mathbf{Rm}_{g_T}$ . Clearly, these estimates should involve the constants arising in (A1)–(A4). Consider the general class of metrics

$$\tilde{g} := f dt^2 + g_{ij} dx^i dx^j, \quad (3.4)$$

which allows us to recover both the Lorentzian ( $f = -n^2$ ) and the Riemannian ( $f = n^2$ ) metrics.

In view of the expressions of the Christoffel symbols and the Riemann curvature

$$\begin{aligned} \tilde{\Gamma}_{\alpha\beta}^{\gamma} &:= \frac{1}{2} \tilde{g}^{\gamma\delta} \left( \frac{\partial \tilde{g}_{\delta\beta}}{\partial x^\alpha} + \frac{\partial \tilde{g}_{\delta\alpha}}{\partial x^\beta} - \frac{\partial \tilde{g}_{\alpha\beta}}{\partial x^\delta} \right), \\ \tilde{R}_{\alpha\beta\delta}^{\zeta} &:= \frac{\partial \tilde{\Gamma}_{\beta\delta}^{\zeta}}{\partial x^\alpha} - \frac{\partial \tilde{\Gamma}_{\alpha\delta}^{\zeta}}{\partial x^\beta} + \tilde{\Gamma}_{\alpha\eta}^{\zeta} \tilde{\Gamma}_{\beta\delta}^{\eta} - \tilde{\Gamma}_{\beta\eta}^{\zeta} \tilde{\Gamma}_{\alpha\delta}^{\eta}, \\ \tilde{R}_{\alpha\beta\gamma\delta} &:= \tilde{g}_{\gamma\zeta} \tilde{R}_{\alpha\beta\delta}^{\zeta}, \quad \tilde{R}_{\alpha\beta} := \tilde{R}_{\alpha\gamma\beta\delta} \tilde{g}^{\gamma\delta}, \end{aligned}$$

we compute explicitly the Christoffel symbols associated with the metric  $\widetilde{g}$ ,

$$\begin{aligned}\widetilde{\Gamma}_{00}^0 &= \frac{1}{2f} \frac{\partial f}{\partial t}, & \widetilde{\Gamma}_{0i}^0 &= \frac{1}{2f} \frac{\partial f}{\partial x^i}, & \widetilde{\Gamma}_{ij}^0 &= -\frac{1}{2f} \frac{\partial g_{ij}}{\partial t}, \\ \widetilde{\Gamma}_{00}^k &= -\frac{1}{2} g^{kl} \frac{\partial f}{\partial x^l}, & \widetilde{\Gamma}_{i0}^k &= \frac{1}{2} g^{kl} \frac{\partial g_{li}}{\partial t}, & \widetilde{\Gamma}_{ij}^k &= \Gamma_{ij}^k,\end{aligned}\tag{3.5}$$

as well as the non-trivial curvature terms

$$\begin{aligned}\widetilde{R}_{ijkl} &= R_{ijkl} - \frac{1}{4f} \left( \frac{\partial g_{ik}}{\partial t} \frac{\partial g_{jl}}{\partial t} - \frac{\partial g_{il}}{\partial t} \frac{\partial g_{jk}}{\partial t} \right), \\ \widetilde{R}_{0jl}^p &= \frac{\partial \Gamma_{jl}^p}{\partial t} - \frac{\partial}{\partial x^j} \left( \frac{1}{2} g^{pq} \frac{\partial g_{ql}}{\partial t} \right) + \frac{1}{2} g^{pq} \frac{\partial}{\partial t} g_{qk} \Gamma_{jl}^k - \Gamma_{jk}^p \left( \frac{1}{2} g^{kq} \frac{\partial g_{lq}}{\partial t} \right) \\ &\quad + \left( \frac{1}{4f} g^{pq} \frac{\partial f}{\partial x^q} \right) \frac{\partial g_{jl}}{\partial t} - \frac{1}{2} g^{pq} \frac{\partial g_{qj}}{\partial t} \frac{1}{2f} \frac{\partial f}{\partial x^l}, \\ \widetilde{R}_{0jl} &= \frac{1}{2} \left( \nabla_l \left( \frac{\partial}{\partial t} g_{ij} \right) - \nabla_i \left( \frac{\partial}{\partial t} g_{lj} \right) \right) + \frac{1}{4f} \left( \frac{\partial f}{\partial x^i} \frac{\partial g_{jl}}{\partial t} - \frac{\partial f}{\partial x^l} \frac{\partial g_{ij}}{\partial t} \right), \\ \widetilde{R}_{i00}^p &= \frac{\partial}{\partial x^i} \left( -\frac{1}{2} g^{pq} \frac{\partial f}{\partial x^q} \right) - \frac{\partial}{\partial t} \left( \frac{1}{2} g^{pq} \frac{\partial g_{qi}}{\partial t} \right) + \Gamma_{il}^p \left( -\frac{1}{2} g^{lq} \frac{\partial f}{\partial x^q} \right) \\ &\quad - \left( \frac{1}{2} g^{pq} \frac{\partial g_{ql}}{\partial t} \right) \left( \frac{1}{2} g^{lr} \frac{\partial g_{ri}}{\partial t} \right) + \frac{1}{2f} \frac{\partial f}{\partial t} \left( \frac{1}{2} g^{pq} \frac{\partial g_{qi}}{\partial t} \right) + \frac{1}{2} g^{pq} \frac{\partial f}{\partial x^q} \frac{1}{2f} \frac{\partial f}{\partial x^i},\end{aligned}$$

and

$$\widetilde{R}_{i0j0} = -\frac{1}{2} \left( \nabla_i \nabla_j f + \frac{\partial^2 g_{ij}}{\partial t^2} \right) + \frac{1}{4} g^{pq} \frac{\partial g_{ip}}{\partial t} \frac{\partial g_{jq}}{\partial t} + \frac{1}{4f} \frac{\partial f}{\partial t} \frac{\partial g_{ij}}{\partial t} + \frac{1}{4f} \frac{\partial f}{\partial x^i} \frac{\partial f}{\partial x^j}.$$

By applying the formulas above to both metrics  $g, g_T$  we estimate the Christoffel symbols, as follows. Recall that the difference  $\Gamma_{\alpha\beta}^{\gamma'} - \Gamma_{g_T, \alpha\beta}^{\gamma'}$  can be regarded as a tensor field on  $M$ , so that the following (Riemannian) norm squared is a scalar field on the manifold  $M$ :

$$|\nabla_{g_T} - \nabla|_T^2 := |\Gamma_{g_T} - \Gamma_T|^2 = (\Gamma_{g_T, \beta\gamma}^{\alpha} - \Gamma_{\beta\gamma}^{\alpha}) (\Gamma_{g_T, \beta'\gamma'}^{\alpha'} - \Gamma_{\beta'\gamma'}^{\alpha'}) g_{T, \alpha\alpha'} g_T^{\beta\beta'} g_T^{\gamma\gamma'}.$$

We need also the expression of the Lie derivative of  $g$  along the vector field  $T$ . By a direct computation from (3.2) we obtain

$$(\mathcal{L}_T g)_{00} = 0, \quad (\mathcal{L}_T g)_{0i} = \frac{1}{n} \frac{\partial n}{\partial x^i}, \quad (\mathcal{L}_T g)_{ij} = \frac{1}{n} \frac{\partial g_{ij}}{\partial t}.\tag{3.6}$$

**Lemma 3.2** (Levi-Cevita connection of the reference metric). *Suppose that  $g$  satisfies Assumptions (A1)-(A2). Then, the covariant derivative of the Lorentzian and Riemannian metrics are comparable, precisely*

$$|\nabla_{g_T} - \nabla|_T = n^2 |\mathcal{L}_T g|_T^2 \leq e^{2K_0} K_1^2 =: K_3.$$

*Proof.* In view of (3.5) the difference  $\Gamma_{g_T} - \Gamma$  depends essentially upon the terms  $\frac{\partial n}{\partial x^i}$  and  $\frac{\partial g_{ij}}{\partial t}$  which precisely appear in the expression of the Lie derivative (3.6). We omit the details.  $\square$

It is important to observe that the difference between the curvature tensors can not be similarly estimated, and that this is one of the main difficulties to deal with in the present work.

For future reference we provide here the expressions of certain curvature coefficients of  $g$  and  $g_T$  in terms of (first-order derivatives of) the lapse function  $n$  and the induced metric  $g_{jk}$ :

$$\begin{aligned} R_{ijkl} &= R_{ijkl}^\Sigma + \frac{1}{4n^2} \left( \frac{\partial g_{ik}}{\partial t} \frac{\partial g_{jl}}{\partial t} - \frac{\partial g_{il}}{\partial t} \frac{\partial g_{jk}}{\partial t} \right), \\ R_{0jil} &= \frac{1}{2} \left( \nabla_l \left( \frac{\partial}{\partial t} g_{ij} \right) - \nabla_i \left( \frac{\partial}{\partial t} g_{lj} \right) \right) + \frac{1}{4n^2} \left( \frac{\partial n^2}{\partial x^i} \frac{\partial g_{jl}}{\partial t} - \frac{\partial n^2}{\partial x^l} \frac{\partial g_{ij}}{\partial t} \right), \\ R_{i0j0} &= \frac{1}{2} \left( \nabla_i \nabla_j (n^2) - \frac{\partial^2 g_{ij}}{\partial t^2} \right) + \frac{1}{4} g^{pq} \frac{\partial g_{ip}}{\partial t} \frac{\partial g_{jq}}{\partial t} + \frac{1}{4n^2} \frac{\partial n^2}{\partial t} \frac{\partial g_{ij}}{\partial t} - \frac{1}{4n^2} \frac{\partial n^2}{\partial x^i} \frac{\partial n^2}{\partial x^j}, \end{aligned}$$

and

$$\begin{aligned} R_{T,ijkl} &= R_{ijkl}^\Sigma - \frac{1}{4n^2} \left( \frac{\partial g_{ik}}{\partial t} \frac{\partial g_{jl}}{\partial t} - \frac{\partial g_{il}}{\partial t} \frac{\partial g_{jk}}{\partial t} \right), \\ R_{T,0jil} &= \frac{1}{2} \left( \nabla_l \left( \frac{\partial}{\partial t} g_{ij} \right) - \nabla_i \left( \frac{\partial}{\partial t} g_{lj} \right) \right) + \frac{1}{4n^2} \left( \frac{\partial n^2}{\partial x^i} \frac{\partial g_{jl}}{\partial t} - \frac{\partial n^2}{\partial x^l} \frac{\partial g_{ij}}{\partial t} \right), \\ R_{T,i0j0} &= \frac{1}{2} \left( \nabla_i \nabla_j (-n^2) - \frac{\partial^2 g_{ij}}{\partial t^2} \right) + \frac{1}{4} g^{pq} \frac{\partial g_{ip}}{\partial t} \frac{\partial g_{jq}}{\partial t} + \frac{1}{4n^2} \frac{\partial n^2}{\partial t} \frac{\partial g_{ij}}{\partial t} + \frac{1}{4n^2} \frac{\partial n^2}{\partial x^i} \frac{\partial n^2}{\partial x^j}, \end{aligned}$$

where  $R_{ijkl}^\Sigma$  denotes the induced curvature tensor on the time slices  $\Sigma = \Sigma_t$ .

## 4 Derivation of the first injectivity radius estimate

In this section we provide a proof of Theorem 3.1.

*Step 1. Radius of definition of the exponential map.* First of all, we note that the injectivity radius of the Riemannian metric  $g|_{\Sigma_0}$  induced on the initial hypersurface  $\Sigma_0 = t^{-1}(0)$  is controled, as follows. Using Assumptions (A3) and (A4), we see that the Riemann curvature of the metric  $g|_{\Sigma_0}$  is bounded and the volume of the unit geodesic ball  $\text{Vol}_{g|_{\Sigma_0}}(\mathcal{B}_{\Sigma_0}(p, 1))$  is bounded below. Therefore, according to [10], there exists a constant  $i_1 = i_1(K_2, v_0)$  such that the injectivity radius of  $g|_{\Sigma_0}$  at the point  $p$  is  $i_1$  at least:

$$\text{Inj}_{g|_{\Sigma_0}}(\Sigma_0, p) \geq i_1.$$

Moreover, according to [16] we can also assume that  $i_1$  is sufficiently small so that, given any  $\varepsilon > 0$  there exists a coordinates  $(x^\alpha)$  defined in a ball with definite

size near  $p$ , with  $x^\alpha(p) = 0$ , such that the metric  $g|_{\Sigma_0}$  is close to the  $n$ -dimensional Euclidian metric  $g_{E'} = \delta_{ij}$  (in these coordinates). More precisely, on the initial slice  $\Sigma_0$  we have

$$e^{-\varepsilon} \delta_{ij} \leq g_{ij}(0, x^1, \dots, x^n) \leq e^\varepsilon \delta_{ij}, \quad (x^1, \dots, x^n) \in B_{E'}(0, i_1),$$

where we have set  $B_{E'}(0, r) := \{(x^1)^2 + \dots + (x^n)^2 < r^2\} \subset \mathbb{R}^n$ . The latter can be regarded as a subset of  $\Sigma_0$  by identifying a point with its coordinates, and we also use the notation  $\mathcal{B}_{E'}(p, r)$  for this Euclidian ball.

We can next introduce some coordinates  $(x^\alpha) = (t, x^j)$  on the manifold, by propagating the coordinates  $(x^j)$  chosen on  $\Sigma_0$  along the integral curve of the vector field  $T$ . This construction allows us to cover the domain  $\Omega$ . From Assumption (A2) (together with (A1) and (3.6)) we deduce that the induced metric on *each slice* of the foliation is comparable with the  $n$ -dimensional Euclidian metric in some time interval  $[-i_2, i_2]$ , that is

$$\begin{aligned} (e^{-\varepsilon} - K i_2) \delta_{ij} &\leq g_{ij}(x) \leq (e^\varepsilon + K i_2) \delta_{ij}, \\ x = (t, x^1, \dots, x^n) &\in [-i_2, i_2] \times B_{E'}(0, i_1), \end{aligned}$$

for some  $K > 0$  depending only on  $K_0, K_1, K_2$ .

We then restrict attention to a smaller radius  $i_2 = i_2(K_0, K_1, K_2) \leq i_1$  such that  $e^{-\varepsilon} - K i_2 > 0$ , and we pick up  $c_1 \geq 0$  sufficiently large so that  $e^{-c_1} \leq e^{-\varepsilon} - K i_2 \leq e^\varepsilon + K i_2 \leq e^{c_1}$ . In turn, in view of Assumption (A1) on the lapse function  $n$  and of the expression (3.3) of the reference Riemannian metric  $g_T$ , the above inequalities imply that the reference Riemannian metric  $g_T$  is comparable to the  $(n + 1)$ -dimensional Euclidean metric:

$$e^{-c_2} \delta_{\alpha\beta} \leq g_{T,\alpha\beta} \leq e^{c_2} \delta_{\alpha\beta}, \quad x = (t, x^1, \dots, x^n) \in [-i_2, i_2] \times B_{E'}(0, i_2)$$

for some constant  $c_2 \geq c_1$  depending upon  $c_1$  and  $K_0$ .

Introducing on the manifold the  $(n + 1)$ -dimensional Euclidian metric  $E$  (which we define in the constructed coordinates  $(x^\alpha)$  and is, of course, independent of the point on the manifold) and the corresponding Euclidian metric ball  $\mathcal{B}_E(p, i_2)$ , we have established

$$e^{-c_2} g_E \leq g_{T,q} \leq e^{c_2} g_E, \quad q \in \mathcal{B}_E(p, i_2). \quad (4.1)$$

In the following we use the notation  $|X|_E$  for the Euclidian norm of a vector  $X$ .

Our first task is to determine the radius of a ball on which the exponential map is well-defined. This radius depends upon the reference vector field  $T$ . Let  $\gamma : [0, s_0] \rightarrow M$  be a geodesic associated with the Lorentzian metric  $g$  and satisfying  $\gamma(0) = p$ . Assume that this geodesic is included in the Euclidian ball  $\mathcal{B}_E(p, i_2)$  (in which we have a good control of the metric  $g_T$ ). Obviously, we have

$$\langle \gamma'(s), \gamma'(s) \rangle_g = \langle \gamma'(0), \gamma'(0) \rangle_g, \quad s \in [0, s_0].$$

On the other hand, to determine the length of  $\gamma'(s)$  with respect to the reference metric  $g_T$ , we proceed as follows:

$$\begin{aligned} \left| \frac{d}{ds} \langle \gamma'(s), \gamma'(s) \rangle_T \right| &= \left| \nabla_{T, \gamma'(s)} (g_T(\gamma'(s), \gamma'(s))) \right| = 2 \left| \langle \nabla_{g_T, \gamma'(s)} \gamma'(s), \gamma'(s) \rangle_T \right| \\ &= 2 \left| \langle (\nabla_{g_T} - \nabla_g)_{\gamma'(s)} \gamma'(s), \gamma'(s) \rangle_T \right| \\ &\leq 2 |\nabla_{g_T} - \nabla_g|_T |\gamma'(s)|_T^3. \end{aligned}$$

So, by Lemma 3.2,  $\left| \frac{d}{ds} |\gamma'(s)|_T^2 \right| \leq 2K_3 |\gamma'(s)|_T^3$ , and, in consequence,

$$\left| \frac{d}{ds} |\gamma'(s)|_T^{-1} \right| \leq K_3.$$

By integration of the above inequality and *provided*  $s$  is small enough so that  $2sK_3 |\gamma'(0)|_T < 1$ , we see that

$$\frac{1}{2} |\gamma'(0)|_T \leq |\gamma'(s)|_T \leq 2 |\gamma'(0)|_T. \quad (4.2)$$

In view of (4.1) this implies

$$\frac{e^{-c_2}}{2} |\gamma'(0)|_E \leq |\gamma'(s)|_E \leq 2e^{c_2} |\gamma'(0)|_E. \quad (4.3)$$

These inequalities hold for all  $s \in [0, 1/(2K_3 |\gamma'(0)|_T)]$  as long as  $\gamma(s) \in \mathcal{B}_E(p, i_2)$ . In particular, by restricting attention to geodesics whose initial vector has unit Euclidian length,  $|\gamma'(0)|_E = 1$ , we see that  $\gamma([0, r_2]) \subset \mathcal{B}_E(p, i_2)$  where  $r_2 := i_2 e^{-c_2}/2$ . In turn, this establishes that the exponential map at the point  $p$  is well-defined on the ball  $B_E(0, r_2)$  with a range included in the geodesic ball  $\mathcal{B}_E(p, i_2)$ .

*Step 2. Conjugate radius estimate.* Our second task is to determine a ball on which the exponential map is a local diffeomorphism, and we therefore need to control the length of a Jacobi field along a geodesic. Let  $\gamma : [0, r_2] \rightarrow M$  be a  $g$ -geodesic satisfying  $\gamma(0) = p$  and  $|\gamma'(0)|_E = 1$ . By the discussion in Step 1 we already know that the curve  $\gamma$  lies in  $\mathcal{B}_E(p, i_2)$  and that  $\max_{s \in [0, r_2]} |\gamma'(s)|_T \leq 2e^{2c_2}$ . Given an arbitrary Jacobi field along  $\gamma$ ,  $J = J(s)$ , satisfying

$$\begin{aligned} J''(s) &= -\mathbf{Rm}(J(s), \gamma'(s))\gamma'(s), \\ J(0) &= 0, \quad |J'(0)|_T = 1, \end{aligned}$$

we need to control its Riemannian length  $F(s) := |J|_T(s)$ , as stated in (4.7) below.

Let  $[0, s_0]$  be the largest subinterval of  $[0, r_2/4]$  in which the inequality  $|J|_T \leq 1$  holds. Using the equation satisfied by the Jacobi field and taking into account the curvature bound (A3), we deduce that, in the interval  $[0, s_0]$ ,

$$\begin{aligned} \left| \frac{d}{ds} \langle \nabla_{\gamma'} J, \nabla_{\gamma'} J \rangle_T \right| &= 2 \left| \langle \nabla_{g_T, \gamma'} \nabla_{\gamma'} J, \nabla_{\gamma'} J \rangle_T \right| \\ &\leq 2 |\nabla_{g_T} - \nabla_g|_T |\gamma'|_T |\nabla_{\gamma'} J|_T^2 + 2K_2 |\gamma'|_T^2 |J|_T |\nabla_{\gamma'} J|_T. \end{aligned}$$

With (4.2) and the covariant derivative estimate in Lemma 3.2, we obtain

$$\left| \frac{d}{ds} |\nabla_{\gamma'} J|_T \right| \leq 4K_3 |\nabla_{\gamma'} J|_T + 8K_2. \quad (4.4)$$

We can next integrate (4.4) over an arbitrary interval  $[0, s] \subset [0, s_0]$ , use the initial condition on the Jacobi field, and obtain

$$1 + \frac{2K_2}{K_3} (1 - e^{-4K_3 s}) \leq |\nabla_{\gamma'} J|_T \leq 1 + \frac{2K_2}{K_3} (e^{4K_3 s} - 1).$$

Assuming that  $r_2$  is small enough so that  $\frac{2K_2}{K_3} (1 - e^{-4K_3 s}) \leq 1/2$  and  $\frac{2K_2}{K_3} (e^{4K_3 s} - 1) \leq$  we infer that

$$\frac{1}{2} \leq |\nabla_{\gamma'} J|_T \leq 2. \quad (4.5)$$

Hence, using this inequality and Lemma 3.2 we find  $\frac{d}{ds} |J|_T \leq 2 + 2K_3 \leq 1$ , so that

$$F(s) = |J|_T(s) \leq (2 + 2K_3) s \leq (2 + 2K_3) r_2. \quad (4.6)$$

Further assuming that  $(2 + 2K_3) r_2 \leq 1$  we conclude that  $s_0 = r_2$ .

Next, we want to improve the rough estimate (4.6). Since

$$\frac{d}{ds} \langle \nabla_{\gamma'} J, J \rangle_T = \langle \nabla_{g_T, \gamma'} \nabla_{\gamma'} J, J \rangle_T + \langle \nabla_{g_T, \gamma'} J, \nabla_{\gamma'} J \rangle_T$$

then by substituting the previous estimates of  $|J|_T(s)$  and  $|\nabla_{\gamma'} J|_T(s)$  and performing similar calculations as above, we get

$$e^{-c_3} \leq \frac{d}{ds} \langle \nabla_{\gamma'} J, J \rangle_T \leq e^{c_3}$$

for some constant  $c_3 > 0$ . By integration this implies

$$e^{-c_3} s \leq \langle \nabla_{\gamma'} J, J \rangle_T \leq e^{c_3} s$$

and we arrive at the following lower bound for the norm of the Jacobi field:

$$F(s) \geq \frac{|\langle \nabla_{\gamma'} J, J \rangle_T|}{|\nabla_{\gamma'} J|_T} \geq \frac{e^{-c_3} s}{2} \geq e^{-c_4} s$$

for some  $c_4 > 0$ .

On the other hand, using again the above estimates we have

$$\begin{aligned} \frac{d}{ds} F &\leq \frac{1}{F} (\langle \nabla_{g_T, \gamma'} J, J \rangle_T + K_3 F^2) \\ &\leq \frac{e^{c_4}}{s} (e^{c_3} s + K_3 (2 + 2K_3)^2 s^2) \leq e^{c_5} \end{aligned}$$

for some constant  $c_5 > 0$ . This leads to the upper bound

$$F(s) \leq e^{c_5} s.$$



In summary, we have established that the norm of the Jacobi field is comparable with  $s$ :

$$e^{-c_4} s \leq F(s) \leq e^{c_5} s, \quad s \in [0, r_2]. \quad (4.7)$$

By the definition of Jacobi fields these inequalities are equivalent to controlling the differential of the exponential map, that is for  $s \in [0, r_2]$

$$e^{-c_4} |W|_T \leq |d\exp_{p, s\gamma'(0)}(W)|_T \leq e^{c_5} |W|_T.$$

We conclude that the pull back of the reference metric to the tangent space at  $p$  satisfies

$$e^{-c_4} g_{T,p} \leq (\mathbf{exp}_p)^* g_T \leq e^{c_5} g_{T,p} \quad (4.8)$$

in the ball  $B_T(0, r_2) \subset T_p M$ .

In particular, since the conjugate radius of the Lorentzian metric is precisely defined from the reference Riemannian metric, these inequalities show that the conjugate radius of the exponential map is  $r_2$  at least.

*Step 3. Injectivity radius estimate.* We are now in a position to establish that  $\mathbf{Inj}_g(M, p, T) \geq r_3 := r_2 e^{-c_2}/4$ . We argue by contradiction and assume that  $\gamma_1 : [0, s_1] \rightarrow M$  and  $\gamma_2 : [0, s_2] \rightarrow M$  are two distinct  $g$ -geodesics satisfying  $\max(s_1, s_2) \leq r_3$  and

$$\begin{aligned} \gamma_1(0) = \gamma_2(0) = p, \quad |\gamma_1'(0)|_T = |\gamma_2'(0)|_T = 1, \\ \gamma_1(s_1) = \gamma_2(s_2) =: q. \end{aligned}$$

We will reach a contradiction and this will establish that the injectivity radius is greater or equal to  $r_3$  (as can be checked by using the fact that the exponential map is at least a local diffeomorphism).

By Step 1 we know that  $\gamma_1, \gamma_2 \subset \mathcal{B}_E(p, 2e^{2c_2} r_3)$ . By concatenating these two curves, we construct a geodesic loop containing  $p$ ,

$$\gamma = \gamma_2^{-1} \cup \gamma_1 : [0, s_1 + s_2] \rightarrow \mathcal{B}_E(p, 2e^{2c_2} r_3),$$

which need not be smooth at  $p$  or  $q$ . Since  $\gamma$  is contained in the image of the ball  $B_T(p, r_2)$  under the exponential map, we can define an homotopy of  $\gamma$  with the origin ( $x = 0$ ), by setting (in the coordinates constructed earlier)

$$\Gamma_\varepsilon(s) = \varepsilon \gamma(s), \quad \varepsilon \in [0, 1].$$

The curves  $\Gamma_\varepsilon : [0, s_1 + s_2] \rightarrow \mathcal{B}_E(p, 2e^{2c_2} r_3)$  satisfy

$$\Gamma_\varepsilon(0) = \Gamma_\varepsilon(s_1 + s_2) = p, \quad \Gamma_0([0, 1]) = p, \quad \Gamma_1 = \gamma.$$

Moreover, we have  $|\Gamma'_\varepsilon(s)|_E \leq \varepsilon 2e^{2c_2} \leq 2e^{2c_2}$  and thus  $|\Gamma'_\varepsilon(s)|_T \leq 2e^{3c_2}$ . In particular, the  $g_T$ -lengths (computed with the reference metric) of the loops  $\Gamma_\varepsilon$  are less than

$$L(\Gamma_\varepsilon, g_T) \leq 2e^{3c_2} r_3 = \frac{r_2}{2}.$$

Since the exponential map is a local diffeomorphism from the ball  $B_T(0, r_2) \subset T_p M$  to the manifold, and in view of the estimate (4.8) on the exponential map, it follows that all the loops  $\Gamma_\varepsilon$  can be lifted to the ball  $B_T(0, r_2)$  in the tangent space with the same origin 0. Consequently, we obtain a *continuous* family of curves  $\widetilde{\Gamma}_\varepsilon : [0, s_1 + s_2] \rightarrow T_p M$  satisfying

$$\widetilde{\Gamma}_\varepsilon(0) = 0, \quad \varepsilon \in [0, 1].$$

At this juncture we observe that, since  $\widetilde{\Gamma}_\varepsilon(s_1 + s_2)$  (for  $\varepsilon \in [0, 1]$ ) all cover the same point  $p$  and since the curve  $\widetilde{\Gamma}_0$  is trivial and the family is continuous,

$$\widetilde{\Gamma}_\varepsilon(s_1 + s_2) = 0, \quad \varepsilon \in [0, 1].$$

It remains to consider the lift of the original geodesic loop  $\gamma$ : under the lifting the geodesics  $\gamma_1, \gamma_2$  are sent to two distinct *line segments* (with respect to the vector space structure) originating at the origin 0 which obviously do not intersect. This is a contradiction and we conclude that, in fact,  $\mathbf{Inj}_g(M, p, T) \geq r_3$  as announced. This completes the proof of Theorem 3.1.

## 5 Convex functions and convex neighborhoods

We establish now the existence of convex functions and convex neighborhoods in  $M$ . Let us recall first some basic definitions. A function  $u$  is said to be *geodesically convex* if the composition of  $u$  with any geodesic is a convex function (of one variable). A set  $\Omega' \subset \Omega''$  is said to be *relatively geodesically convex in  $\Omega''$*  if, given any points  $p, q \in \Omega'$  and any geodesic (segment)  $\gamma$  from  $p$  to  $q$  contained in  $\Omega''$ , one has  $\gamma \subset \Omega'$ . A set  $\Omega'$  is said to be *geodesically convex in  $\Omega''$*  if  $\Omega'$  is relatively geodesically convex in  $\Omega''$  and, in addition, for any  $p, q$ , there exists a unique geodesic  $\gamma$  connecting  $p$  and  $q$  and lying in  $\Omega'$ .

We denote by  $d_T$  the distance function associated with the reference Riemannian metric  $g_T$ .

**Theorem 5.1** (Existence of geodesically convex functions). *Let  $(M, g)$  be a differentiable  $(n + 1)$ -manifold endowed with a Lorentzian metric  $g$ , satisfying the regularity assumptions (A1)-(A4) for some point  $p \in M$  and some future-oriented, unit, time-like vector field  $T$ , and let  $g_T$  be the reference Riemannian metric associated with  $T$ . Then, for any  $\varepsilon \in (0, 1)$  there exists a positive constant  $r_0$  depending only upon  $\varepsilon$ , the foliation bounds  $K_0, K_1$ , the curvature bound  $K_2$ , the volume bound  $v_0$ , and the dimension of the manifold and there exists a smooth function  $u$  defined on  $\mathcal{B}_T(p, r_0)$  such that*

$$\begin{aligned} (1 - \varepsilon) d_T(p, \cdot)^2 &\leq u \leq (1 + \varepsilon) d_T(p, \cdot)^2, \\ (2 - \varepsilon) g_T &\leq \nabla^2 u \leq (2 + \varepsilon) g_T. \end{aligned}$$

Hence, the function  $u$  above is equivalent to the Riemannian distance function from  $p$  and is geodesically convex for the Lorentzian metric. In the proof

given below, the function  $u$  is the Riemannian distance function associated with a new time-like vector field (denoted by  $N$  in the proof below). The following corollary is immediate and provides us with a control of the radius of convexity, which generalizes Whitehead theorem from Riemannian geometry [23, 6].

**Corollary 5.2** (Existence of geodesically convex neighborhoods). *Under the assumptions of Theorem 5.1, for any  $0 < r < r_0$  there exists a set  $\Omega_r \subset \Omega$  which is geodesically convex in  $\mathcal{B}_T(p, 2r_0)$  and satisfies*

$$\mathbf{exp}_p(\mathcal{B}_T(0, r)) \subset \Omega_r \subset \mathbf{exp}_p(\mathcal{B}_T(0, (1 + \delta)r)).$$

Moreover, one can always choose  $\Omega_r$  so that

$$\mathcal{B}_T(p, r) \subset \Omega_r \subset \mathcal{B}_T(p, (1 + \delta)r),$$

where  $\mathcal{B}_T(p, r)$  is the geodesic ball determined by the reference Riemannian metric.

*Proof of Theorem 5.1. Step 1. Synchronous coordinate system.* Given  $\varepsilon > 0$ , by applying the injectivity radius estimate in Theorem 3.1 to points near  $p$ , we see that there exists a constant  $r_0$  depending on  $K_0, K_1, K_2, v_0, \varepsilon, n$  such that for any  $q \in \mathcal{B}_T(p, 2r_0)$  the injectivity radius at  $q$  is  $2r_0$  at least, and we can assume that

$$e^{-\varepsilon} g_{T,q} \leq (\mathbf{exp}_q)^\star g_T \leq e^\varepsilon g_{T,q}, \quad \mathcal{B}_T(0, r_0) \subset T_q M, \quad q \in \mathcal{B}_T(p, 2r_0).$$

Let  $\gamma = \gamma(s)$  be the (backward) time-like geodesic satisfying  $\gamma(0) = p$  and  $\gamma'(0) = -T_p$ , and consider the (past) point  $q := \gamma(r_0/2)$ . The future null cone at  $q$  with radius  $r_0$  (the orientation being determined by the vector field  $T$ ) is defined by

$$C_q(r_0) := \left\{ V \in T_q M \mid |V|_{g_{T,q}} < r_0, |V|_{g_q}^2 < 0, \langle V, T \rangle > 0 \right\}.$$

Observe that the  $g_T$ -length of  $\gamma$  between  $p$  and  $q$  is approximately  $r_0/2$  and that the norm  $|\gamma'|_T$  is almost 1, while  $|\gamma'(q)|_{g_q}^2 = 1$  and  $\langle -\gamma', T \rangle_g > 0$ . By the injectivity radius estimate in Theorem 3.1 the exponential map at  $q$  is a diffeomorphism from  $C_q(r_0)$  onto its image which, moreover, contains the original point  $p$ .

Next, introduce the set of vectors that are “almost” parallel to  $T$ :

$$C_q(r_0, \varepsilon) := \left\{ V \in T_q M \mid |V|_{T,q} < r_0, \langle V, T \rangle_{g_q} > 0, \frac{\langle V, V \rangle_{g_q}}{\langle V, V \rangle_{T,q}} > 1 - \varepsilon \right\}.$$

The notation  $c(\varepsilon) > 0$  is used for constants that depend only on  $K_0, K_1, K_2, v_0, n, \varepsilon$  and satisfy  $\lim_{\varepsilon \rightarrow 0} c(\varepsilon) = 0$ . We claim that there is constant  $c(\varepsilon) > 0$  such that

$$\mathcal{B}_T(p, c(\varepsilon)r_0) \subset \mathbf{exp}_q(C_q(r_0, \varepsilon)). \quad (5.1)$$

Actually, we have  $\mathcal{B}_T(p, c(\varepsilon)r_0) \subset \mathcal{B}_T(q, (\frac{1}{2} + c(\varepsilon))r_0)$ , hence

$$\mathcal{B}_T(p, c(\varepsilon)r_0) \subset \mathbf{exp}_q\left(\mathcal{B}_T(0, \left(\frac{1}{2} + c(\varepsilon)\right)r_0)\right).$$

Since the metrics  $g_{T,0}$  and  $g_{T,q}$  are comparable (under the exponential map at  $q$ ) we see that geodesics  $\sigma$  connecting  $q$  and points of  $\mathcal{B}_T(p, c(\varepsilon)r_0)$  make an angle  $\leq c(\varepsilon)$  with  $-\gamma'(q)$  at the point  $q$  (as measured by the metric  $g_{T,q}$ ). By reducing the constant  $c(\varepsilon)$  if necessary, the claim is proved.

Let  $\tau$  be the Lorentzian distance from  $q$ : it is defined on  $\mathbf{exp}_q(C_q(r_0))$  and is a smooth function on  $\mathbf{exp}_q(C_q(r_0)) \setminus \{p\}$ . Using the claim (5.1) we deduce that  $\tau$  is smooth in the ball  $\mathcal{B}_T(p, c(\varepsilon)r_0)$  and satisfies

$$\left(\frac{1}{2} - c(\varepsilon)\right)r_0 < \tau < \left(\frac{1}{2} + c(\varepsilon)\right)r_0 \quad \text{in the ball } \mathcal{B}_T(p, c(\varepsilon)r_0). \quad (5.2)$$

It is clear also that

$$|\nabla\tau|_g^2 = -1, \quad \nabla^2\tau(\nabla\tau, \cdot) = 0.$$

We now introduce a new foliation. Let  $(z^i)$  be coordinates on the level set hypersurface  $\tau = \tau(p)$ , and by following the integral curves of the (unit, time-like) vector field

$$N := \nabla\tau$$

let us construct coordinates  $(z^\alpha)$  with  $z_0 := \tau$  in which the Lorentzian metric  $g$  takes the simple form

$$g = -(dz^0)^2 + g_{ij} dz^i dz^j.$$

Let  $g_N = \langle \cdot, \cdot \rangle_N$  be a (new) reference Riemannian metric based on the vector field  $N$ .

By Lemma 3.2 using the equation satisfied by (future)  $g$ -geodesics  $\sigma$  we see that

$$\left| \frac{d}{d\tau} \log |\sigma'(\tau)| \right| \leq K_3 r_0.$$

(Recall that we allow  $r_0$  to depend upon  $\varepsilon$ .) This inequality shows that the vector field  $N$  makes an angle  $\leq c(\varepsilon)$  with  $T$ , everywhere on  $\mathbf{exp}_q(C_q(r_0, \varepsilon))$ . From this, we conclude that the two metrics are comparable:

$$(1 - c(\varepsilon))g_T \leq g_N \leq (1 + c(\varepsilon))g_T \quad \text{in the cone } \mathbf{exp}_q(C_q(r_0, \varepsilon)).$$

*Step 2. Hessian comparison theorem and curvature bound for the reference metric  $g_N$ .* Since  $p \in \mathbf{exp}_q(C_q(r_0))$ , let  $\sigma : [0, \tau(p)] \rightarrow M$  be the future time-like geodesic connecting  $q$  to  $p$ , and let  $\tilde{V}$  be the Jacobi field defined along  $\sigma$  such that

$$\tilde{V}(0) = 0, \quad \tilde{V}(\tau(p)) = V,$$

where  $V \in T_p M$  satisfies the orthogonality condition  $\langle \nabla\tau, V \rangle = 0$ . Then we have

$$\begin{aligned} -\nabla^2\tau(V, V) &= -\langle \tilde{V}, \nabla_{\nabla\tau} \tilde{V} \rangle = \langle \tilde{V}, \nabla_{\frac{\partial}{\partial\tau}} \tilde{V} \rangle \\ &= \int_0^{\tau(p)} \langle \nabla_{\frac{\partial}{\partial\tau}} \tilde{V}, \nabla_{\frac{\partial}{\partial\tau}} \tilde{V} \rangle_g - R(\sigma', \tilde{V}, \sigma', \tilde{V}) =: I(\tilde{V}, \tilde{V}). \end{aligned}$$

Recall that in the absence of conjugate points Jacobi fields minimize the index form  $I(V, V)$  among all vector fields with fixed boundary values. By applying a standard comparison technique from Riemannian geometry on the orthogonal space  $(\nabla\tau)^\perp$  (on which the Lorentzian metric induces a Riemannian metric) we control the Hessian of  $\tau$  in terms of the curvature bound  $K_2$ :

$$\frac{\sqrt{K_2(1+c(\varepsilon))}}{\tan \sqrt{K_2(1+c(\varepsilon))}\tau} g|_{(\nabla\tau)^\perp} \leq (-\nabla^2\tau)|_{(\nabla\tau)^\perp} \leq \frac{\sqrt{K_2(1+c(\varepsilon))}}{\tanh \sqrt{K_2(1+c(\varepsilon))}\tau} g|_{(\nabla\tau)^\perp}. \quad (5.3)$$

Since  $-\nabla_{ij}^2\tau = \frac{1}{2}\frac{\partial g_{ij}}{\partial\tau}$ , we deduce from (5.3) that

$$\frac{g_{ij}}{\tau} \leq \frac{\partial g_{ij}}{\partial\tau} \leq \frac{3g_{ij}}{\tau} \quad \text{in the cone } \mathbf{exp}_q(C_q(r_0)). \quad (5.4)$$

Combining (5.4) with the curvature formulas derived in Section 3, i.e.

$$\begin{aligned} R_{ijkl} &= R_{ijkl}^\Sigma + \frac{1}{4} \left( \frac{\partial g_{ik}}{\partial\tau} \frac{\partial g_{jl}}{\partial\tau} - \frac{\partial g_{il}}{\partial\tau} \frac{\partial g_{jk}}{\partial\tau} \right), \\ R_{0jil} &= \frac{1}{2} \left( \nabla_l \left( \frac{\partial}{\partial\tau} g_{ij} \right) - \nabla_i \left( \frac{\partial}{\partial\tau} g_{lj} \right) \right), \\ R_{i0j0} &= -\frac{1}{2} \frac{\partial^2 g_{ij}}{\partial\tau^2} + \frac{1}{4} g^{pq} \frac{\partial g_{ip}}{\partial\tau} \frac{\partial g_{jq}}{\partial\tau}, \end{aligned}$$

we conclude that

$$\left| \frac{\partial^2 g_{ij}}{\partial\tau^2} \right| \leq \frac{C}{\tau^2} \quad \text{on } \mathbf{exp}_q(C_q(r_0)). \quad (5.5)$$

Finally, relying on the formulas for the curvature of the reference Riemannian metric  $g_N$ , we obtain

$$|\mathbf{Rm}_{g_N}|_N \leq \frac{C}{\tau^2} \quad \text{on } \mathbf{exp}_q(C_q(r_0)).$$

(Observe that, as could have been expected, the upper bound blows-up as one approach the point  $q$  which is the base point in our definition of the distance.) In particular, this implies the following curvature bound near the point  $p$ :

$$|\mathbf{Rm}_{g_N}|_N \leq Cr_0^{-2} \quad \text{on the ball } \mathcal{B}_T(p, c(\varepsilon)r_0).$$

*Step 3. Constructing geodesically convex functions.* Since the metrics  $g_T$  and  $g_N$  are comparable, the volume ratio  $(1/r_0^{n+1}) \mathbf{Vol}_{g_N} \mathcal{B}_N(p, c(\varepsilon)r_0)$  is uniformly bounded (above and) below. By applying the theory for Riemannian metrics in [10], the injectivity radius of the metric  $g_N$  is bounded from below by  $c(\varepsilon)r_0$ . Let

$$u(x) := d_{g_N}(p, x)^2$$

be the (square) of the distance function associated with the Riemannian metric  $g_N$ , which is a smooth function defined on the geodesic ball  $\mathcal{B}_N(p, c(\varepsilon)r_0)$ . By the standard Hessian comparison theorem for Riemannian manifold we have

$$(2 - \varepsilon) g_{N,\alpha\beta} \leq \nabla_{g_N,\alpha} \nabla_{g_N,\beta} u \leq (2 + \varepsilon) g_{N,\alpha\beta} \quad \text{on the ball } \mathcal{B}_N(p, c(\varepsilon)r_0).$$

In terms of the original Lorentzian metric  $g$ , the Hessian of the function  $u$  is

$$\nabla_\alpha \nabla_\beta u = \nabla_{g_N,\alpha} \nabla_{g_N,\beta} u + (\Gamma_{g_N,\alpha\beta}^\gamma - \Gamma_{g,\alpha\beta}^\gamma) \frac{\partial u}{\partial x^\alpha}.$$

Since  $|\Gamma_{g_N} - \Gamma|_N \leq C \sup |\frac{\partial g_{ii}}{\partial \tau}| \leq C'$  by the estimate (5.4) and since also  $|\nabla u|_N \leq 2d_{g_N}$  on  $\mathcal{B}_N(p, r_0)$ , we conclude that

$$(2 - \varepsilon) g_{N,\alpha\beta} \geq \nabla_\alpha \nabla_\beta u \geq (2 + \varepsilon) g_{N,\alpha\beta} \quad \text{in the ball } \mathcal{B}_N(p, c(\varepsilon)r_0).$$

This completes the proof of Theorem 5.1.  $\square$

## 6 Injectivity radius of null cones

We turn our attention now to null cones within foliated Lorentzian manifolds. Our main result (Theorem 6.1 below) provides a lower bound for the null injectivity radius under the main assumption that the exponential map is defined in some ball and the null conjugate radius is already controled. Hence, contrary to the presentation in Section 3 our main assumption (see (A3') below) is not directly stated as a curvature bound. However, under additional assumptions, it is known that the conjugate radius estimate can be deduced from an  $L^p$  curvature bound, so that our result is entirely relevant for the applications.

Indeed, in a series of fundamental papers [17, 18, 19], Klainerman and Rodnianski assumed on an  $L^2$  curvature bound and estimated the null conjugate and injectivity radii for Ricci-flat Lorentzian  $(3+1)$ -manifolds. Our result in the present section is a continuation of the recent work [19] and covers a general class of Lorentzian manifolds with arbitrary dimension, while our proof is local and geometric and so conceptually simple.

We use the terminology and notation introduced in Section 2. In particular, a point  $p \in M$  and a reference vector field  $T$  are given, and  $N_p^-$  denotes the past null cone in the tangent cone at  $p$ . The null exponential map  $\mathbf{exp}_p^N : B_T^N(0, r) \rightarrow M$  is defined over a subset of this cone,

$$B_T^N(0, r) := B_T(0, r) \cap N_p^-,$$

and allows us to introduce the (past) null injectivity radius  $\mathbf{Null Inj}_g(M, p, T)$ .

We also set

$$\mathcal{B}_T^N(p, r) := \mathbf{exp}_p^N(B_T^N(0, r)).$$

We consider a domain  $\Omega \subset M$  containing some point  $p$  on a final slice  $\Sigma_0$  and foliated as

$$\Omega = \bigcup_{t \in [-1, 0]} \Sigma_t, \quad p \in \Sigma_0. \quad (6.1)$$

We assume that there exist positive constants  $K_0, K_1, K_2$  such that

$$e^{-K_0} \leq n \leq e^{K_0} \quad \text{in } \Omega, \quad (\text{A1})$$

$$|\mathcal{L}_T g|_T \leq K_1 \quad \text{in } \Omega, \quad (\text{A2})$$

the null conjugate radius at  $p$  is  $r_0$  (at least) and the null exponential map satisfies

$$e^{-K_2} g_{T,p} |_{\mathbb{B}_T^N(0,r_0)} \leq \left( \exp_p^N \right)^* (g_T |_{\mathbb{B}_T^N(0,r_0)}) \leq e^{K_2} g_{T,p} |_{\mathbb{B}_T^N(0,r_0)} \quad (\text{A3}')$$

and, finally, there exists a coordinate system on the initial slice  $\Sigma_{-1}$  such that the metric  $g |_{\Sigma_{-1}}$  is comparable to the  $n$ -dimensional Euclidian metric  $g_{E'}$  in these coordinates:

$$e^{-K_0} g_{E'} \leq g |_{\Sigma_{-1}} \leq e^{K_0} g_{E'} \quad \text{in } \mathbb{B}_{\Sigma_{-1}, E'}(p, r_0). \quad (\text{A4}')$$

We refer to  $K_2$  as the *effective conjugate radius constant*.

**Theorem 6.1** (Injectivity radius of null cones). *Let  $M$  be a differentiable  $(n + 1)$ -manifold, endowed with a Lorentzian metric  $g$  satisfying the regularity assumptions (A1), (A2), (A3'), and (A4') at some point  $p$  and for some foliation (3.1). Then, there exists a positive constant  $i_0$  depending only upon the foliation bounds  $K_0, K_1$ , the null conjugate radius  $r_0$ , the effective conjugate radius constant  $K_2$ , and the dimension  $n$  such that the null injectivity radius of the metric  $g$  at  $p$  satisfies*

$$\text{Null Inj}_g(M, p, T) \geq i_0.$$

It is interesting to compare the assumptions above with the ones in Section 3. Assumptions (A1) and (A2) are concerned with the property of the foliation and were already required in Section 3.

Assumption (A3') should be viewed as a weaker version of the  $L^\infty$  curvature condition (A3). Recall that, under the assumptions of Theorem 3.1 which included a curvature bound, an analogue of (A3') valid in the whole of  $\Omega$  was already established in (4.8). It is expected that (A3') is still valid when the curvature in every spacelike slice is solely bounded in some  $L^m$  space.

Indeed, at least when the spatial dimension is  $n = 3$  and the manifold is Ricci-flat, according to Klainerman and Rodnianski [17, 18] Assumption (A3') is a consequence of the following  $L^2$  curvature bound

$$\|\mathbf{Rm}_g\|_{L^2(\Sigma_{-1}, g_T)} \leq K'_2 \quad (6.2)$$

for some constant  $K'_2 > 0$ .

Assumption (A4') concerns the metric on the initial hypersurface and is only slightly stronger than the volume bound (A4). Furthermore, according to Anderson [1] and Petersen [22] the property (A4') is also a consequence of the curvature bound

$$\|\mathbf{Rm}_g\|_{L^m(\Sigma_{-1}, g_T)} \leq K'_2 \quad (6.3)$$

for  $m > n/2$  and some constant  $K'_2 > 0$  and a volume lower bound at every scale

$$r^{-n} \mathbf{Vol}_{g|_{\Sigma_0}}(\mathbb{B}_{\Sigma_0}(p, r)) \geq v_0, \quad r \in (0, r_0]. \quad (6.4)$$

In summary, by combining Theorem 6.1 above with the results in [19, 16] we conclude:

**Corollary 6.2** (Einstein field equations of general relativity). *Let  $(M, g)$  be a Lorentzian  $(3 + 1)$ -manifold satisfying the vacuum Einstein equation*

$$\mathbf{Ric}_g = 0. \quad (6.5)$$

*Suppose that near some point  $p \in M$  there exists a foliation  $\Omega$  of the form (6.1) satisfying Assumptions (A1)-(A2) and such that the  $\mathbf{L}^2$  curvature assumption (6.2) holds on the initial spacelike hypersurface  $\Sigma_{-1}$ . Then, there exists a positive constant  $i_0$  depending only upon the foliation bounds  $K_0, K_1$  and the curvature bound  $K'_2$  such that the null injectivity radius satisfies*

$$\mathbf{Null Inj}_g(M, p, T) \geq i_0.$$

*Proof of Theorem 6.1. Step 1. Localization of the past null cone  $\mathcal{N}^-(p)$  between two flat null cones.* Assumption (A3') provides us with a bound on the null conjugate radius, we need to control the null cut locus radius. We proceed as in Section 4 and introduce coordinates near the point  $p$  such that  $x^\alpha(p) = 0$ . Precisely, relying on Assumptions (A1), (A2), and (A4'), we determine the coordinates  $x = (x^\alpha)$  so that  $x^0 = t$  and the spatial coordinates  $(x^j)$  are transported (via the gradient of the function  $t$ ) from the coordinates prescribed on the initial slice  $\Sigma_{-1}$ . Then, the Lorentzian metric reads  $g = -n^2 dt^2 + g_{ij} dx^i dx^j$  and satisfies for some  $C_0, C_1 > 0$

$$\frac{1}{C_0} \leq n^2 \leq C_0, \quad \frac{1}{C_1} \delta_{ij} \leq g_{ij} \leq C_1 \delta_{ij}, \quad (6.6)$$

for all  $-r_0 < t \leq 0$  and  $(x^1)^2 + \dots + (x^n)^2 \leq (r_0)^2$ , and in these coordinates the reference Riemannian metric  $g_T$  is comparable to the  $(n + 1)$ -dimensional Euclidian metric  $g_E := dt^2 + (dx^1)^2 + \dots + (dx^n)^2$ :

$$\frac{1}{C_1} g_E \leq g_T \leq C_1 g_E. \quad (6.7)$$

Denote by  $\mathcal{B}_E(q, r)$  the Euclidean ball with center  $q$  and radius  $r$ . Note that these inequalities holds within a neighborhood of  $p$  in  $\Omega$ . The forthcoming bounds will hold in a neighborhood of the past null cone only. To simplify the notation we set

$$c_0 := \frac{1}{C_0}, \quad c_1 := \frac{1}{C_1}.$$

In each time slice of parameter value  $t = a$  we introduce the  $n$ -dimensional Euclidian ball with radius  $b$

$$\mathcal{A}_{<b}^a := \left\{ t = a, \quad (x^1)^2 + \dots + (x^n)^2 < b^2 \right\} \subset \Sigma_a,$$

which is centered around the point  $p'$  with coordinates  $(a, 0, \dots, 0)$ . We also define  $\mathcal{A}_{>b}^a, \mathcal{A}_{[c,d]}^a \dots$  similarly.

For any point  $q$  in a slice  $\Sigma_{t_0}$  satisfying  $-r_0 \leq t_0 < 0$  and  $x^1(q)^2 + \dots + x^n(q)^2 < c_1^2 t_0^2$  we consider the line (for the Euclidian metric) connecting  $q$  to  $p$ :

$$\gamma(\tau) = \left( \tau, \frac{\tau}{t_0} x^1(q), \dots, \frac{\tau}{t_0} x^n(q) \right), \quad \tau \in [t_0, 0].$$



This is a timelike curve for the Lorentzian metric  $g$ , since

$$|\gamma'(\tau)|^2 = -n^2 + g_{ij} \frac{x^i(q)}{t_0} \frac{x^j(q)}{t_0} < -c_0 + c_1 < 0,$$

which shows that

$$\mathcal{A}_{<c_1|t|}^t \subset \mathcal{J}^-(p), \quad t \in (-r_0, 0).$$

On the other hand, we claim that the larger Euclidian cone  $\mathcal{A}_{<c_1|t|}^t$  contains the null cone, in other words

$$\mathcal{A}_{\geq c_1|t|}^t \subset (\mathcal{N}^-(p) \cup \mathcal{J}^-(p))^c, \quad t \in (-c_1 r_0, 0).$$

Indeed, arguing by contradiction we suppose there exist a time  $t_0 \in (-c_1 r_0, 0)$  and a point  $q \in \mathcal{A}_{\geq c_1 t_0}^{t_0}$  connected to  $p$  by a causal curve  $\gamma = \gamma(s)$  with  $\gamma(0) = p$ . After reparametrizing (in time) the curve is necessary we can assume that  $\gamma(\tau) = (\tau, x^j(\tau))$  for some  $t'_0 \leq \tau \leq 0$ , as long as the point  $\gamma(\tau)$  lies in the coordinate system under consideration. For this part of the curve at least we have

$$0 \geq |\gamma'|^2 = -n^2 + g_{ij} \frac{dx^i}{d\tau} \frac{dx^j}{d\tau},$$

which by (6.6) implies that  $(\frac{dx^1}{d\tau})^2 + \dots + (\frac{dx^n}{d\tau})^2 < C_1 C_0$ . Therefore, after integration we find

$$(x^1(q)^2 + \dots + x^n(q)^2)^{1/2} (t'_0) \leq \sqrt{C_0 C_1} t'_0 \leq \sqrt{C_0 c_1} r_0 < r_0.$$

Hence, we can choose  $t'_0 = t_0$ , the whole curve lies in the system of coordinates, and is parametrized in the form  $\gamma(\tau) = (\tau, x(\tau))$ , ( $\tau \in [t_0, 0]$ ). Moreover, we have  $|x(t_0)| \leq \sqrt{C_1 C_0} |t_0| < C_1 |t_0|$ , which contradicts our assumption  $q \in \mathcal{A}_{\geq c_1 t_0}^{t_0}$ .

In conclusion, we have localized the slices of the past null cone within "annulus" regions:

$$\mathcal{N}^-(p) \cap \Sigma_t \subset \mathcal{A}_{[c_1|t|, C_1|t|]}^t, \quad t \in [-c_1 r_0, 0].$$

*Step 2. The past null cone  $\mathcal{N}^-(p)$  viewed as a graph with bounded slope.* We now obtain a Lipschitz continuous parametrization of the null cone. For any fixed  $q \in \mathcal{A}_{\leq c_1^2 r_0}^{-c_1 r_0}$  we consider the vertical curve passing through  $q$ :

$$\gamma_q(\tau) = (\tau, x^1(q), \dots, x^n(q)), \quad \tau \in [-c_1 r_0, 0].$$

By Step 1 we know that there exists  $\tau_q$  such that  $\gamma_q(\tau_q) \in \mathcal{N}^-(p)$ . Moreover,  $\tau_q$  is unique since  $\mathcal{N}^-(p)$  is achronal, and this defines a map

$$F : \mathcal{A}_{\leq c_1^2 r_0}^{-c_1 r_0} \rightarrow \mathcal{N}^-(p)$$

such that  $F(q) = \gamma_q(\tau_q)$ . It is obvious  $F(-c_1 r_0, 0) = p$ .

We claim that the map  $F$  is Lipschitz continuous with Lipschitz constant less than  $C_1$ , as computed with the Euclidean metric  $E$ . Namely, by contradiction, suppose that  $|F(q_1) - F(q_2)|_E > C_1 |q_1 - q_2|_E$  for some  $q_1, q_2 \in \mathcal{A}_{\leq c_1^2 r_0}^{-c_1 r_0}$ , then by (6.7) in Step 1,  $F(q_1)$  would be chronologically related to  $F(q_2)$  and this would contradict the fact that  $\mathcal{N}^-(p)$  is achronal. Moreover, from Step 1 it follows that

$$F(\mathcal{A}_{\leq c_1^2 r_0}^{-c_1 r_0}) \supset \mathcal{N}^-(p) \cap \mathcal{B}_E(p, c_1^3 r_0).$$

*Step 3. Constructing an homotopy of curves on the null cone.* Suppose that  $\gamma_1, \gamma_2$  are two (past) null geodesics from  $p$  satisfying

$$\begin{aligned} \gamma_1(0) &= \gamma_2(0), & |\gamma_1'(0)|_T &= |\gamma_2'(0)|_T = 1, \\ \gamma_1(s_1) &= \gamma_2(s_2). \end{aligned}$$

We claim that  $\max(s_1, s_2) > c_1^6 r_0$ , which will establish the desired injectivity bound by setting  $i_0 = c_1^6 r_0$ .

We argue by contradiction and assume that  $\max(s_1, s_2) < c_1^6 r_0$ . Taking into account Assumption (A2) and applying exactly the same arguments as in Step 1 of Section 4 we see that the  $g_T$ -lengths of the curves  $\gamma_1, \gamma_2$  satisfy

$$L(\gamma_j, g_T) \leq s_j e^{CC_1 s_j} \leq c_1^{5+3/4} r_0 \quad (j = 1, 2).$$

By Step 1 of the present proof we know that the Euclidean lengths of  $\gamma_1, \gamma_2$  satisfy

$$L(\gamma_j, g_E) \leq c_1^{5+1/4} r_0 \quad (j = 1, 2).$$

In particular,  $\gamma_1, \gamma_2 \subset \mathcal{N}^-(p) \cap \mathcal{B}_E(p, c_1^5 r_0)$  and we can thus concatenate the curve  $\gamma_1, \gamma_2$  and obtain

$$\gamma := \gamma_2^{-1} \cup \gamma_1 : [0, s_1 + s_2] \rightarrow \mathcal{N}^-(p) \cap \mathcal{B}_E(p, c_1^5 r_0).$$

Since  $F(\mathcal{A}_{\leq c_1^2 r_0}^{-c_1 r_0}) \supset \mathcal{N}^-(p) \cap \mathcal{B}_E(p, c_1^3 r_0)$ , there exists a smooth family of curves  $\sigma_\varepsilon : [0, s_1 + s_2] \rightarrow \mathcal{N}^-(p)$  such that

$$\begin{aligned} \sigma_1 &= \gamma, & \sigma_0 &= p, \\ \sigma_\varepsilon(0) &= \sigma_\varepsilon(s_1 + s_2) = p, & \varepsilon &\in [0, 1]. \end{aligned}$$

Specifically, we choose

$$\sigma_\varepsilon(s) := F(\varepsilon F^{-1} \gamma(s)),$$

where the multiplication by  $\varepsilon$  is defined by relying on the linear structure of  $\mathcal{A}_{\leq c_1^2 r_0}^{-c_1 r_0} \approx B_{\mathbb{R}^n}(0, c_1^2 r_0)$ . Equivalently, by setting  $x^i(s) = x^i(\gamma(s))$  we have the explicit formula

$$\sigma_\varepsilon(s) = F\left(-c_1 r_0, \varepsilon x^1(s), \dots, \varepsilon x^n(s)\right).$$

It is clear that the Euclidean and  $g_T$ -lengths of  $\sigma_\varepsilon$  satisfy

$$\begin{aligned} L(\sigma_\varepsilon, g_E) &\leq \varepsilon(1 + C_1)L(\gamma, g_E) \leq c_1^{4+1/8} r_0, \\ L(\sigma_\varepsilon, g_T) &\leq c_1^{3+5/8} r_0. \end{aligned}$$

By Assumption (A3') on the null conjugate radius, we can lift to the null cone of the tangent space  $T_p M$  the continuous family of loops  $\sigma_\varepsilon$ , and we obtain a continuous family of curves  $\widetilde{\sigma}_\varepsilon$  defined on  $[0, s_1 + s_2]$  such that

$$\widetilde{\sigma}_\varepsilon(0) = 0, \quad L(\widetilde{\sigma}_\varepsilon, g_{T,p}) \leq c_1^3 r_0.$$

Observe that the property  $L(\widetilde{\sigma}_\varepsilon, g_{T,p}) \leq c_1^3 r_0 \leq r_0$  guarantees the existence of this continuous lift. By continuity, all of the curves  $\widetilde{\sigma}_\varepsilon$  are loops containing 0. As observed earlier in the proof for the case of bounded curvature,  $\widetilde{\sigma}_1$  consists of two distinct segments which clearly can not form a closed loop and we have reached a contradiction.  $\square$

## 7 Injectivity radius of an observer in a Lorentzian manifold

### Main result

We are now in a position to discuss and prove Theorem 1.1 stated in the introduction. As we have seen in the proof of the previous section, once the injectivity radius is controlled, one can construct a foliation satisfying certain “good” properties. On the other hand, the concept of injectivity radius is clearly independent of any prescribed foliation. As this is more natural, we will now present a general result which avoids to assume a priori the existence of a foliation. This will be achieved by relying on purely geometric and intrinsic quantities and constructing coordinates adapted to the geometry. Such a result is conceptually very important in the applications. The result and proof in this section should be viewed as a Lorentzian generalization of Cheeger, Gromov, and Taylor’s technique [10], originally developed for Riemannian manifolds.

Let  $(M, g)$  be a differentiable  $(n + 1)$ -manifold endowed with a Lorentzian metric tensor  $g$ , and consider a point  $p \in M$  and a vector  $T \in T_p M$  with  $g_p(T, T) = -1$ . That is, we now fix a single observer located at the point  $p$ . As explained in Section 2 the vector  $T$  induces an inner product  $g_T = \langle \cdot, \cdot \rangle_T$  on the tangent space  $T_p M$ . We assume that the exponential map  $\mathbf{exp}_p$  is defined in some ball  $B_T(0, r_0) \subset T_p M$  determined by this inner product, which is of course always true in a sufficiently small ball. Controlling the geometry at the point  $p$  precisely amounts to estimating the size of this radius  $r_0$  where the exponential map is defined and has some good property. We restrict attention to the geodesic ball  $\mathcal{B}_T(p, r_0) := \mathbf{exp}_p(B_T(0, r_0))$ ; recall that these sets depend upon the vector  $T$  given at  $p$ .

As explained in the introduction, by  $g$ -parallel translating the vector  $T$  at  $p$  along a geodesic  $\gamma$  from  $p$ , we can define get a future-oriented unit time-like vector field  $T - \gamma$  defined along this geodesic. To this vector field and the Lorentzian metric  $g$  we can associate a reference Riemannian metric  $g_{T_\gamma}$  along the geodesic. In turn, this allows us to compute the norm  $|\mathbf{Rm}_g|_{T_\gamma}$  of the Riemann curvature tensor along the geodesic.

Of course, whenever two such geodesics  $\gamma, \gamma'$  meet away from  $p$ , the corresponding vectors  $T_\gamma$  and  $T_{\gamma'}$  are generally *distinct*. If we consider the family of all such geodesics we therefore obtain a (generally) multi-valued vector field defined in the geodesic ball  $\mathcal{B}_T(p, r_0)$ . We use the same letter  $T$  to denote this vector field. In turn we can still compute the Riemann curvature norm  $|\mathbf{Rm}_g|_T$  by taking into account every value of  $T$ .

The key objective of the present section is the study of the geometry of the local covering  $\mathbf{exp}_p : B_T(0, r_0) \rightarrow \mathcal{B}_T(p, r_0)$  and to compare the Lorentzian metric  $g$  defined on the manifold  $M$  with the reference Riemannian metrics  $g_T$ . As we will see in the proof below, it will be convenient to pull the metric “upstairs” on the tangent space at  $p$ , using the exponential map. Indeed, this will be possible once we will have estimated the conjugate radius (in Step 1 of the proof below) and will know that the exponential map is non-degenerate on  $B_T(0, r_0)$ . Pulling back the Lorentzian metric  $g$  on  $M$  by the exponential map we get a Lorentzian metric  $g = \mathbf{exp}_p^* g$  defined in the tangent space, on the ball  $B_T(0, r_0)$ . We use the same letter  $g$  to denote this metric. Then, the geometry in the tangent space is particularly simple, since the  $g$ -geodesics on  $M$  passing through  $p$  are radial straightline in  $B_T(0, r_0)$ .

A third view point could be adopted by restricting attention within the cut-locus from the point  $p$ , and by imposing the curvature assumption within the cut-locus only.

We are in a position to prove the main result of the present paper that was stated in Theorem 1.1.

*Proof of Theorem 1.1.* After scaling we may assume that  $r_0 = 1$ , and so we need to show

$$\mathbf{Inj}_g(M, p, T) \geq c(n) \mathbf{Vol}_g(\mathcal{B}_T(p, c(n))). \quad (7.1)$$

*Step 1. Estimates for the metric  $g_T$  and its covariant derivative.* Let  $E_0 = T, E_1, \dots, E_n$  be an orthonormal frame at the origin in  $T_p M$  for the Lorentzian metric  $g$ . By  $g$ -parallel transporting this basis along along a radial geodesic  $\gamma = \gamma(r)$ , satisfying  $\gamma(0) = 0, |\gamma'(0)|_T = 1$ , we get an orthonormal frame defined along the geodesic. We use the same letters  $E_\alpha$  to denote these vector fields. Since

$$\frac{d}{dr} \langle E_\alpha, E_\beta \rangle_g = 0,$$

we infer that

$$|E_i|_T^2 = |E_i|_g^2 = 1 \quad \text{along the geodesic.}$$

The same argument also implies

$$|\gamma'(r)|_T^2 = |\gamma'(0)|_T^2 = 1, \quad |\gamma'(r)|_g^2 = |\gamma'(0)|_g^2 = 1, \quad (7.2)$$

and  $\gamma'(r) = c^\alpha E_\alpha(r)$  with constant (in  $r$ ) scalars  $c^\alpha$  and  $\sum |c^\alpha|^2 = |\gamma'(0)|_T = 1$ . We used here that, by definition,  $\gamma'$  is  $g$ -parallel transported.

Let  $V = a^\alpha(r) E_\alpha(r)$  be a Jacobi field along a radial geodesic  $\gamma = \gamma(r)$ , with  $V(0) = 0$  and  $|V'(0)|_T = 1$ . Then, the Jacobi equation takes the form

$$(a^\alpha)''(r) = -\langle E_\alpha, R(E_\beta, E_\gamma) E_\delta \rangle_T c^\beta c^\delta a^\gamma(r).$$

Since

$$-2 \sum_\alpha (a_\alpha'^2 + a_\alpha^2) \leq \frac{d}{dr} \left( \sum_\alpha a_\alpha'^2 + a_\alpha^2 \right) \leq 2 \sum_\alpha (a_\alpha'^2 + a_\alpha^2),$$

we obtain  $|V'(r)|_T \leq e^r$  and thus  $|V(r)|_T \leq (e^r - 1)$ .

By substituting this result into the above formulas, the estimate can be improved again. Indeed, by computing and estimating the second-order derivative  $\frac{d}{dr} \sum_\alpha a_\alpha' a_\alpha$  as we did for the Jacobi field estimate of Section 4, we can check that

$$r - C(n) r^2 \leq \left( \sum |a_\alpha|^2(r) \right)^{1/2} \leq (e^r - 1) \quad \text{along the geodesic.}$$

Denote by  $g_0, g_{T,0}$  the Lorentzian and the Riemannian metrics at the origin 0 (which are nothing but the metrics at the point  $p$ ), and let  $y^0, \dots, y^n$  be Cartesian coordinates on  $B_T(0, 1)$ , with  $\langle \frac{\partial}{\partial y^\alpha}, \frac{\partial}{\partial y^\beta} \rangle_{g_0}(0) = \eta_{\alpha\beta}$ . Assuming that the radius under consideration is sufficiently small so that  $(1 - C(n)|y|) < 1$  we conclude from the Jacobi field estimate that the exponential map is non-degenerate and that the metric along the geodesic are comparable. In turn, since this is true for every radial geodesic, we can define the pull back of the metric to the tangent space and the conclusion hold in the whole ball  $B_T(0, 1)$ , that is

$$(1 - C(n)|y|) g_{T,0} \leq g_{T,y} \leq (1 + C(n)|y|) g_{T,0}, \quad y \in B_T(0, 1). \quad (7.3)$$

By construction of the metric  $g_T$  we have  $\nabla_{g_T} - \nabla_g = \nabla_g T * T$  (schematically) and  $\nabla T(0) = 0$ , and it is useful to control the covariant derivative too. To this end, write the radial vector field as

$$\frac{\partial}{\partial r} = \frac{y^\alpha}{r} \frac{\partial}{\partial y^\alpha}, \quad r := \left( \sum |y^\alpha|^2 \right)^{1/2},$$

with  $|\frac{\partial}{\partial r}|_T^2 \equiv 1$  (as stated already in (7.2)). Using that  $|\nabla T|_T^2 = \nabla_\alpha T^\xi \nabla_\beta T^\eta g_{T,\xi\eta} g_T^{\alpha\beta}$  and computing the derivative of  $|\nabla T|_T^2$  along radial geodesics, we find

$$\frac{d}{dr} |\nabla T|_T^2 \leq C(n) |\nabla T|_T^3 + 2 \langle \nabla_{\frac{\partial}{\partial r}} \nabla T, \nabla T \rangle_T.$$

By using that

$$\nabla_{\frac{\partial}{\partial r}} T = 0, \quad \left[ \frac{\partial}{\partial r}, \frac{\partial}{\partial y^\alpha} \right] = -\frac{1}{r} \frac{\partial}{\partial y^\alpha} + \frac{y^\alpha}{r^2} \frac{\partial}{\partial r},$$

we obtain

$$\nabla_{\frac{\partial}{\partial r}} \nabla_{\frac{\partial}{\partial y^\alpha}} T^\gamma = -\frac{1}{r} \nabla_{\frac{\partial}{\partial y^\alpha}} T^\gamma + R\left(\frac{\partial}{\partial r}, \frac{\partial}{\partial y^\alpha}\right) T^\gamma,$$

and therefore, thanks to the curvature assumption,

$$\frac{d}{dr} |\nabla T|_T^2 \leq -\frac{2}{r} |\nabla T|_T^2 + C(n) |\nabla T|_T^3 + C(n) |\nabla T|_T.$$

This implies the following bound for the covariant derivative

$$|\nabla T|_T(y) \leq C(n) |y|, \quad |y| \leq 1/C(n), \quad (7.4)$$

which also provides a bound for the difference  $\nabla_{g_T} - \nabla_g$ .

*Step 2. Estimate of the injectivity radius of  $g$  on  $B_T(0, c(n))$ .*

Since the curvature on  $B_T(0, 1)$  is bounded and that  $|\nabla_{g_T} - \nabla_g|_T^2 \leq C(n) = 1/c(n)$  on the ball  $B_T(0, c(n))$  we can follow the argument in Section 4 and bound from below the conjugate radius for any point in the ball  $B_T(0, 3c(n)/4)$ .

Next, given any point  $y \in B_T(0, c(n)/2)$ , let  $\gamma_1$  and  $\gamma_2$  be two geodesics which meet at their end points and have “short” length with respect to the metric  $g_T$  (or  $g_{T,0}$ ). By using the linear structure on  $B_T(0, 1)$  (a subset of the vector space  $T_p M$ ) we can construct an homotopy of the loop  $\gamma_1 \cup \gamma_2^{-1}$  to the origin, such that each curve have also “short” length for the metric  $g_T$ . By lifting the homotopy to the tangent space  $T_y B_T(0, 1)$  and by relying on the conjugate radius bound, we reach a contradiction as was done in Section 4.

In summary, there exists a universal constant  $C(n) = 1/c(n)$  depending only on the dimension such that the injectivity radius at each point  $y$  of  $B_T(0, c(n))$  is bounded from below by  $4c(n)$ . Moreover, by the Jacobian field estimate again, we can prove the ball  $B_{T,p}(0, c(n)) \subset T_p M$  defined by the Euclidean metric  $g_{T,p}$  is covered by  $\exp_y(B_{T,y}(0, 3c(n)))$ , where  $B_{T,y}(0, 3c(n)) \subset T_y T_p M$  is a ball of radius  $3c(n)$  defined by metric  $g_{T,y}$ , and any two points in  $B_{T,p}(0, c(n))$  can be connected by a  $g$  geodesic which is totally contained in  $B_{T,p}(0, 2c(n))$ . Further arguments are now required to arrive at the desired bound (7.1).

*Step 3. Riemannian metric  $g_N$  induced on  $B_T(0, 2c(n))$ .* Consider a geodesic  $\gamma$  satisfying  $\gamma(0) = 0$  and  $\gamma'(0) = -T$ , and let us set

$$\gamma(c(n)/2) =: q, \quad \tau := d_g(\cdot, q) - d_g(q, 0).$$

Then, by following exactly the same arguments as in the main proof of Section 5, we construct a normal coordinate system (of definite size) such that  $g = -d\tau^2 + g_{ij} dx^i dx^j$  and  $g_N = d\tau^2 + g_{ij} dx^i dx^j$ , and such that the corresponding reference Riemannian metric satisfies the following properties:

- (i)  $(1 - c(n)) g_N \leq g_T \leq (1 + c(n)) g_N$ ,
- (ii)  $g_N$  has bounded curvature ( $\leq C(n)$ ) (see (5.5)), and

(iii) for any fixed  $y_0 \in B_T(0, c(n))$  the distance function  $d_{g_N}(y_0, \cdot)^2$  is strictly  $g$ -convex on the ball  $B_T(0, 2c(n))$  and, more precisely,

$$(2 + c(n)) g_N \geq \nabla_g^2 d_{g_N}^2(y_0, \cdot) \geq (2 - c(n)) g_N \quad \text{on } B_T(0, 2c(n))$$

for any  $y_0 \in B_T(0, c(n))$ .

The Hessian of the distance function (defined by the Riemannian metric  $g_N$ ) is naturally computed using the covariant derivatives defined by the Lorentzian metric  $g$ .

*Step 4.* Suppose that  $p_1, \dots, p_N$  are distinct pre-images of  $p$  in the ball  $B_T(0, c(n))$ . We claim that any  $p' \in \mathcal{B}_T(p, c(n))$  has at least  $N$  distinct pre-images in  $B_T(0, 1)$ , and refer to this property as a “lower semi-continuity” property.

Generalizing the terminology in [10], we use the notation  $a \underset{(g_{T,0}, A)}{\sim} b$  when two curves  $a, b$  defined on  $M$  and with the same endpoints are homotopic through a family of curves whose lift have  $g_{T,0}$ -lengths  $\leq A$ . Relying on the lift and the linear structure, we see that, for any curve  $\xi$  starting from  $p$  with (after lifting through 0)  $g_{T,0}$ -length  $A \leq 1$ , there exists a unique  $g$ -geodesic  $\gamma_\xi$  (with the same end points as  $\xi$ ) defined on  $M$  such that  $\xi \underset{(g_{T,0}, A)}{\sim} \gamma_\xi$ . This fact establishes a one-to-one correspondence between the following three concepts:

- (i) equivalence class of curves through  $p$  with  $g_{T,0}$ -lengths  $\leq 3c(n)$ ,
- (ii) radial geodesic segments of  $g_{T,0}$ -lengths  $\leq 3c(n)$ , and
- (iii) points in the ball  $B_T(0, 3c(n)) \subset T_p M$ .

Let  $\sigma$  be a  $g$ -geodesic connecting  $p$  to  $p'$  in  $\mathcal{B}_T(p, c(n))$ . Observe that the images of the lines  $\overline{Op}_i$  by the exponential map,  $\sigma_i = \mathbf{exp}_p(\overline{Op}_i)$ , are distinct geodesic loops through  $p$ . Denote by  $\tilde{\sigma}_i$  the lift of  $\sigma_i \cup \sigma$  through 0, and denote by  $p'_i$  the end point of  $\tilde{\sigma}_i$ . Then, it is clear that all the points  $p'_i$  ( $i = 1, \dots, N$ ) are the pre-images of  $p'$  in  $B_T(0, 1/2)$ . We claim that they are distinct.

Indeed, assuming that  $p'_i = p'_j$  for some  $i \neq j$ , we would find  $\sigma \cup \sigma_i \underset{g_{T,0}, 2c(n)}{\sim} \sigma \cup \sigma_j$ , which gives

$$\sigma_i \underset{g_{T,0}, 3c(n)}{\sim} \sigma^{-1} \cup \sigma \cup \sigma_i \underset{g_{T,0}, 3c(n)}{\sim} \sigma^{-1} \cup \sigma \cup \sigma_j \underset{g_{T,0}, 3c(n)}{\sim} \sigma_j.$$

This would imply  $\sigma_i \underset{g_{T,0}, 3c(n)}{\sim} \sigma_j$  and, therefore,  $p_i = p_j$ , which is a contradiction. In short, this argument shows that the “cancellation law” holds for the homotopy class of “not too long” curves.

*Step 5.* Suppose that there exist two distinct  $g$ -geodesics  $\gamma_1 : [0, l_1] \rightarrow M$  and  $\gamma_2 : [0, l_2] \rightarrow M$  satisfying

$$\gamma_1(0) = \gamma_2(0) = p, \quad |\gamma'_1(0)|_T^2 = |\gamma'_2(0)|_T^2 = 1,$$

and meeting at their endpoints, that is:  $\gamma_1(l_1) = \gamma_2(l_2)$ . Then, let  $l := l_1 + l_2$  and  $\gamma := \gamma_2^{-1} \cup \gamma_1 : [0, l] \rightarrow M$ . Our aim is to prove that

$$l \geq c(n) \mathbf{Vol}_g(\mathcal{B}_T(p, c(n))),$$

which will give us the desired injectivity radius.

From the loop  $\gamma$  we define a map  $\pi_\gamma : B_T(0, c(n)) \rightarrow B_T(0, 2c(n))$  as follows: for any  $y \in B_T(0, c(n))$ , the point  $\pi_\gamma(y)$  is the end point of the lift  $\mathbf{exp}_p(\overline{Oy}) \cup \gamma$  (through the origin). If one would have  $\pi_\gamma(y) = y$  then by the cancellation law established in Step 4, we would have  $\gamma \underset{g_{T,0,2c(n)}}{\sim} 0$ , which is a contradiction. So, the map  $\pi_\gamma$  has no fixed point.

Without loss of generality, we assume that  $l \leq c(n)^5$ . Let  $N = \lfloor c(n)^3/l \rfloor$  be the largest integer less than  $c(n)^3/l$ , and let us use the notation  $2\gamma = \gamma \circ \gamma$ , etc.

**Claim.** The classes  $[\gamma], [2\gamma], \dots, [N\gamma]$  are distinct homotopy classes for the relation  $\underset{g_{T,0,c(n)^2}}{\sim}$ .

If this were not true, then by the cancellation law we would have  $[j\gamma] \underset{g_{T,0,c(n)^2}}{\sim} 0$  for some  $1 \leq j \leq N$ . We already know that all  $\pi_\gamma^i$  is defined from  $B_T(0, c(n)^2)$  to  $B_T(0, c(n))$  for  $i \leq j$ . Since for any  $y \in B_T(0, c(n)^2)$  we have

$$\mathbf{exp}_p(\overline{Oy}) \cup j\gamma \underset{g_{T,0,c(n)}}{\sim} \mathbf{exp}_p(\overline{Oy}),$$

which implies that  $\pi_\gamma^j = id$ . We use here the notation  $\pi_\gamma^2 := \pi_\gamma \circ \pi_\gamma$ , etc.

Then, we define a function  $u : B_T(0, c(n)) \rightarrow \mathbb{R}$  by

$$u(y) = d_g^2(0, y) + d_g^2(0, \pi_\gamma y) + \dots + d_g^2(0, \pi_\gamma^{j-1} y).$$

Since  $\pi_\gamma^j = id$ , it is easy to see  $u(\pi_\gamma y) = u(y)$  for any  $y \in B_T(0, c(n))$ . That is to say,  $u$  is  $\pi_\gamma$ -invariant. By Step 3,  $u$  is strictly  $g$ -geodesically convex on  $B_T(0, c(n))$ . More precisely, since for any  $g$ -geodesic  $\xi : [0, s_0] \rightarrow B_T(0, c(n))$ ,  $\pi_\gamma^i \xi$  are still  $g$ -geodesics in  $B_T(0, c(n))$ , and

$$\begin{aligned} \frac{d^2}{ds^2} u(\xi(s)) &= \nabla^2 d_g^2(0, \cdot)(\xi'(s), \xi'(s)) + \dots + \nabla^2 d_g^2(0, \cdot)(d\pi_\gamma^{j-1}(\xi'(s)), d\pi_\gamma^{j-1}(\xi'(s))) \\ &\geq \widetilde{g}(\xi'(s), \xi'(s)) > 0. \end{aligned}$$

Observe that

$$u|_{B_T(0, c(n))^c} \geq j(1 - c(n))^2(c(n) - \frac{2lc(n)^3}{l})^2 \geq \frac{jc(n)^2}{2},$$

and

$$u(0) \leq j(jl)^2 \leq jc(n)^5 < \frac{jc(n)^2}{2},$$

so the minimum of function  $u$  over  $\overline{B_T(0, c(n))}$  is only achieved at an interior point, say  $y_0 \in B_T(0, c(n))$ . Then by  $\pi_\gamma$  invariance of  $u$ , we have  $u(\pi_\gamma y_0) = u(y_0) < jc(n)^2/2$ , and this implies  $\pi_\gamma(y_0) \in B_T(0, c(n))$ . By the injectivity radius estimate at  $y_0 \in (T_p M, g)$ , there exists a  $g$ -geodesic connecting  $y_0$  to  $\pi_\gamma(y_0)$ ,



which is contained in  $B_{T,p}(0, 2c(n))$ . By using the strong  $g$ -geodesic convexity of  $u$ , we conclude that  $\pi_\gamma y_0 = y_0$ . This contradicts the fact that  $\pi_\gamma$  has no fixed point, and the claim is proved.

*Step 6.* The pull back of the volume element of  $g$  is the same as the one of  $g_T$ . By combining this observation with our results in Steps 4 and 5 we find

$$\mathbf{Vol}_{g_T}(B_T(0, 1)) \geq \frac{c(n)^3}{l} \mathbf{Vol}_g(B_T(p, c(n))),$$

which gives

$$l \geq c(n) \frac{\mathbf{Vol}_g(B_T(p, c(n)))}{\mathbf{Vol}_{g_T}(B_T(0, 1))} \geq c(n) \mathbf{Vol}_g(B_T(p, c(n))).$$

The proof of Theorem 1.1 is completed.  $\square$

## 8 Volume comparison for future or past cones

In Riemannian geometry, under a Ricci curvature lower bound, Bishop-Gromov's volume comparison theorem allows one to compare the volume of small and large balls in a sharp and qualitative manner. Let us return to Step 2 of Section 5, where we introduced the index form associated with the synchronous coordinate system on time-like geodesics. By noticing that the index form is symmetric and that Jacobi fields minimize the index form (in some sense), we can extend the method of proof of the index comparison theorem. However, in a general Lorentzian manifold, since the index form we needed (without imposing a restriction on the geodesics) is non-symmetric, we need to adapt the method of the index comparison theorem, as follows.

**Theorem 8.1** (Volume comparison theorem for cones). *Let  $(M, g)$  be a globally hyperbolic, Lorentzian  $(n + 1)$ -manifold. Fix  $p \in M$  and a vector  $T \in T_p M$  with  $g_p(T, T) = -1$ , and suppose that the exponential map  $\mathbf{exp}_p$  is defined on the ball  $B_T(0, r_0) \subset T_p M$  (determined by the reference inner product  $g_T$  at  $p$ ). Suppose also that the Ricci curvature satisfies on  $B_T(p, r_0)$*

$$\mathbf{Ric}_g(V, V) \geq -n K_2 |V|_g^2 \quad \text{for all time-like vector fields } V.$$

Then for any  $0 < r < s < r_0$  the inequality

$$\frac{\mathbf{Vol}_g(\mathcal{FC}(p, r))}{\mathbf{Vol}_g(\mathcal{FC}(p, s))} \geq \frac{\mathbf{Vol}_{K_2}(B(r))}{\mathbf{Vol}_{K_2}(B(s))}$$

holds, with  $\mathcal{FC}(p, r) := \mathbf{exp}_p(\mathcal{FC}(p, r))$  and

$$\mathcal{FC}(p, r) := \left\{ 0 < |V|_{g_{T,0}} < r_0, |V|_g^2 < 0, \langle T, V \rangle_{g_{T,0}} < 0 \right\}$$

and  $\mathbf{Vol}_{K_2}(B(r))$  is the volume of the ball with radius  $r$ ,  $B(r)$  (analogous to  $\mathcal{B}_T(p, r) \subset M$ ), in the simply-connected Lorentzian  $(n + 1)$ -manifold with constant curvature  $K_2$  (that is,  $R_{\alpha\beta\gamma\delta} = -K_2(g_{\alpha\gamma}g_{\beta\delta} - g_{\alpha\delta}g_{\beta\gamma})$ ).

More generally, if  $\Sigma$  is a subset in unit sphere  $S^n$  such that  $|V|_g^2 < 0, \langle T, V \rangle_g < 0$  for all  $V \in \Sigma$ , then the inequality

$$\frac{\mathbf{Vol}_g(\mathcal{FC}_\Sigma(p, r))}{\mathbf{Vol}_g(\mathcal{FC}_\Sigma(p, s))} \geq \frac{\mathbf{Vol}_{K_2}(B(r))}{\mathbf{Vol}_{K_2}(B(s))}$$

holds with  $\mathcal{FC}_\Sigma(p, r) := \mathbf{exp}_p(\mathcal{FC}_\Sigma(p, r))$  and

$$\mathcal{FC}_\Sigma(p, r) := \left\{ V \in \mathcal{FC}(p, r) / \frac{V}{|V|_{g_T}} \in \Sigma \right\}.$$

This result will be used shortly to control the injectivity radius of null cones, but is also of independent interest. For definiteness we state the result for future cones.

*Proof.* Let  $\gamma : [0, s_0] \rightarrow M$  be a future-oriented time-like geodesic satisfying  $\gamma(0) = p$  and  $|\gamma'(0)|_{g_T} = -1$ . We are going to use the standard technique to compute the rate of change of the volume element along  $\gamma$ . Given  $s_1 \in (0, s_0)$  assume that any point in  $(0, s_1]$  is neither a conjugate point nor a cut point with respect to  $p$ . Let  $v_0 = \gamma'(s_1), v_1, v_2, \dots, v_n$  be an orthonormal basis at  $\gamma(s_1)$  with respect to  $g_{\gamma(s_1)}$ . Let also  $V_\alpha$  be the Jacobi field defined on  $[0, s_1]$  and satisfying  $V_\alpha(0) = 0$  and  $V_\alpha(s_1) = v_\alpha$ . Clearly  $V_0 = (s/s_1)\gamma'$ , and the vectors  $V_i$  and  $\nabla_{\gamma'} V_i$  are orthogonal to  $\gamma'$  for all  $i \geq 1$ .

Consider the Jacobian of the exponential map  $\varphi(s) := J(d\mathbf{exp}_{\gamma(s)})$ , which is given by

$$\varphi(s)^2 = \frac{|\gamma'(s) \wedge V_1(s) \wedge \dots \wedge V_n(s)|_g^2}{s^{2n} |\gamma'(0) \wedge V_1(0) \wedge \dots \wedge V_n(0)|_g^2}.$$

Denote by  $\varphi_{K_2}(s)$  the corresponding quantity in the simply connected Lorentzian  $(n + 1)$ -manifold with constant curvature  $-K_2$ . Define the index form

$$I_{s_1}(X, Y) := \int_0^{s_1} \left( \langle \nabla_{\gamma'} X, \nabla_{\gamma'} Y \rangle_g - \mathbf{Rm}_g(\gamma', X, \gamma', Y) \right) ds,$$

where  $X, Y$  are vector fields along  $\gamma$  and  $\mathbf{Rm}_g(\gamma', X, \gamma', Y) := -\langle \mathbf{Rm}_g(\gamma', X)\gamma', Y \rangle_g$ . Observe that  $I_{s_1}$  is symmetric in  $X, Y$ . It is easy to see

$$\begin{aligned} \frac{d}{ds} \Big|_{s=s_1} \log \varphi^2 &= \sum_i \langle V'_i(s_1), V_i(s_1) \rangle_g - \frac{2n}{s_1} \\ &= \sum_i I_{s_1}(V_i, V_i) - \frac{2n}{s_1}. \end{aligned}$$

Let  $E_i(s)$  be the parallel transport of  $v_i$  along  $\gamma$ . Since there are no conjugate points along  $\gamma$ , the Jacobi field minimizes the index form among all vector fields

with fixed boundary values. This is the same as in Riemannian geometry. The reason is that the length of time-like geodesic without conjugate points is locally maximizing among all nearby time-like curves with the same end points. Let  $\tilde{V}_i(s) = \frac{\sinh s}{\sinh s_1} E_i(s)$ , then  $I_{s_1}(V_i, V_i) \leq I_{s_1}(\tilde{V}_i, \tilde{V}_i)$  and

$$\begin{aligned} \frac{d}{ds} \Big|_{s=s_1} \log \frac{\varphi^2}{(\varphi_{K_2})^2} &\leq - \sum_i \int_0^{s_1} \frac{(\sinh s)^2}{(\sinh s_1)^2} (\mathbf{Rm}_g(\gamma', E_i, \gamma' E_i) - K_2) \\ &= - \int_0^{s_1} \frac{(\sinh s)^2}{(\sinh s_1)^2} (\mathbf{Ric}_g(\gamma', \gamma') - n K_2) ds \leq 0. \end{aligned}$$

The following is a simple but very important observation due to Gromov, which we now extend to a globally hyperbolic Lorentzian manifold. Let  $A$  be the star-shaped domain (with respect to 0) in  $T_p M$ , such that  $\mathbf{exp}_p : A \cap B_T(0, r_0)$  is a diffeomorphism on its image and the image of  $\partial A \cap B_T(0, r_0)$  is set of cut locus (in  $\mathcal{B}_T(p, r_0)$ ). Let  $\chi_A$  be the characteristic function of  $A$ . Since  $\varphi(s)/\varphi_{K_2}(s)$  is decreasing in  $s$  we see that  $\chi_A \varphi/\varphi_{K_2}$  is also decreasing in  $s$ . Now, we get two functions on  $B_T(0, r_0)$ , whose quotient is decreasing along radial geodesics. Observe that  $M$  is globally hyperbolic, so any point in  $\mathcal{FC}(p, r_0)$  is connected to  $p$  by a maximizing time-like geodesic. This also implies that the integration of  $\chi_A \varphi$  over  $B_T(0, s)$  gives the volume  $\mathbf{Vol}_g(\mathcal{B}_T(p, s))$ . Then, by integrating  $\chi_A \varphi$  and  $\varphi_{K_2}$  over  $B_T(0, s)$  and after a simple calculation we deduce that  $\mathbf{Vol}_g(\mathcal{FC}(p, s))/\mathbf{Vol}_{K_2}(B(s))$  is decreasing in  $s$ . The case of the ratio  $\mathbf{Vol}_g(\mathcal{FC}_\Sigma(p, s))/\mathbf{Vol}_{K_2}(B(s))$  is similar. The proof of the theorem is completed.  $\square$

We are now in a position to prove :

**Corollary 8.2** (Injectivity radius based on the volume of a future cone). *Let  $M$  be a manifold satisfying the assumptions in Theorem 1.1 and assumed to be globally hyperbolic, and let  $T \in T_p M$  be a reference vector. Let  $\Sigma$  be a subset in the unit sphere  $S^n$  included in the future cone  $N_p^+$ . If  $\mathbf{Vol}_g(\mathcal{FC}_\Sigma(p, r_0)) \geq v_0 > 0$ , then the inequality*

$$\frac{\mathbf{Inj}_g(M, p, T)}{r_0} \geq c(\Sigma) \frac{v_0}{r_0^{n+1}}$$

holds, where  $\mathcal{FC}_\Sigma(p, r_0) := \mathbf{exp}_p(\mathcal{FC}_p(r_0))$  with

$$\mathcal{FC}_p(r_0) := \left\{ 0 < |V|_T < r_0, \langle T, V \rangle_T < 0, |V|_g^2 < 0, \frac{V}{|V|_T} \in \Sigma \right\},$$

and the constant  $c(\Sigma)$  depends only on the distance (measured by  $T$ ) of  $\Sigma$  to the null cone.

*Proof.* First, we recall there is a constant  $C(\Sigma)$  depending only on the distance of  $\Sigma$  to the null cone, such that  $\mathbf{Ric}(\gamma', \gamma') \geq -C(\Sigma)|\gamma'|_g^2$  for any time-like geodesic  $\gamma$  with  $\gamma'(0) \in \Sigma$ . By the volume comparison theorem for future cone established in Theorem 8.1 we have

$$\frac{\mathbf{Vol}_g(\mathcal{FC}_\Sigma(p, c(n)r_0))}{\mathbf{Vol}_g(\mathcal{FC}_\Sigma(p, r_0))} \geq C(\Sigma),$$

and combining this result with Theorem 1.1, the corollary follows.  $\square$

## 9 Final remarks

### Regularity of Lorentzian metrics

Following the strategy proposed in the present paper, we now “transfer” to the Lorentzian metric the regularity available on a reference Riemannian metric. Clearly, the regularity obtained in this manner depends on the way the reference Riemannian metric is constructed. The interest of our approach below is to provide a simple derivation: using harmonic-like coordinates for the Riemannian metric we see immediately that the Lorentzian metric has uniformly bounded first-order derivatives. For the optimal regularity achievable with Lorentzian metrics we refer to Anderson [3].

**Proposition 9.1** (Regularity in harmonic-like coordinates). *Under the assumptions and notation of Theorem 1.1, define*

$$r_1 := c(n) \frac{\mathbf{Vol}_g(\mathcal{B}_T(p, c(n) r_0))}{r_0^{n+1}} r_0,$$

where  $c(n)$  is the constant determined in this theorem. Then for any  $\varepsilon > 0$  there exist a constant  $c_1(n, \varepsilon)$  with  $\lim_{\varepsilon \rightarrow 0} c_1(n, \varepsilon) = 0$  and a coordinate system  $(x^\alpha)$  satisfying  $x^\alpha(p) = 0$  and defined for all  $(x^0)^2 + (x^1)^2 + \dots + (x^n)^2 < (1 - \varepsilon)^2 r_1^2$ , such that

$$\begin{aligned} |g_{\alpha\beta} - \eta_{\alpha\beta}| &\leq c_1(n, \varepsilon), \\ r_1 |\partial g_{\alpha\beta}| &\leq c_1(n, \varepsilon), \end{aligned} \tag{9.1}$$

where  $\eta_{\alpha\beta}$  is the Minkowski metric in these coordinates.

*Proof.* By scaling we may assume  $r_1 = 1$ . By Step 1 in the proof of Theorem 1.1, we know that the Riemannian metric  $g_T$  is equivalent to the Riemannian metric  $g_{T,0}$  on  $\mathcal{B}_T(0, 4c(n))$ . By considering a lift and using again the results in Step 1 this implies

$$\mathcal{B}_T(p, c(n)) \subset \mathcal{B}_T(q, 3c(n)) \quad q \in \mathcal{B}_T(p, c(n)).$$

Applying the same argument as in Theorem 1.1, we deduce that the injectivity radius of any point in  $\mathcal{B}_T(p, c(n))$  is bounded from below by  $c(n)$ . As in Step 3 in the proof of Theorem 1.1 (or in Step 2 of Section 5), we see that there exists a synchronous coordinate system  $(y^\alpha) = (\tau, y^j)$  of definite size around  $p$  such that the metrics  $g = -d\tau^2 + g_{ij} dy^i dy^j$  and  $g_N = d\tau^2 + g_{ij} dy^i dy^j$  (the Riemannian metric constructed therein) satisfy the following properties on the geodesic ball  $\mathcal{B}_T(p, c(n))$ :

- (a)  $(1 - c(n)) g_N \leq g_T \leq (1 + c(n)) g_N$ ,
- (b)  $g_N$  has bounded curvature ( $\leq 1/c(n)$ ),

$$(c) \quad |\tau| + \frac{1}{|\tau|} + |\nabla^2 \tau|_N \leq 1/c(n).$$

(In particular, this implies  $|\nabla_{g_N} g|_N < 1/c(n)$ .) Since the volume  $\mathbf{Vol}_{g_N}(\mathcal{B}_T(p, c(n)))$  is bounded from below, it follows from [10] that the injectivity radius of  $g_N$  at  $p$  is bounded from below by  $c(n)$ . By the theorem in [16] on the existence of harmonic coordinates, for any small  $\varepsilon > 0$  there exists an harmonic coordinate system  $(x^\alpha)$  with respect to the Riemannian metric  $g_N$  such that  $\sum_\alpha |x^\alpha|^2 < (1-\varepsilon)^2$  and for every  $0 < \gamma < 1$

$$|g_{N,\alpha\beta} - \delta_{\alpha\beta}| < c_1(n, \varepsilon), \quad |\partial g_N| < 1/c(n), \quad |\partial g_N|_{C^\gamma} < 1/c(n, \varepsilon, \gamma).$$

In the construction of harmonic coordinates, we may also assume that  $|\frac{\partial}{\partial y_0} - \frac{\partial}{\partial \tau}|_{g_{T_0}} < c_1(n, \varepsilon)$ .

Since  $|\nabla_{g_N} g|_N < 1/c(n)$  and that, in these coordinates,  $|\nabla_{g_N}| \leq 1/c(n)$ , we have  $|\partial g| < 1/c(n)$ . Finally, to estimate the metric we write  $|g_{\alpha\beta} - \eta_{\alpha\beta}|_p < c_1(n, \varepsilon)$  and  $|\partial g| < 1/c(n)$  and we conclude that  $|g_{\alpha\beta} - \eta_{\alpha\beta}| < \frac{1}{c(n)}\varepsilon + c_1(n, \varepsilon)$ . The proof is completed.  $\square$

## Pseudo-Riemannian manifolds

Finally, we would like discuss pseudo-Riemannian manifolds  $(M, g)$  (also referred to as semi-Riemannian manifolds). Consider a differentiable manifold  $M$  endowed with a symmetric, non-degenerate covariant 2-tensor  $g$ . We assume that the signature of  $g$  is  $(n_1, n_2)$ , that is,  $n_1$  negative signs and  $n_2$  positive signs. Riemannian and Lorentzian manifolds are special cases of pseudo-Riemannian manifolds. Fix  $p \in M$  and an orthonormal family  $T$  consisting of  $n_1$  vectors  $E_1, E_2, \dots, E_{n_1} \in T_p M$  such that  $\langle E_i, E_j \rangle_g = -\delta_{ij}$ . Based on this family, we can define a reference inner product  $g_T$  on  $T_p M$  by generalizing our construction in the Lorentzian case, and by using this inner product we can then define the ball  $B_T(0, s) \subset T_p M$ . By parallel translating  $E_1, E_2, \dots, E_{n_1}$  along radial geodesics from the origin in  $T_p M$ , we obtain vector fields  $E_1, E_2, \dots, E_{n_1}$  defined in the tangent space (or multi-valued vector fields on the manifold). This also induces a (multi-valued) Riemannian metric  $g_T$  as was explained before.

The following corollary immediately follows by repeating the proof of Theorem 1.1. We note that the curvature covariant derivative bound imposed below is probably superfluous and could probably be removed by introducing a foliation based on certain synchronous-type coordinates, as we did in Section 5 for Lorentzian manifolds. On the other hand, to the best of our knowledge this is the first injectivity radius estimate for pseudo-Riemannian manifolds.

**Corollary 9.2** (Injectivity radius of pseudo-Riemannian manifolds). *Let  $(M, g)$  be a differentiable pseudo-Riemannian  $n$ -manifold with signature  $(n_1, n_2)$ , and let  $p \in M$  and  $T = (E_1, \dots, E_{n_1})$  be a family of vectors in  $T_p M$  satisfying  $g(E_i, E_j) = -\delta_{ij}$ . Suppose that the exponential map  $\mathbf{exp}_p$  is defined on  $B_T(0, r_0) \subset T_p M$  and that*

$$|\mathbf{Rm}_g|_T \leq r_0^{-2}, \quad |\nabla \mathbf{Rm}_g|_T \leq r_0^{-3} \quad \text{on } B_T(0, r_0).$$

Then, there exists a positive constant  $c(n)$  such that

$$\frac{\mathbf{Inj}_g(M, p, T)}{r_0} \geq c(n) \frac{\mathbf{Vol}_g(\mathcal{B}_T(p, c(n) r_0))}{r_0^n},$$

where  $\mathcal{B}_T(p, r) = \mathbf{exp}_p(\mathcal{B}_T(0, r))$  is the geodesic ball at  $p$  with radius  $r$ .

*Proof.* Without loss of generality we assume  $r_0 = 1$ . In local coordinate system  $y^\alpha$ , let

$$E_i =: E_i^\beta \frac{\partial}{\partial y^\beta}, \quad E_{i\alpha} = E_i^\beta g_{\alpha\beta}, \quad i = 1, \dots, n_1,$$

then  $g_{T, \alpha\beta} = g_{\alpha\beta} + 2 \sum_{i=1}^{n_1} E_{i\alpha} E_{i\beta}$ . By the same computations as in the proof of Theorem 1.1 we obtain

$$|\nabla E_i|_T \leq \frac{1}{c(n)},$$

$$|g_T - g_{T,0}| + |g - \eta| < c(n) \quad \text{on the ball } \mathcal{B}_T(0, c(n)),$$

where  $\eta_{\alpha\beta} := \mp \delta_{\alpha\beta}$  (a minus sign for  $\alpha \leq n_1$ , and a plus sign for  $\alpha > n_1$ ). In view of the computations in [13] (Theorem 4.11 and Corollary 4.12) we deduce that  $|\partial g| < r/c(n)$ , where  $r^2 = (y^1)^2 + \dots + (y^n)^2$ . Since  $d_{g_{T,0}}^2(y_0, y) = |y - y_0|^2$ , we have for any point  $y_0 \in \mathcal{B}_T(0, c(n))$

$$\nabla_{\alpha\beta}^2 d_{g_{T,0}}^2(y_0, \cdot) \geq \delta_{\alpha\beta} = g_{T,0} \quad \text{on the ball } \mathcal{B}_T(0, c(n)).$$

Since the metric  $g_{T,0}$  plays the same role as  $g_N$  (cf. the proof of Theorem 1.1), all arguments can be carried out and this completes the proof of the corollary.  $\square$

## Acknowledgements

The first author (BLC) was partially supported by Sun Yat-Sen University via China-France-Russia collaboration grant (No. 34000-3275100), the Ecole Normale Supérieure de Paris, the French Foreign Ministry, and the Institut des Hautes Études Scientifiques (IHES, Bures-sur-Yvette). The second author (PLF) was partially supported by the A.N.R. (Agence Nationale de la Recherche) through the grant 06-2-134423 entitled “*Mathematical Methods in General Relativity*” (MATH-GR), and by the Centre National de la Recherche Scientifique (CNRS).

## References

- [1] M.T. ANDERSON, Convergence and rigidity of manifolds under Ricci curvature bounds, *Invent. Math.* 102 (1990), 429–445.
- [2] M.T. ANDERSON, On long-time evolution in general relativity and geometrization of 3-manifolds, *Commun. Math. Phys.* 222 (2001), 533–567.

- [3] M.T. ANDERSON, Regularity for Lorentz metrics under curvature bounds, *Jour. Math. Phys.* 44 (2003), 2994–3012.
- [4] A. BESSE, *Einstein manifolds*, Ergebnisse de Math. Series 3, Springer Verlag, 1987.
- [5] J. CHEEGER, Finiteness theorems for Riemannian manifolds, *Amer. J. Math.* 92 (1970), 61–94.
- [6] J. CHEEGER AND D. EBIN, *Comparison theorems in Riemannian geometry*, North-Holland, Amsterdam-Oxford, American Elsevier Pub., New York, 1975.
- [7] J. CHEEGER AND M. GROMOV, Collapsing Riemannian manifolds while keeping their curvature bounded. I, *J. Diff. Geom.* 23 (1986), 309–346.
- [8] J. CHEEGER AND M. GROMOV, Collapsing Riemannian manifolds while keeping their curvature bounded. II, *J. Diff. Geom.* 32 (1990), 269–298.
- [9] J. CHEEGER, K. FUKAYA, AND M. GROMOV, Nilpotent structures and invariant metrics on collapsed manifolds, *J. Amer. Math. Soc.* 5 (1992), 327–372.
- [10] J. CHEEGER, M. GROMOV, AND M. TAYLOR, Finite propagation speed, kernel estimates for functions of the Laplace operator, and the geometry of complete Riemannian manifolds, *J. Differential Geom.* 17 (1982) 15–53.
- [11] S.Y. CHENG, P. LI, AND S.T. YAU, Heat equations on minimal submanifolds and their applications, *Amer. J. Math.* 106 (1984), 1033–1065.
- [12] D.M. DETURCK AND J.L. KAZDAN, Some regularity theorems in Riemannian geometry, *Ann. Sci. École Norm. Sup.* 14 (1981), 249–260.
- [13] R.S. HAMILTON, A compactness property for solution of the Ricci flow, *Amer. J. Math.* 117 (1995), 545–572.
- [14] S. HAWKING AND G.F. ELLIS, *The large scale structure of space-time*, Cambridge Univ. Press, 1973.
- [15] E. HEINTZE AND H. KARCHER, A general comparison theorem with applications to volume estimates for submanifolds, *Ann. Sci. Ecole Norm. Sup.* 11 (1978), 451–470.
- [16] J. JOST AND H. KARCHER, Geometrische Methoden zur Gewinnung von a-priori-Schranken für harmonische Abbildungen, *Manuscripta Math.* 40 (1982), 27–77.
- [17] S. KLAINERMAN AND I. RODNIANSKI, Ricci defects of microlocalized Einstein metrics, *J. Hyperbolic Differ. Equa.* 1 (2004), 85–113.
- [18] S. KLAINERMAN AND I. RODNIANSKI, Rough solutions of the Einstein-vacuum equations, *Ann. of Math.* 161 (2005), 1143–1193.

- [19] S. KLAINERMAN AND I. RODNIANSKI, On the radius of injectivity of null hypersurfaces, J. Amer. Math. Soc., to appear.
- [20] R. PENROSE, *Techniques of differential topology in relativity*, CBMS-NSF Region. Conf. Series Appl. Math., Vol. 7, 1972.
- [21] S. PETERS, Convergence of Riemannian manifolds, *Compositio Math.* 62 (1987), 3–16.
- [22] P. PETERSEN, Convergence theorems in Riemannian geometry, in “Comparison Geometry” (Berkeley, CA, 1992–93), MSRI Publ. 30, Cambridge Univ. Press, 1997, pp. 167–202.
- [23] J.H.C. WHITEHEAD, Convex regions in the geometry of paths, *Quart. J. Math. Oxford* 3 (1932), 33–42.