

From Stationary Phase to Steepest Descent

Spyridon Kamvissis

Dedicated to Percy Deift on his 60th birthday

0. INTRODUCTION

The so-called nonlinear stationary-phase-steepest-descent method for the asymptotic analysis of Riemann-Hilbert factorization problems has been very successful in providing

(i) rigorous results on long time, long range and semiclassical asymptotics for solutions of completely integrable equations and correlation functions of exactly solvable models,

(ii) asymptotics for orthogonal polynomials of large degree,

(iii) the eigenvalue distribution of random matrices of large dimension (and related universality results),

(iv) proofs of important results in combinatorial probability (e.g. the limiting distribution of the length of longest increasing subsequence of a permutation, under uniform distribution).

Even though the stationary phase idea was first applied to a Riemann-Hilbert problem and a nonlinear integrable equation by Its ([I], 1982) the method became systematic and rigorous in the work of Deift and Zhou ([DZ], 1993). As a recognition

2000 *Mathematics Subject Classification*. Primary 37K40.

of the fruitfulness of the method, Percy Deift was invited to give a plenary address to the recent ICM in Madrid, on the subject of universality in mathematics and physics. Of course, the main mathematical tool in proving universality theorems has been the nonlinear stationary-phase-steepest-descent method.

In analogy to the linear stationary-phase and steepest-descent methods, where one asymptotically reduces the given exponential integral to an exactly solvable one, in the nonlinear case one asymptotically reduces the given Riemann-Hilbert problem to an exactly solvable one.

Our aim here is to clarify the distinction between the stationary-phase idea and the steepest-descent idea, stressing the importance of actual steepest-descent contours in some problems. We claim that the distinction partly mirrors the self-adjoint / non-self-adjoint dichotomy of the underlying Lax operator. To this aim we first have to review some of the main groundbreaking ideas (due to Percy Deift and his collaborators) appearing in the self-adjoint case; then we describe recent results ([KMM], [KR]) in the non-self-adjoint case, that we see as a natural extension. We mostly use the defocusing / focusing nonlinear Schrödinger equation as our working model, but we also digress to the KdV at some point.

We stress both here and in the main text that an extra feature appearing only in the nonlinear asymptotic theory is the Lax-Levermore variational problem, discovered in 1979, before the work of Its, Deift and Zhou, but closely related to the so-called "g-function" which is catalytic in the process of deforming Riemann-Hilbert factorization problems to exactly solvable ones.

1. THE LINEAR METHOD

Suppose one considers the Cauchy problem for the linearized KdV equation: $u_t + u_{xxx} = 0$. It can of course be solved via Fourier transforms. The end result of the Fourier method is an exponential integral. To understand the long time asymptotic behavior of the integral one needs to apply the stationary-phase method (see e.g. [E]). The underlying principle, going back to Stokes and Kelvin, is that the dominating contribution comes from the vicinity of the *stationary phase points*.

Through a local change of variables at each stationary phase point and using integration by parts we can calculate each contributing integral asymptotically to all orders with exponential error. It is essential here that the phase¹ $x\xi - \xi^3 t$ is real and that the stationary phase points are real.

On the other hand, suppose we have something like the Airy exponential integral

$$(1) \quad \text{Ai}(z) = \frac{1}{\pi} \int_0^\infty \cos\left(\frac{s^3}{3} + zs\right) ds,$$

and we are interested in $z \rightarrow \infty$. Set $s = z^{1/2}t$ and $x = z^{3/2}$.

$$(2) \quad \text{Ai}(x^{2/3}) = \frac{x^{1/3}}{2\pi} \int_{-\infty}^\infty \exp\left(ix\left(\frac{t^3}{3} + t\right)\right) dt.$$

The phase is $h(t) = \frac{t^3}{3} + t$ and the zeros of $h'(t) = (t^2 + 1)$ are $\pm i$. As they are not real, and since the integrand is analytic, one must deform the integral off the real line, along particular paths. These are referred to as *steepest descent paths*. They are given by the simple characterization

$$(3) \quad \text{Im}h(t) = \text{constant}.$$

In our particular example, the curves of steepest descent are the imaginary axis and the two branches of a hyperbola. By deforming to one of these branches, we finally end up with integrals which can be analyzed directly, using the so-called Laplace's method (which is simpler than the stationary phase method). We thus recover asymptotics valid to all orders.

The nonlinear method generalizes the ideas above, but also employs new ones.

2. THE NONLINEAR METHOD

(i) The Stationary Phase Idea

Consider the defocusing nonlinear Schrödinger equation

$$(4) \quad \begin{aligned} i\partial_t \psi + \partial_x^2 \psi - |\psi|^2 \psi &= 0, \\ \text{under } \psi(x, 0) &= \psi_0(x), \end{aligned}$$

¹ ξ is the spectral variable

where the initial data function lies in, say, Schwartz space. The analog of the Fourier transform is the scattering coefficient $r(\xi)$ for the Dirac operator

$$L = \begin{pmatrix} i\partial_x & i\psi_0(x) \\ -i\psi_0^*(x) & -i\partial_x \end{pmatrix}.$$

Suppose we are now interested in the long time behavior of the solution to (4). The inverse scattering problem can be posed in terms of a Riemann-Hilbert factorization problem.

THEOREM. There exists a 2×2 matrix Q with analytic entries in the upper and lower open half-planes, such that the normal limits Q_+, Q_- , as ξ approaches the real line from above or below respectively, exist and satisfy

$$(5) \quad Q_+(\xi) = Q_-(\xi) \begin{pmatrix} 1 - |r(\xi)|^2 & -r^*(\xi)e^{-2i\xi x - 4i\xi^2 t} \\ r(\xi)e^{2i\xi x + 4i\xi^2 t} & 1 \end{pmatrix}, \quad \text{Im}\xi = 0,$$

and $\lim_{\xi \rightarrow \infty} Q(\xi) = I.$

The solution to (4) is recovered via

$$(6) \quad \psi(x, t) = -2 \lim_{\xi \rightarrow \infty} \xi Q_{12}(\xi).$$

It was first realized by Its [I, IN], that the leading order behavior of the long time asymptotics for the solution of (4) can be described by replacing the problem (5) by a "local" model Riemann-Hilbert problem located in a small neighborhood of the *stationary phase point* $\xi_0 = -\frac{x}{4t}$ satisfying $\Theta'(\xi_0) = 0$ where $\Theta = \xi x + 2\xi^2 t$, or, equivalently to a problem similar to (5) but where ξ is replaced by the constant value ξ_0 (see (7) below). But no idea was given on how to show that this solution of the full problem and of the model problem are actually close to each other. To show how to do this, was the work of Deift and Zhou.² The basic ideas of [DZ] have been used in all works on the stationary-phase-steepest-descent-method since. They include:

1. Appropriate lower/diagonal/upper factorizations of jump matrices.
2. Equivalence of the solvability of inhomogeneous matrix Riemann-Hilbert

problems to the invertibility of associated singular integral operators. This idea

²See [DZ] for a comprehensive review of the history of the problem before the use of Riemann-Hilbert problem techniques.

goes at least back to Gohberg [CG]. A crucial contribution of Beals and Coifman [BC] was to make this precise for contours with self-intersections (which they needed in their study of the inverse scattering of first order systems). These ideas were further developed by Zhou [Z] who provided a very useful existence theorem for matrix Riemann-Hilbert problems with jumps and jump contours satisfying some special Schwarz reflection type symmetries and an integral formula expressing the solution of the Riemann-Hilbert problem in terms of the inverse of a particular weighted Cauchy operator depending on a given factorization of the jump matrix and thus taking advantage of the factorization mentioned above. Perturbing Zhou's formula provides a nice way to show that under some conditions, small changes in the jump data result in small changes in the solution.

3. Introduction and solution of auxiliary scalar problems.

Following analyticity and the above ideas one ends up with a problem on a *small cross centered at the stationary phase point*. Using a rescaling the Riemann-Hilbert problem is rescaled to a new problem on an infinite cross. After deforming the components of the cross back to the real line, it is equivalent to the following problem on the real line:

$$(7) \quad H_+(\xi) = H_-(\xi) \exp(-i\xi^2 \sigma_3) \begin{pmatrix} 1 - |r(\xi_0)|^2 & -r^*(\xi_0) \\ r(\xi_0) & 1 \end{pmatrix} \exp(i\xi^2 \sigma_3),$$

$$H(\xi) \sim \xi^{i\nu \sigma_3},$$

where ν is a constant depending only on ξ_0 and $\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ is a Pauli matrix.

So the jump matrix $\begin{pmatrix} 1 - |r(\xi)|^2 & -r^*(\xi) \\ r(\xi) & 1 \end{pmatrix}$ of the original problem is replaced by its value at ξ_0 .

Problem (7) can be solved explicitly. Written in terms of the new unknown $H(\xi) \exp(-i\xi^2 \sigma_3)$ it has a constant jump and can thus be reduced to a first order linear matrix ODE ([I]).

(ii). The finite-gap g-function mechanism and a "shock" phenomenon with no linear analogue.

An important step halfway between the leap from the "stationary phase" idea to the general definition of a "steepest descent contour" is the introduction of the so-called finite-gap g -function mechanism. The g -function was introduced in [DZ95] in the special case of genus 0 and in [DVZ94] in the special case of genus 1 but the full force of the finite-gap g -function idea and the connection to the Lax-Levermore variational problem was first explored in the analysis of the KdV equation [DVZ97]

$$u_t - 6uu_x + \epsilon^2 u_{xxx} = 0,$$

$$u(x, 0) = u_0(x),$$

in the limit as $\epsilon \rightarrow 0$. Assume for simplicity, that the initial data are real analytic, positive and consist of a "hump" of unit height.

The associated RH problem is

$$S_+(z) = S_-(z) \begin{pmatrix} 1 - |r(z)|^2 & -r^*(z)e^{-\frac{izx-4iz^3t}{\epsilon}} \\ r(z)e^{\frac{izx+4iz^3t}{\epsilon}} & 1 \end{pmatrix}, \quad \text{Im}z = 0,$$

and $\lim_{z \rightarrow \infty} S(z) = (1, 1)$,

where r is the reflection coefficient for the Schrödinger operator with potential u_0 .

The solution of KdV is recovered via

$$u(x, t; \epsilon) = -2i\epsilon \frac{\partial}{\partial x} S_1^1(x, t; \epsilon),$$

where S_1^1 is the residue of the first entry of S at infinity. The reflection coefficient r also depends on ϵ . In fact, the WKB approximation is

$$r(z) \sim -ie^{-\frac{2i\rho(z)}{\epsilon}} \chi_{[0,1]}(z)$$

$$1 - |r(z)|^2 \sim e^{-\frac{2\tau(z)}{\epsilon}},$$

where

$$\rho(z) = x_+ z + \int_{x_+}^{\infty} [z - (z^2 - u_0(x))^{1/2}] dx,$$

$$\tau(z) = \text{Re} \int (u_0(x) - z^2)^{1/2} dx$$

and $x_+(z)$ is the largest solution of $u_0(x_+) = z^2$.

[DVZ97] introduce the following change of variables $\hat{S}(z) = S(z)e^{\frac{ig(z)\sigma_3}{\epsilon}}$ where g is a scalar function defined by the following conditions.

1. g is analytic off the interval $[0, 1]$, the normal limits g_+, g_- of g exist along $[0, 1]$ and g vanishes at infinity.

2. "Finite gap ansatz". Define $h(z) = g_+(z) + g_-(z) - 2\rho + 4tz^3 + xz$. There exists a finite set of disjoint open real intervals ("bands") $I_j \in [0, 1]$ such that

3a. For $z \in \cup_j I_j$, we have $-\tau < (g_+ - g_-)/2i < 0$ and $h' = 0$.

3b. For $z \in [0, 1] \setminus \cup_j I_j$, we have $2i\tau = g_+ - g_-$ and $h' < 0$.

The conditions above are meant to determine not only g but also the band-gap structure in $[0, 1]$. In general (for any data u_0) it is not true that the above conditions can be satisfied. It is believed however that under the condition of analyticity a g-function satisfying the "finite gap ansatz" exists. (In fact [K00] gives a proof of the "finite gap ansatz" in the analogous problem of the continuum Toda equations.) Assuming that there is a g-function satisfying the three conditions above one can show that the RH problem reduces to one supported on the bands I_j with jumps of the form

$$\begin{pmatrix} 0 & -ie^{-ih(z)/\epsilon} \\ -ie^{ih(z)/\epsilon} & 0 \end{pmatrix},$$

and in fact, because of (iib), $h(z)$ is a real constant on each band I_j . This RH problem can be solved explicitly via theta functions. The details in [DVZ97] involve the so-called "lens"-argument: auxiliary contours are introduced near pieces of the real line (one below and one above each band/gap) and appropriate factorizations and analytic extensions are used, very similarly to the subsection above. The conditions for g above are chosen precisely to make the lens argument work.

As is remarked in [DVZ94] the fact that the new RH problem is on slits "is a new and essentially nonlinear feature of our nonlinear stationary phase method". Unlike [DVZ94] where it is defined explicitly via an integral formula, in [DVZ97] the g-function is only defined implicitly via the conditions above, which may or may not admit a solution.

(iii) The Lax-Levermore Variational Problem

The g-function satisfying conditions (i), (ii), (ia), (ib) can be written as

$$g(z) = \int \log(z - \eta) d\mu(\eta)$$

where μ is a continuous measure supported in $\cup_j I_j$. In a sense, the reduction of the given RH problem to an explicitly solvable one depends on the existence of a particular measure. Conditions (i), (ii), (ia), (ib) turn out to be equivalent to a maximization problem for logarithmic potentials under a particular external field depending on $x, t, u_0(x)$ over positive measures with an upper constraint. This is related to the famous Lax-Levermore Variational Problem [LL]. Even though it appears as an afterthought in [DVZ97] (though clearly serving as inspiration), it seems that its analysis is essential for the justification of the method (as in [K00]).

3. STEEPEST DESCENT CONTOURS

Having reviewed some essential ideas in the previous sections, we are ready to consider the focusing NLS equation, following [KMM].

$$i\hbar\partial_t\psi + \frac{\hbar^2}{2}\partial_x^2\psi + |\psi|^2\psi = 0,$$

under data $\psi(x, 0) = \psi_0(x)$.

Note that the Lax operator

$$L = \begin{pmatrix} i\hbar\partial_x & -i\psi_0(x) \\ -i\psi_0^*(x) & -i\hbar\partial_x \end{pmatrix},$$

is non-self-adjoint. We shall see that the deformation of the semiclassical RH problem can be no more confined to a small neighborhood of the real axis but is instead fully two-dimensional. A *steepest descent contour* needs to be discovered!³

For simplicity consider the very specific data $\psi_0(x) = A \operatorname{sech} x$ where $A > 0$. Let $x_-(\eta) < x_+(\eta)$ be the two solutions of $\operatorname{sech}^2(x) + \eta^2 = 0$. Also assume that $\hbar = A/N$ and consider the limit $N \rightarrow \infty$. It is known that the reflection coefficient is identically zero and that the eigenvalues of L lie uniformly placed on the imaginary segment $[-iA, iA]$. In fact the eigenvalues are the points $\lambda_j = i\hbar(j + 1/2), j = 0, \dots, N - 1$ and their conjugates. The norming constants oscillate between -1 and 1 .

³By the way, in the long time asymptotics for the above with $\hbar = 1$ a collisionless shock phenomenon is also present; for x, t in the shock region the deformed RH problem is supported on a vertical imaginary slit. (See [K96].) But here, we rather focus on the semiclassical problem $\hbar \rightarrow 0$ which is far more complicated.

The associated RH problem is a meromorphic problem with no jump: to find a rational function with prescribed residues at the poles λ_j and their conjugates. It can be turned into a holomorphic problem by constructing two loops, one denoted by C say, encircling the λ_j and one, C^* , encircling their conjugates. We redefine the unknown 2×2 matrix inside the loops so that the poles vanish (there is actually a discrete infinity of choices, corresponding to an infinity of analytic interpolants of the norming constants, see below) and thus arrive at a nontrivial jump across the two loops, encircling the segments $[0, iA]$ and $[-iA, 0]$ respectively. This is a trivial deformation, valid for any h (not necessarily small). The discrete nature of the spectrum of L is mirrored in the discrete nature of the jump matrices: they involve a logarithmic integral with respect to a discrete measure. We sometimes refer to this as a discrete Riemann-Hilbert problem.

THEOREM. Let $d\mu = (\rho^0(\eta) + (\rho^0)^*(\eta^*))d\eta$, where $\rho^0 = i$ is the asymptotic density of eigenvalues supported on the linear segment $[0, iA]$. Set $X(\lambda) = \pi(\lambda - iA)$.

Letting M_+ and M_- denote the limits of M on $\Sigma = C \cup C^*$ from left and right respectively, we define the Riemann-Hilbert factorization problem

$$M_+(\lambda) = M_-(\lambda)J(\lambda),$$

where

$$\begin{aligned} J(\lambda) &= v(\lambda), \lambda \in C, \\ &= \sigma_2 v(\lambda^*)^* \sigma_2, \lambda \in C^*, \\ \lim_{\lambda \rightarrow \infty} M(\lambda) &= I, \end{aligned}$$

and

$$(8) \quad v(\lambda) = \begin{pmatrix} 1 & -i \exp\left(\frac{1}{h} \int \log(\lambda - \eta) d\mu(\eta)\right) \exp\left(-\frac{1}{h}(2i\lambda x + 2i\lambda^2 t - X(\lambda))\right) \\ 0 & 1 \end{pmatrix}.$$

Then the solution of the initial value problem for the focusing NLS equation is given by $\psi(x, t) = 2i \lim_{\lambda \rightarrow \infty} (\lambda M_{12})$.

Note that in the statement of the theorem the measure in the logarithmic integral is now continuous. We have effectively substituted a discrete set of eigenvalues by its continuous limiting density. This is only valid as $h \rightarrow 0$ and the rigorous

justification of the discrete-to-continuum passage is far from trivial, especially near the point 0 where the loops C, C^* hit the eigenvalue spike.

The analysis in [KMM] makes use of all the ideas described in the previous sections (factorization, lenses, the weighted Cauchy operator, an auxiliary scalar problem), but it also takes care of the fact that while the loops can be deformed anywhere away from the poles as long as h is not small, they have to be eventually located at a very specific position in order to asymptotically simplify the RH problem, as $h \rightarrow 0$. Appropriately, the definition of a g-function has to be generalized. Not only will it introduce the division of the loop into arcs, called "bands" and "gaps", but it must implicitly select a contour. Rather than giving the complicated set of equations and inequalities defining the g-function, we will rather focus on the associated variational problem; it is not a maximization problem but rather a maximin problem. Here's the setting.

Let $\mathbb{H} = \{z : \text{Im}z > 0\}$ be the complex upper-half plane and $\bar{\mathbb{H}} = \{z : \text{Im}z \geq 0\} \cup \{\infty\}$ be the closure of \mathbb{H} . Let also $\mathbb{K} = \{z : \text{Im}z > 0\} \setminus \{z : \text{Re}z = 0, 0 < \text{Im}z \leq A\}$. In the closure of this space, $\bar{\mathbb{K}}$, we consider the points ix_+ and ix_- , where $0 \leq x < A$ as distinct. In other words, we cut a slit in the upper half-plane along the segment $(0, iA)$ and distinguish between the two sides of the slit. The point infinity belongs to $\bar{\mathbb{K}}$, but not \mathbb{K} . Define $G(z; \eta)$ to be the Green's function for the upper half-plane

$$G(z; \eta) = \log \frac{|z - \eta^*|}{|z - \eta|}$$

and let $d\mu^0(\eta)$ be the nonnegative measure $-id\eta$ on the segment $[0, iA]$ oriented from 0 to iA . The star denotes complex conjugation. Let the "external field" ϕ be defined by

$$\phi(z) = - \int G(z; \eta) d\mu^0(\eta) - \text{Re}(\pi(iA - z) + 2i(zx + z^2t)),$$

where, without loss of generality $x > 0$.

Let \mathbb{M} be the set of all positive Borel measures on $\bar{\mathbb{K}}$, such that both the free energy

$$E(\mu) = \int \int G(x, y) d\mu(x) d\mu(y), \quad \mu \in \mathbb{M}$$

and $\int \phi d\mu$ are finite. Also, let

$$V^\mu(z) = \int G(z, x) d\mu(x), \quad \mu \in \mathbb{M}.$$

be the Green's potential of the measure μ . The weighted energy of the field ϕ is

$$E_\phi(\mu) = E(\mu) + 2 \int \phi d\mu,$$

for any $\mu \in \mathbb{M}$.

Now, given any curve F in $\bar{\mathbb{K}}$, the equilibrium measure λ^F supported in F is defined by

$$E_\phi(\lambda^F) = \min_{\mu \in M(F)} E_\phi(\mu),$$

where $M(F)$ is the set of measures in \mathbb{M} which are supported in F , provided such a measure exists.

It turns out that the finite gap ansatz is equivalent to the existence of a so-called S-curve joining the points 0_+ and 0_- and lying entirely in $\bar{\mathbb{K}}$. By S-curve we mean an oriented curve F such that the equilibrium measure λ^F exists, its support consists of a finite union of analytic arcs and at any interior point of $\text{supp}\mu$ the so called S-property is satisfied⁴

$$\frac{d}{dn_+}(\phi + V^{\lambda^F}) = \frac{d}{dn_-}(\phi + V^{\lambda^F}),$$

The appropriate variational problem is: seek a "continuum"⁵ C such that

$$(9) \quad E_\phi(\lambda^C) = \max_{F \in \mathbb{F}} E_\phi(\lambda^F) = \max_{F \in \mathbb{F}} \min_{\mu \in M(F)} E_\phi(\mu),$$

where \mathbb{F} is the set of continua lying in $\bar{\mathbb{K}}$. The existence of a nice S-curve follows from the existence of a continuum C maximizing the equilibrium measure, in particular the associated Euler-Lagrange equations and inequalities.

⁴ $\frac{d}{dn_+}, \frac{d}{dn_-}$ are the normal outward derivatives on each side respectively

⁵ a compact connected set containing $0_+, 0_-$

Problem (9) is the non-self-adjoint analogue of the Lax-Levermore problem and a nonlinear analogue of (3).

4. JUSTIFICATION: EXISTENCE OF THE STEEPEST DESCENT PATH

EXISTENCE THEOREM [KR]. For the external field ϕ , there exists a continuum $F \in \mathbb{F}$ such that the equilibrium measure λ^F exists and

$$E_\phi[F](= E_\phi(\lambda^F)) = \max_{F \in \mathbb{F}} \min_{\mu \in M(F)} E_\phi(\mu).$$

REGULARITY THEOREM [KR]. The continuum F is in fact an S-curve, so long as it does not touch the spike $[0, iA]$ at more than a finite number of points.

If F touches the spike $[0, iA]$ at more than a finite number of points, a conceptual revision is required. We briefly discuss this issue in the next section.

Here are the main ideas of the proofs. Let ρ_0 be the distance between compact sets E, F in $\bar{\mathbb{K}}$ defined as

$$\rho_0(E, F) = \max_{z \in E} \min_{\zeta \in F} \rho_0(z, \zeta).$$

Introduce the Hausdorff metric on the set $I(\bar{\mathbb{K}})$ of closed non-empty subsets of $\bar{\mathbb{K}}$:
 $\rho_{\mathbb{K}}(A, B) = \sup(\rho_0(A, B), \rho_0(B, A)).$

Compactness of \mathbb{F} is the necessary first ingredient to prove existence of a maximizing contour. The second ingredient is semicontinuity of the energy functional that takes a given continuum F to the equilibrium energy on this continuum:

$$\mathbb{E} : F \rightarrow E_\psi[F] = E_\psi(\lambda^F) = \inf_{\mu \in M(F)} (E(\mu) + 2 \int \psi d\mu).$$

For regularity, the crucial step is

THEOREM [KR]. Let F be the maximizing continuum of and λ^F be the equilibrium measure. Let x, t be such that F does not touch the spike $[0, iA]$ at more than a finite number of points. Let μ be the extension of λ^F to the lower complex plane via $\mu(z^*) = -\mu(z)$. Then, if V is the logarithmic potential of μ ,

$$\begin{aligned} \operatorname{Re} \left(\int \frac{d\mu(u)}{u-z} + V'(z) \right)^2 &= \operatorname{Re}(V'(z))^2 - 2 \operatorname{Re} \int \frac{V'(z) - V'(u)}{z-u} d\mu(u) \\ &\quad + \operatorname{Re} \left[\frac{1}{z^2} \int 2(u+z)V'(u) d\mu(u) \right]. \end{aligned}$$

PROOF: By taking variations with respect to the equilibrium measure.

It is now easy to see that the support of the equilibrium measure of the maximizing continuum is characterized by

$$\operatorname{Re} \int^z (R_\mu)^{1/2} dz = 0,$$

$$\text{where } R_\mu(z) = (V'(z))^2 - 2 \int_{\operatorname{supp}\mu} \frac{V'(z) - V'(u)}{z - u} d\mu(u)$$

$$+ \frac{1}{z^2} \left(\int_{\operatorname{supp}\mu} 2(u + z)V'(u) d\mu(u) \right).$$

The S-property follows easily and this proves the Regularity Theorem.

It is worth mentioning here the recent work of Tovbis, Venakides and Zhou [TVZ], which examines the initial value problem for the focusing NLS (in the semiclassical limit) under two different classes of initial data. Under one of these classes, no eigenvalues exist, hence the eigenvalue spike is missing. In such a case our argument above would prove a regularity theorem without the extra assumption on the maximizing contour (that it does not touch the spike $[0, iA]$ at more than a finite number of points).

5. CROSSING THE EIGENVALUES BARRIER AND THE QUESTION OF SECONDARY CAUSTICS.

If F touches the spike $[0, iA]$ at more than a finite number of points, regularity cannot be proved as above because variations cannot be taken. In [KR] we have included a rough idea on how to extend the above proof. A more detailed argument is forthcoming [K07]. Since a complete proof is not published yet we simply summarize our general plan.

One wishes to somehow allow the contour F go through the spike $[0, iA]$. One problem arising is that (the complexification of) the external field is not analytic across the segment $[-iA, iA]$. What is true, however, is that V is analytic in a Riemann surface consisting of infinitely many sheets, cut along the line segment $[-iA, iA]$. So, the appropriate underlying space for the (doubled up) variational problem should now be a non-compact Riemann surface, say \mathbb{L} . Now, compactness

is the crucial element in the proof of a maximizing continuum. But we can indeed compactify the Riemann surface \mathbb{L} by mapping it to a subset of the complex plane and compactifying the complex plane. The other problem, of course, is whether the amended variational problem (with the modified field defined on the Riemann surface and with the possibility of F not enclosing all the original eigenvalues) is still appropriate for the semiclassical NLS. The argument goes roughly as follows:

(i) Proof of the existence of an S-curve F in \mathbb{L} along the lines of [KR].

(ii) Deformation of the original discrete Riemann-Hilbert problem to the set \hat{F} consisting of the projection of F to the complex plane. At first sight, it is clear that \hat{F} may not encircle the spike $[0, iA]$. It is however possible to append S-loops (not necessarily with respect to the same branch of the external field) and end up with a sum of S-loops, which we still denote by \hat{F} , that *does* encircle the spike $[0, iA]$. The original discrete Riemann-Hilbert problem can be trivially deformed to a discrete Riemann-Hilbert on a union of S-loops. All this is possible even in the case where \hat{F} self-intersects.

(iii) Deform the discrete Riemann-Hilbert problem to the continuous one with the right band/gap structure (on \hat{F} ; according to the projection of the equilibrium measure on F), which is then explicitly solvable via theta functions. Both the discrete-to-continuous approximation and the opening of the lenses needed for this deformation are justified as in [KMM] (see also the article [LM] mentioned below for the delicate study of the Riemann-Hilbert problem near the points where \hat{F} crosses the spike). The g-function is defined by the same Thouless-type formula with respect to the equilibrium measure (cf. section 2(iii)). It satisfies the same conditions as in [KMM] (measure reality and variational inequality) on bands (where the branch of the field turns out to be irrelevant) and on gaps (where the inequalities are satisfied according to the branch of the external field).

Once the proposed plan is implemented in detail, we will have a complete proof that the solution of the initial value problem for the focusing NLS equation admits a finite genus representation, asymptotically as $h \rightarrow 0$, at least in the case of the

simplest data $u(x, 0) = Asechx$. But the arguments above also hold for a large class of "semiclassical soliton ensembles" initial data defined precisely in section VIII of [KR].

It is worth noting here the recent paper [LM] which also addresses the issue of the target contour hitting the eigenvalue barrier. ([LM] does not really mention a variational problem and prefers to consider directly the conditions (equations and inequalities) for the g -function.) This very interesting paper does not prove the existence of an appropriate target contour but instead contains a numerical and theoretical discussion of the issue of the target contour hitting the eigenvalue barrier.

In [LM] the "band" part of the contour is defined not as the support of an equilibrium measure but instead it is considered as the trajectory of a quadratic differential (as in [KMM]). It is noted numerically that such a trajectory may hit the barrier $[0, iA]$. It is then proposed that the inequalities defining the "gap" part of the target contour be amended when it passes the barrier $[0, iA]$ and the actual amendment is justified numerically and theoretically. As a conclusion it is claimed that the mechanism of the second "caustic" (a caustic appears when the topology of \hat{F} changes as x, t vary) is different from the mechanism of the first caustic.

We simply note here that the extra conditions suggested in [LM] appear naturally in the framework we have introduced above. The different gap inequalities in [LM] correspond to gap inequalities in different sheets, as viewed from our perspective. We thus prefer to say that the mechanism for any "caustic" (caused by the change of the topology of \hat{F} as x, t vary) is essentially the same, independently of whether the maximizing contour has crossed the spike $[0, iA]$ or not. In any case the real issue here is not whether we have a first or second or higher order caustic, but whether the maximizing contour has hit the spike $[0, iA]$. It so happens that for the very specific data $Asechx$ the second caustic appears after the maximizing

contour has crossed the spike $[0, iA]$ once, but in general this does not have to be the case.⁶

6. CONCLUSION

In the asymptotic analysis of Riemann-Hilbert problems arising from integrable systems where the associated Lax operator is non-self-adjoint, the computation of non-trivial steepest descent contours is essential. The two main components of a rigorous proof of asymptotic formulae are:

(i) Proof of the existence and regularity of such steepest descent contours.

(ii) Given (i), a rigorous proof of the asymptotic validity of the deformation of the given Riemann-Hilbert problem to one with jumps across the steepest descent contour.

In this review paper we have presented some methods and results, contained in [KR] and [KMM], achieving (i) and (ii) for some specific cases of the initial value problem for the focusing integrable nonlinear Schrödinger equation in the semiclassical limit. We expect that these methods and results may be useful in the treatment of Riemann-Hilbert problems arising in the analysis of general complex or normal random matrices [WZ]. Although there have been simple cases of non-self-adjoint problems where the target contour can be explicitly computed without any recourse to a variational problem (which of course is always there; see e.g. [K96], [KSVW], [TVZ]), we believe that global results can in general only be justified by proving existence and regularity for a solution of a maximin variational problem in two dimensions.

7. REFERENCES

[BC] R.Beals, R.Coifman, Scattering and inverse scattering for first order systems. *Comm. Pure Appl. Math.* v.37 (1984), no. 1, pp.39–90.

⁶Also, it is worth recalling Chapter 6 of [KMM] where it is noticed that, depending on the choices of the parameters σ and J defined there, the maximizing contour may hit the barrier $[0, iA]$ even before the first caustic, even for the simplest *Asechx* data. So the time where the barrier is first hit is definitely not an intrinsic parameter of the original problem.

[CG] K.F.Clancey, I.Gohberg, Factorization of matrix functions and singular integral operators. *Operator Theory: Advances and Applications*, 3. Birkhäuser Verlag, Basel-Boston, Mass., 1981.

[DVZ94] P.Deift, S.Venakides, X.Zhou, The Collisionless Shock Region for the Long-Time Behavior of Solutions of the KdV equation, *Comm. Pure Appl. Math.*, v.47, 1994, pp.199-206.

[DVZ97] P.Deift, S.Venakides, X.Zhou, New Results in Small Dispersion KdV by an Extension of the Steepest Descent Method for Riemann-Hilbert Problems, *Inter.Math.Res.Notices*, 1997, pp.286-299.

[DZ] P.Deift, X.Zhou, A Steepest Descent Method for Oscillatory Riemann-Hilbert Problems, *Annals of Mathematics*, v.137, n.2, 1993, pp.295-368.

[DZ95] P.Deift, X.Zhou, Asymptotics for the Painlevé II equation, *Comm. Pure Appl. Math.* v.48, 1995, no. 3, pp.277–337.

[E] A.Erdélyi, *Asymptotic Expansions*, Dover 1956.

[I] A.R.Its, *Asymptotics of Solutions of the Nonlinear Schrödinger equation, and Isomonodromic Deformations of Systems of Linear Differential Equations*, *Sov. Math.Dokl.*, v.24, n.3, 1982, pp.14–18.

[IN] A.R.Its, V.Novokshenov, *The Isomonodromic Deformation Method in the Theory of Painlevé Equations*, *Lecture Notes in Math.*, 1191, Springer Verlag, Berlin 1986.

[K07] S.Kamvissis, in preparation.

[K96] S.Kamvissis, Long time behavior for the focusing nonlinear Schroedinger equation with real spectral singularities, *Comm. Math. Phys.* v.180, no. 2, 1996, pp.325–341.

[KMM] S.Kamvissis, K.McLaughlin, P.Miller, *Semiclassical Soliton Ensembles for the Focusing Nonlinear Schrödinger Equation*, *Annals of Mathematics Studies*, v.154, Princeton University Press, 2003.

[KR] S.Kamvissis, E.Rakhmanov, Existence and Regularity for an Energy Maximization Problem in Two Dimensions, *Jour. Math. Phys.*, v.46, n.8, 083505, 2005.

[K00] A.Kuijlaars, On the finite-gap ansatz in the continuum limit of the Toda lattice, *Duke Math. J.* v.104, no. 3, 2000, pp.433–462.

[KSVW] A.Kuijlaars, H.Stahl, W.Van Assche, F.Wielonsky, Asymptotique des approximants de Hermite-Padé quadratiques de la fonction exponentielle et problèmes de Riemann-Hilbert, *CRAS* v.336, 2003, pp.893-896.

[LL] P.D.Lax, C.D.Levermore, The Zero Dispersion Limit for the KdV Equation, I-III, *Comm. Pure Appl. Math.*, v.36, 1983, pp.253-290, pp.571-593, pp.809-829.

[LM] G.Lyng, P.Miller, The N-soliton of the focusing nonlinear Schrödinger equation for N large, *Comm. Pure Appl. Math.*, v.60, 2007, pp. 951-1026.

[TVZ] A.Tovbis, S.Venakides, X.Zhou, On semiclassical (zero dispersion limit) solutions of the focusing nonlinear Schrödinger equation, *Comm. Pure Appl. Math.* 57, n. 7, 2004, pp.877–985.

[WZ] P.Wiegmann, A.Zabrodin, Large scale correlations in normal non-Hermitian matrix ensembles, *J. Phys. A* v.36, no. 12, 2003, pp.3411–3424.

[Z] X.Zhou, The Riemann–Hilbert problem and inverse scattering, *SIAM J. Math. Anal.*, v.20, n.4, 1989, pp. 966–986.

DEPARTMENT OF APPLIED MATHEMATICS, UNIVERSITY OF CRETE, KNOSSOS, GREECE
E-mail address: `spyros@tem.uoc.gr`