

BUNDLES OF RANK 2 WITH SMALL CLIFFORD INDEX ON ALGEBRAIC CURVES

H. LANGE AND P. E. NEWSTEAD

ABSTRACT. In this paper, we construct stable bundles E of rank 2 on suitably chosen curves of any genus $g \geq 12$ with maximal Clifford index such that the Clifford index of E takes the minimum possible value for curves with this property.

1. INTRODUCTION

In a previous paper [8] (see also [4, 5, 9]), we constructed examples of curves for which the rank-2 Clifford index $\text{Cliff}_2(C)$ is strictly less than the classical Clifford index, thus producing counter-examples to a conjecture of Mercat [10]. The purpose of the present paper is to improve [8, Theorem 1.1] by substantially weakening the hypotheses; the new result is best possible and enables us to construct examples of curves C of any genus $g \geq 12$ for which the Clifford index $\text{Cliff}(C)$ takes its maximum possible value $\lfloor \frac{g-1}{2} \rfloor$, while the rank-2 Clifford index $\text{Cliff}_2(C)$ satisfies $\text{Cliff}_2(C) = \frac{1}{2} \text{Cliff}(C) + 2$, which is the minimum possible value for curves of Clifford index $\text{Cliff}(C)$.

To state the results, we recall first the definition of $\text{Cliff}_n(C)$. For any vector bundle E of rank n and degree d on C , we define

$$\gamma(E) := \frac{1}{n} (d - 2(h^0(E) - n)) = \mu(E) - 2\frac{h^0(E)}{n} + 2.$$

If C has genus $g \geq 4$, we then define, for any positive integer n ,

$$\text{Cliff}_n(C) := \min_E \left\{ \gamma(E) \mid \begin{array}{l} E \text{ semistable of rank } n \\ h^0(E) \geq 2n, \mu(E) \leq g - 1 \end{array} \right\}$$

(this invariant is denoted in [7, 8, 9] by γ'_n). Note that $\text{Cliff}_1(C) = \text{Cliff}(C)$ is the usual Clifford index of the curve C . Moreover, as observed in [7, Proposition 3.3 and Conjecture 9.3], the conjecture of [10] can be restated in a slightly weaker form as

Conjecture. $\text{Cliff}_n(C) = \text{Cliff}(C)$.

2000 *Mathematics Subject Classification.* Primary: 14H60; Secondary: 14J28.

Key words and phrases. Algebraic curve, stable vector bundle, Clifford index, K3-surface.

Both authors are members of the research group VBAC (Vector Bundles on Algebraic Curves). They would like to thank the Isaac Newton Institute, where a part of this paper was written during the Moduli Spaces programme.

In fact, for $n = 2$, this form of the conjecture is equivalent to the original (see [9, Proposition 2.7]).

Our main theorem can now be stated.

Theorem 3.3. *Suppose that g, s, d are integers such that*

$$s \geq -1, \quad g \geq 2s + 14 \quad \text{and} \quad d = g - s.$$

Then there exists a curve C of genus g having $\text{Cliff}(C) = \lfloor \frac{g-1}{2} \rfloor$ and a stable vector bundle E of rank 2 and degree d on C with $\gamma(E) = \frac{g-s}{2} - 2$. Hence

$$\text{Cliff}_2(C) \leq \frac{g-s}{2} - 2 < \text{Cliff}(C).$$

This theorem is a substantially strengthened version of [8, Theorem 1.1]; the hypotheses are now best possible in the sense that the theorem fails for $g \leq 2s + 13$. The stronger hypotheses in the original theorem were needed to ensure that certain K3-surfaces contained no effective divisors D such that $D^2 = 0$ or $D^2 = -2$. In the present paper, our K3-surfaces may contain such divisors, but we are able to control these and show that they do not affect the calculations required to prove the theorem. The proof of the theorem itself is essentially the same as that of [8, Theorem 1.1]; we give it in full for the sake of clarity and to demonstrate how the hypotheses are used.

As a corollary to Theorem 3.3 we have

Theorem 3.8. *Let γ be an integer, $\gamma \geq 5$. Then there exists a curve C with $\text{Cliff}(C) = \gamma$ such that*

$$\text{Cliff}_2(C) = \frac{\gamma}{2} + 2.$$

Moreover C can be taken to have genus either $2\gamma + 1$ or $2\gamma + 2$.

Following an extended discussion of curves on certain K3 surfaces in section 2, the proofs of the theorems are given in section 3. We finish with some open questions in section 4.

2. SOME CURVES ON A K3-SURFACE

Let g, d, s be integers with

$$(2.1) \quad d = g - s > 0, \quad g \geq 0, \quad g \geq 2s + 13 \quad \text{and} \quad (d, g) \neq (7, 4).$$

Note that

$$d^2 - 12g = g(g - 2s - 12) + s^2 > 0.$$

It follows from [6, Theorem 6.1,2] that there exists a smooth K3-surface S of type $(2, 3)$ in \mathbb{P}^4 containing a smooth curve C of genus g and degree d with

$$\text{Pic}(S) = H\mathbb{Z} \oplus C\mathbb{Z},$$

where H denotes the hyperplane bundle. In particular, we have

$$H^2 = 6, \quad C \cdot H = d \text{ and } C^2 = 2g - 2.$$

Proposition 2.1. *Suppose (2.1) holds, $g \geq 2$ and $g + s > 2$. Then the curve C is an ample divisor on S .*

Proof. We show that $C \cdot D > 0$ for any effective divisor D on S which we may assume to be irreducible. So let $D \sim mH + nC$ be an irreducible curve on S . We have

$$C \cdot D = m(g - s) + n(2g - 2).$$

Note first that, since H is a hyperplane,

$$(2.2) \quad D \cdot H = 6m + (g - s)n > 0.$$

If $m, n \geq 0$, then one of them has to be positive and then clearly $C \cdot D > 0$. The case $m, n \leq 0$ contradicts (2.2).

Suppose $m > 0$ and $n < 0$. Then, using (2.2) and (2.1), we have

$$C \cdot D = m(g - s) + n(2g - 2) > -\frac{n}{6}(g(g - 2s - 12) + s^2 + 12) > 0.$$

Finally, suppose $m < 0$ and $n > 0$. Then, since we assumed D irreducible, we have $D^2 \geq -2$ and

$$(2.3) \quad nC \cdot D = -mD \cdot H + D^2 \geq -mD \cdot H - 2 \geq -m - 2.$$

If $m \leq -3$, then $nC \cdot D > 0$. If $m = -1$, we have

$$C \cdot D = -(g - s) + n(2g - 2) \geq g + s - 2 > 0.$$

The same argument works for $m = -2$, $n \geq 2$. Finally, if $m = -2$ and $n = 1$, we have

$$D^2 = (C - 2H)^2 = 2g - 2 - 4d + 24 = 4s - 2g + 22.$$

So $D^2 = -2$ if and only if $g = 2s + 12$, contradicting (2.1). Thus $D^2 \geq 0$ and (2.3) implies that $C \cdot D > 0$. \square

We now investigate the possible existence of (-2) -curves on S . Note that, if D is an irreducible effective divisor on S , we have

$$\chi(D) = \frac{D^2}{2} + 2 \geq 1,$$

with equality if and only if D is a (-2) -curve. It follows that a fixed component of any effective divisor must be a (-2) -curve. Note that any irreducible (-2) -curve F has

$$h^0(S, F) = 1, \quad h^1(S, F) = h^2(S, F) = 0$$

(see [11]).

Proposition 2.2. *Suppose that (2.1) holds and let F be an irreducible (-2) -curve on S . Then one of the following holds:*

- $F \cdot H \geq d - 5$;

- $s = -3$, $F \cdot H = d - 6$, $F \sim C - H$;
- $s = -3$, $g \equiv 0 \pmod{3}$, $F \cdot H = d - 6$, $F \sim \frac{g}{3}H - C$;
- $s \geq -1$, $g = 4s + 16$, $F \cdot H = d - 8$, $F \sim (s + 4)H - C$;
- $s \geq 1$ and odd, $g = \frac{5}{2}(s + 5)$, $F \cdot H = d - 10$, $F \sim \frac{s+5}{2}H - C$.

Proof. Write $F \sim mH + nC$ and

$$r := F \cdot H = 6m + dn.$$

The condition $F^2 = -2$ translates to

$$3m^2 + dmn + (g - 1)n^2 = -1.$$

Inserting $m = \frac{r-dn}{6}$, this gives

$$(2.4) \quad n^2[d^2 - 12(g - 1)] = r^2 + 12.$$

Suppose first that $n^2 \geq 4$ and $r \leq d - 6$. In order to get a contradiction, it is enough to have

$$4[d^2 - 12(g - 1)] > (d - 6)^2 + 12,$$

which gives

$$d^2 + 4d > 16g.$$

Inserting $d = g - s$, this is equivalent to

$$g(g - 2s - 12) + s^2 - 4s > 0.$$

This holds by (2.1).

It remains to consider the case $n^2 = 1$. If $r \leq d - 12$, then in order to get a contradiction, it is enough to have

$$d^2 - 12(g - 1) > (d - 12)^2 + 12,$$

which means $8d > 4g + 48$. Inserting $d = g - s$, this is equivalent to $g > 2s + 12$, which is valid by (2.1).

The equation (2.4) with $n^2 = 1$ implies that $r - d$ is even. So we need to consider the cases $r = d - 6$, $r = d - 8$ and $r = d - 10$.

If $r = d - 6$, (2.4) reduces to $d = g + 3$, so $s = -3$ and $m = \frac{d-6+d}{6}$, giving the second and third cases of the statement.

Suppose $r = d - 8$. Then (2.4) says

$$d^2 - 12(g - 1) = (d - 8)^2 + 12,$$

which reduces to $4d = 3g + 16$ or equivalently to $g = 4s + 16$. If $n = 1$, the formula $m = \frac{r-d}{6}$ gives $m = -\frac{4}{3}$, a contradiction. We are left with the case $n = -1$ and

$$m = \frac{r+d}{6} = \frac{2d-8}{6} = s + 4.$$

The condition $s \geq -1$ follows from (2.1).

Finally, if $r = d - 10$, (2.4) reduces to $5d = 3g + 25$ or equivalently to $g = \frac{5}{2}(s + 5)$. So $m = \frac{d-10+d}{6}$. Again $n = 1$ gives a contradiction, so $n = -1$ and

$$m = \frac{2d - 10}{6} = \frac{s + 5}{2}.$$

The condition $s \geq 1$ follows from (2.1). \square

Corollary 2.3. *Suppose (2.1) holds with $s \geq -1$. Then the linear system $|C - H|$ is without fixed components.*

Proof. Observe first that $|C - H|$ is effective and has $h^0(C - H) \geq 3$, since $(C - H)^2 = 2s + 4 \geq 2$. Assume $|C - H|$ admits fixed components. Choose one of them and denote it by F . Note that F is a (-2) -curve. So we may write

$$C - H \sim M + F.$$

Then

$$2 < M \cdot H = (C - H) \cdot H - F \cdot H = d - 6 - F \cdot H.$$

So

$$F \cdot H \leq d - 9.$$

By Proposition 2.2, the only possibility is

$$s \geq 1, \quad g = \frac{5}{2}(s + 5), \quad F \sim \frac{s + 5}{2}H - C.$$

In this case,

$$\begin{aligned} M \cdot C &= \left(2C - \frac{s + 7}{2}H\right) \cdot C = 4g - 4 - \frac{s + 7}{2}d \\ &= 10s + 46 - \frac{s + 7}{4}(3s + 25) \\ &= -\frac{1}{4}(3s^2 + 6s - 9) \leq 0. \end{aligned}$$

This contradicts Proposition 2.1. \square

Corollary 2.4. *Suppose (2.1) holds with $s \geq -1$. Let D be an effective divisor on S with $h^0(S, D) \geq 2$ and $h^0(S, C - D) \geq 2$. Then the linear systems $|D|$ and $|C - D|$ have no fixed components.*

Proof. Since the statement is symmetric in D and $C - D$, it is sufficient to prove the corollary for $C - D$.

Suppose F is a (-2) -curve in the base locus of $|C - D|$. We may write

$$C - D \sim M + F.$$

Since $h^0(S, M) = h^0(S, C - D) \geq 2$, we have

$$3 \leq M \cdot H = (C - D) \cdot H - F \cdot H = d - D \cdot H - F \cdot H.$$

Since $h^0(S, D) \geq 2$, we have $D \cdot H \geq 3$. So

$$1 \leq F \cdot H \leq d - 6.$$

By Proposition 2.2, the case $F \cdot H = d - 6$ cannot occur since we are assuming $s \geq -1$ and we are left with the possibilities

$$(2.5) \quad g = 4s + 16, \quad F \cdot H = d - 8, \quad F \sim (s + 4)H - C$$

and

$$(2.6) \quad g = \frac{5}{2}(s + 5), \quad F \cdot H = d - 10, \quad F \sim \frac{s + 5}{2}H - C.$$

Moreover, since $|D|$ and $|C - D - F|$ are both effective, so is $|C - F|$. It follows from Proposition 2.1 that $(C - F) \cdot C > 0$.

For (2.5), we have

$$\begin{aligned} (C - F) \cdot C &= (2C - (s + 4)H) \cdot C = 4g - 4 - (s + 4)d \\ &= 16s + 60 - (s + 4)(3s + 16) \\ &= -(3s^2 + 12s + 4). \end{aligned}$$

This contradicts the fact that $(C - F) \cdot C > 0$ except when $s = -1$.

For (2.6), we argue similarly. We have

$$\begin{aligned} (C - F) \cdot C &= \left(2C - \frac{s + 5}{2}H\right) \cdot C = 2(5s + 25) - 4 - \frac{s + 5}{4}(3s + 25) \\ &= -\frac{1}{4}(3s^2 - 59). \end{aligned}$$

Since s is odd and $s \geq 1$, this is a contradiction except for $s = 1$ and $s = 3$.

This leaves us with the three possibilities

$$(2.7) \quad (g, s) = (12, -1), (15, 1), (20, 3).$$

In these cases, it is not sufficient to consider $(C - F) \cdot C$. However, in all three cases, we can show that the two conditions

$$(2.8) \quad h^0(D) \geq 2, \quad h^0(C - D - F) \geq 2$$

lead to a contradiction. Note that (2.8) implies that $D \cdot H \geq 3$ and $(C - D - F) \cdot H \geq 3$ and hence

$$(2.9) \quad F \cdot H + 3 \leq (C - D) \cdot H \leq C \cdot H - 3.$$

Similarly, using Proposition 2.1, we obtain

$$(2.10) \quad F \cdot C + 1 \leq (C - D) \cdot C \leq C \cdot C - 1.$$

Suppose first that $(g, s) = (12, -1)$, so that (2.1) and (2.5) give

$$C \cdot H = 13, \quad F \cdot H = 5, \quad F \sim 3H - C, \quad F \cdot C = 17.$$

Writing $C - D \sim mH + nC$, (2.9) and (2.10) give

$$(2.11) \quad 8 \leq 6m + 13n \leq 10$$

and

$$(2.12) \quad 18 \leq 13m + 22n \leq 21.$$

Now $13 \times (2.11) - 6 \times (2.12)$ gives

$$-22 \leq 37n \leq 22,$$

so $n = 0$. But now (2.11) gives an immediate contradiction.

Next suppose that $(g, s) = (15, 1)$. Then (2.1) and (2.6) give

$$C \cdot H = 14, F \cdot H = 4, F \sim 3H - C, F \cdot C = 14.$$

So (2.10) gives

$$(2.13) \quad 15 \leq 14m + 28n \leq 27.$$

Since $14m + 28n$ is divisible by 14, this is an immediate contradiction.

The final case $(g, s) = (20, 3)$ is a little more complicated (but also more interesting). Here (2.1) and (2.6) give

$$C \cdot H = 17, F \cdot H = 7, F \sim 4H - C, F \cdot C = 30.$$

So (2.9) and (2.10) give

$$(2.14) \quad 10 \leq 6m + 17n \leq 14$$

and

$$(2.15) \quad 31 \leq 17m + 38n \leq 37.$$

Now $17 \times (2.14) - 6 \times (2.15)$ gives

$$-52 \leq 61n \leq 52,$$

i.e. $n = 0$. Now (2.14) gives $m = 2$, which also satisfies (2.15). Hence we must have $C - D \sim 2H$. But then $|C - D|$ does not have a fixed component. This is a contradiction. \square

We now consider curves D on S with $D^2 = 0$.

Proposition 2.5. *Suppose that (2.1) holds with $s \geq -1$ and let D be an effective divisor with $D^2 = 0$ and without fixed components. Then $D \sim rE$ for some integer r , where E is irreducible with $E^2 = 0$ and $D = E_1 + \dots + E_r$ with $E_i \sim E$. Moreover one of the following holds:*

- $s \geq 0, g = 4s + 13, E \sim (s + 3)H - C$ or $E \sim 3C - 4H$;
- $s \geq 4$ and even, $g = \frac{5s}{2} + 11, E \sim \frac{s+4}{2}H - C$ or $E \sim 3C - 5H$.

Proof. By a result in [11] (see [3, Proposition 2.1] for a statement), $D = E_1 + \dots + E_r \sim rE$ as in the statement. We need only check that E has one of the stated forms. For this, let $E \sim mH + nC$, so that

$$(2.16) \quad E^2 = 6m^2 + 2dmn + (2g - 2)n^2.$$

For an integer solution of the equation $E^2 = 0$, we require the discriminant $d^2 - 6(2g - 2)$ of (2.16) to be a perfect square. So suppose

$$d^2 - 6(2g - 2) = g^2 - (2s + 12)g + s^2 + 12 = t^2$$

for some $t \geq 0$, i.e.

$$(2.17) \quad (g - s - 6)^2 - t^2 = 12s + 24.$$

Write $g - s - 6 = t + 2b$. Since $s \geq -1$, (2.1) implies that $b > 0$ and $t \geq \max\{s + 7 - 2b, 0\}$. The equation (2.17) gives

$$(2.18) \quad b(t + 2b) = 3s + 6 + b^2,$$

so that

$$(2.19) \quad b^2 \geq b(s + 7) - 3s - 6$$

On the other hand, since $bt \geq 0$, $b^2 = 3s + 6 - bt \leq 3s + 6$. Combining this with (2.19), we get

$$(2.20) \quad b(s + 7) \leq 6s + 12.$$

If $b \geq 6$, (2.20) gives an immediate contradiction. For $3 \leq b \leq 5$, we can calculate t directly from (2.18) and show that $t + 2b < s + 7$. This leaves us with $b = 1$ and $b = 2$.

When $b = 1$, (2.18) gives $t = 3s + 5$ and $g = t + 2b + s + 6 = 4s + 13$. The equation $E^2 = 0$ (see (2.16)) now gives

$$\frac{m}{n} = \frac{-d \pm t}{6} = -\frac{4}{3} \text{ or } -(s + 3).$$

When $b = 2$, we get similarly $t = \frac{3s+2}{2}$, $g = \frac{5s}{2} + 11$ and

$$\frac{m}{n} = -\frac{5}{3} \text{ or } -\frac{s+4}{2}.$$

The restrictions on s come from (2.1). To see in each case that there is an effective divisor E in the given divisor class, one checks that $E \cdot H > 0$. Since E is primitive, it must also be irreducible. \square

Corollary 2.6. *Suppose that (2.1) holds with $s \geq -1$ and that D and $C - D$ are effective divisors without fixed components. Then*

- (i) $D^2 \neq 0$, $(C - D)^2 \neq 0$;
- (ii) $h^0(C, D|_C) = h^0(S, D) = \frac{D^2}{2} + 2$.

Proof. (i) Suppose that $(C - D)^2 = 0$. By the proposition, we have $C - D = rE$ with E as in the statement. Moreover $r \geq 1$ since $C - D$ is effective and $E \cdot C \geq 0$ (in fact $E \cdot C > 0$ in view of Proposition 2.1). Since also $E^2 = 0$, we have

$$D^2 = C^2 - 2rE \cdot C = C \cdot (C - 2rE) \leq C \cdot (C - 2E).$$

Using the values of E from the proposition, we see that $D^2 < 0$, contradicting the assumption that D has no fixed components. Interchanging D and $C - D$ in this argument, we obtain a similar contradiction when $D^2 = 0$.

(ii) By (i), $(C - D)^2 > 0$, so the results of [11] ([3, Proposition 2.1]) apply to show that the general member of $|C - D|$ is smooth and irreducible and

$$h^1(S, D - C) = h^1(S, C - D) = 0.$$

Moreover, $D - C$ is not effective, so $h^0(S, D - C) = 0$. The first equality in (ii) now follows from the cohomology sequence

$$0 \rightarrow H^0(S, D - C) \rightarrow H^0(S, D) \rightarrow H^0(C, D|_C) \rightarrow H^1(S, D - C).$$

For the second equality, we note that (i) implies that $h^1(S, D) = 0$ and $h^2(S, D) = h^0(S, -D) = 0$, so

$$h^0(S, D) = \chi(D) = \frac{D^2}{2} + 2.$$

□

3. PROOF OF THEOREMS

In this section we prove our main theorems. We start with a lemma.

Lemma 3.1. *Suppose that (2.1) holds with $s \geq -1$. Then $H|_C$ is a generated line bundle on C with $h^0(C, H|_C) = 5$ and*

$$S^2 H^0(C, H|_C) \rightarrow H^0(C, H^2|_C)$$

is not injective.

Proof. Consider the exact sequence

$$0 \rightarrow \mathcal{O}_S(H - C) \rightarrow \mathcal{O}_S(H) \rightarrow \mathcal{O}_C(H|_C) \rightarrow 0.$$

$H - C$ is not effective, since $(H - C) \cdot H = 6 - d < 0$. So we have

$$0 \rightarrow H^0(S, H) \rightarrow H^0(C, H|_C) \rightarrow H^1(S, H - C) \rightarrow 0.$$

Now

$$(C - H)^2 = 2g - 2 - 2d + 6 = 2s + 4 \geq 2,$$

from which it follows that $|C - H|$ is effective. Since $|C - H|$ has no fixed component by Corollary 2.3, it follows that its general element is smooth and irreducible (see [11] or [3, Proposition 2.1]). Hence $h^1(S, H - C) = 0$ and therefore $h^0(C, H|_C) = h^0(S, H) = 5$. The last assertion follows from the fact that S is contained in a quadric. □

Corollary 3.2. *Suppose that (2.1) holds with $s \geq -1$ and $\text{Cliff}(C) = \lfloor \frac{g-1}{2} \rfloor$. Then there exists a stable vector bundle of rank 2 and degree $g - s$ on C with $h^0(E) = 4$.*

Proof. Note that $g - s < 2(\text{Cliff}(C) + 2)$. The result now follows from the lemma and [8, Lemma 3.3]. □

Theorem 3.3. *Suppose that g, s, d are integers such that*

$$(3.1) \quad s \geq -1, \quad g \geq 2s + 14 \quad \text{and} \quad d = g - s.$$

Then there exists a curve C of genus g having $\text{Cliff}(C) = \lfloor \frac{g-1}{2} \rfloor$ and a stable vector bundle E of rank 2 and degree d on C with $\gamma(E) = \frac{g-s}{2} - 2$. Hence

$$\text{Cliff}_2(C) \leq \frac{g-s}{2} - 2 < \text{Cliff}(C).$$

Proof. Let S and C be as at the beginning of section 2. In view of Corollary 3.2, it is sufficient to prove that $\text{Cliff}(C) = \left\lfloor \frac{g-1}{2} \right\rfloor$. Since C is ample by Proposition 2.1, it follows from [1, Proposition 3.3] that $\text{Cliff}(C)$ is computed by a pencil. If $\text{Cliff}(C) < \left\lfloor \frac{g-1}{2} \right\rfloor$, it then follows from [2] (see also [3, Proposition 3.1]) that there is an effective divisor D on S such that $D|_C$ computes $\text{Cliff}(C)$ and satisfying

$$(3.2) \quad h^0(S, D) \geq 2, \quad h^0(S, C - D) \geq 2 \quad \text{and} \quad \deg(D|_C) \leq g - 1.$$

By Corollaries 2.4 and 2.6, we have

$$\text{Cliff}(C) = \text{Cliff}(D|_C) = D \cdot C - D^2 - 2.$$

To obtain a contradiction, it is therefore sufficient to prove that

$$D \cdot C - D^2 - 2 \geq \left\lfloor \frac{g-1}{2} \right\rfloor.$$

Writing $D \sim mH + nC$ with $m, n \in \mathbb{Z}$, we have $D \cdot C - D^2 - 2 = f(m, n)$, where

$$f(m, n) := -6m^2 + (1 - 2n)dm + (n - n^2)(2g - 2) - 2.$$

We therefore require to prove that

$$(3.3) \quad f(m, n) \geq \left\lfloor \frac{g-1}{2} \right\rfloor.$$

By Corollaries 2.4 and 2.6, we have $D^2 > 0$. Also, by (3.2), $D \cdot H \geq 3$ and $(C - D) \cdot H \geq 3$, hence $D \cdot H \leq d - 3$. These inequalities and $\deg(D|_C) \leq g - 1$ translate to

$$(3.4) \quad 3m^2 + mnd + n^2(g - 1) > 0,$$

$$(3.5) \quad 3 \leq 6m + nd \leq d - 3,$$

$$(3.6) \quad md + (2n - 1)(g - 1) \leq 0.$$

We shall prove that (3.4) – (3.6) imply (3.3).

Denote by

$$a := \frac{1}{6}(d + \sqrt{d^2 - 12(g - 1)}) \quad \text{and} \quad b := \frac{1}{6}(d - \sqrt{d^2 - 12(g - 1)})$$

the solutions of the equation $6x^2 - 2dx + 2g - 2 = 0$. Note that $d^2 > 12(g - 1)$. So a and b are positive real numbers; moreover, substituting $g = d + s$, we see that, since $s \geq -1$ and $d \geq s + 14$,

$$(d - 12)^2 < d^2 - 12(g - 1) < (d - 6)^2.$$

Hence

$$(3.7) \quad 1 < b < 2.$$

Moreover, if $n \neq 0$, (3.4) holds if and only if

$$(3.8) \quad \frac{m}{n} < -a \quad \text{or} \quad \frac{m}{n} > -b.$$

If $n < 0$ and $\frac{m}{n} > -b$, then (3.5) implies that $3 < n(d - 6b) < 0$, because $n < 0$ and $d - 6b = \sqrt{d^2 - 12(g - 1)} > 0$, which gives a contradiction. Similarly, if $n > 0$ and $\frac{m}{n} < -a$, we obtain $3 < n(d - 6a) < 0$, again a contradiction. In view of (3.8), it remains to consider the three possibilities

- $n < 0, m > -an$;
- $n > 0, m > -bn$;
- $n = 0$.

In each case, we use (3.6) to prove (3.3).

If $n < 0$ and $m > -an$, we get from (3.6)

$$-an < m \leq \frac{(g-1)(1-2n)}{d} < \frac{(1-2n)d}{12},$$

since $d^2 > 12(g-1)$. For a fixed n , $f(m, n)$ is strictly increasing as a function of m for $m \leq \frac{(1-2n)d}{12}$ and therefore

$$\begin{aligned} f(m, n) &> f(-an, n) \\ &= \frac{d^2 - 12(g-1) + d\sqrt{d^2 - 12(g-1)}}{6} \cdot (-n) - 2 \\ &\geq \frac{d^2 - 12(g-1) + d\sqrt{d^2 - 12(g-1)}}{6} - 2. \end{aligned}$$

The inequality (3.3) therefore holds if

$$d^2 - 15g + 3 + d\sqrt{d^2 - 12(g-1)} \geq 0.$$

Since $d^2 > 12g$, it is therefore sufficient to prove that

$$d^2 - 15g + 3d + 3 \geq 0,$$

or equivalently $g(g - 2s - 12) + s^2 - 3s + 3 \geq 0$. This is certainly true under our hypotheses.

If $n > 0$ and $m > -bn$, (3.6) and (3.7) give

$$(3.9) \quad -(2n-1) \leq m \leq -\frac{(g-1)(2n-1)}{d}.$$

For a fixed $n \geq 1$, $f(m, n)$ is strictly decreasing for $m \geq -\frac{(2n-1)d}{12}$ and hence throughout the range (3.9) (whenever this range is non-empty).

So

$$\begin{aligned}
f(m, n) - \frac{g-1}{2} &\geq f\left(-\frac{(g-1)(2n-1)}{d}, n\right) - \frac{g-1}{2} \\
&= \frac{g-1}{2}(2n-1)^2 \left(1 - \frac{12(g-1)}{d^2}\right) - 2 \\
&\geq \frac{g-1}{2}(2n-1)^2 \left(1 - \frac{g-1}{g}\right) - 2 \\
&= \frac{g-1}{2g}(2n-1)^2 - 2 \geq 0 \text{ for } n \geq 2.
\end{aligned}$$

If $n = 1$, then (3.9) gives $m = -1$ and

$$(3.10) \quad f(-1, 1) = d - 8 \geq \left\lceil \frac{g-1}{2} \right\rceil \text{ for } g \geq 2s + 14.$$

Finally, suppose $n = 0$. Then

$$f(m, 0) = -6m^2 + dm - 2.$$

As a function of m this takes its maximum value at $\frac{d}{12}$. By (3.4) and (3.6),

$$1 \leq m \leq \frac{g-1}{d} \leq \frac{d}{12}.$$

So $f(m, 0)$ takes its minimal value in the allowable range at $m = 1$. Hence

$$(3.11) \quad f(m, 0) \geq f(1, 0) = d - 8 \geq \left\lceil \frac{g-1}{2} \right\rceil \text{ for } g \geq 2s + 14.$$

□

Remark 3.4. The case $s = -1$, g even, is [4, Theorem 3.7]. The case $s = -2$, g odd (not included in our theorem) is [4, Theorem 1.4].

Remark 3.5. The result of Theorem 3.3 is best possible in the sense that it fails for $g = 2s + 13$. In this case

$$\gamma(E) = \frac{g-s}{2} - 2 < \frac{1}{2} \left\lceil \frac{g-1}{2} \right\rceil + 2,$$

which contradicts [7, Proposition 3.8] if $\text{Cliff}(C) = \left\lceil \frac{g-1}{2} \right\rceil$. The points of failure in the proof are when $(m, n) = (1, 0)$ and $(m, n) = (-1, 1)$ (see (3.10) and (3.11)), i.e. for $D \sim H$ and $D \sim C - H$. In fact $H|_C$ contributes to $\text{Cliff}(C)$, so, when $g = 2s + 13$,

$$\text{Cliff}(C) \leq d - 8 < \left\lceil \frac{g-1}{2} \right\rceil.$$

When $g = 2s + 14$ or $g = 2s + 15$, we have $d = \text{Cliff}(C) + 8$, so that $H|_C$ computes the Clifford index. Thus $\text{Cliff}(C)$ is realised by an embedding of C in \mathbb{P}^4 , although the Clifford dimension of C is 1, i.e. $\text{Cliff}(C)$ is computed by a pencil (a fact used in the proof of Theorem 3.3).

Corollary 3.6. *For $g \geq 12$, there exists a curve C of maximal Clifford index $\lfloor \frac{g-1}{2} \rfloor$ such that*

$$\text{Cliff}_2(C) = \frac{1}{2} \left\lfloor \frac{g-1}{2} \right\rfloor + 2.$$

Proof. Taking $s = \lfloor \frac{g-14}{2} \rfloor$ in the theorem, we obtain

$$\text{Cliff}_2(C) \leq \frac{g-s}{2} - 2 = \frac{1}{2} \left\lfloor \frac{g-1}{2} \right\rfloor + 2 = \frac{1}{2} \text{Cliff}(C) + 2.$$

For the opposite inequality, see [7, Proposition 3.8]. \square

Remark 3.7. The result also holds for $g = 11$ [4, Theorem 1.4]. For $g \leq 10$, we have $\text{Cliff}(C) \leq 4$ for all C and $\text{Cliff}_2(C) = \text{Cliff}(C)$ by [7, Proposition 3.8].

Finally, we can express Corollary 3.6 in terms of $\text{Cliff}(C)$ rather than g . Although this is technically a corollary of Theorem 3.3, it is of sufficient interest for us to state it as a theorem.

Theorem 3.8. *Let γ be an integer, $\gamma \geq 5$. Then there exists a curve C with $\text{Cliff}(C) = \gamma$ such that*

$$\text{Cliff}_2(C) = \frac{\gamma}{2} + 2.$$

Moreover C can be taken to have genus either $2\gamma + 1$ or $2\gamma + 2$.

Proof. For $\gamma \geq 6$, this is a restatement of Corollary 3.6. For $\gamma = 5$, we need also Remark 3.7. \square

4. OPEN QUESTIONS

The following question (Mercat's conjecture for rank 2 and general C – see [10] and [9, Proposition 2.7]) remains open.

Question 4.1. *Is it true that $\text{Cliff}_2(C) = \text{Cliff}(C)$ for the general curve C of any genus?*

Farkas and Ortega conjectured in [4] that the answer to this question is yes and proved this for $g \leq 19$ (for a proof when $g \leq 16$, see [4, Theorem 1.7]). If the answer is yes, we can ask a more precise question, the answer to which is known only for $g \leq 10$ (or equivalently $\text{Cliff}(C) \leq 4$ (see [7, Proposition 3.8])).

Question 4.2. *Is it true that $\text{Cliff}_2(C) = \text{Cliff}(C)$ whenever C is a Petri curve?*

It may be noted that none of the curves constructed in this paper or in [4, 5, 8, 9] is general (they all lie on K3 surfaces with Picard number 2). Some of the curves are definitely not Petri (in particular those of Corollary 3.6 and Theorem 3.8); however it remains possible that some are Petri.

Note also that, for any γ , there exist curves with

$$\text{Cliff}_2(C) = \text{Cliff}(C) = \gamma$$

(for example, smooth plane curves of degree $\gamma + 4$ – see [7, Proposition 8.1]).

Question 4.3. *Suppose $\frac{\gamma}{2} + 2 < \gamma' < \gamma$. Does there exist a curve C with $\text{Cliff}(C) = \gamma$ and $\text{Cliff}_2(C) = \gamma'$?*

REFERENCES

- [1] C. Ciliberto, G. Pareschi: *Pencils of minimal degree on curves on a K3-surface*. J. reine angew. Math. 460 (1995), 15-36.
- [2] R. Donagi, D. Morrison: *Linear systems on K3-sections*. J. Diff. Geom. 29 (1989), 49-64.
- [3] G. Farkas: *Brill-Noether loci and the gonality stratification of \mathcal{M}_g* . J. reine angew. Math. 539 (2001), 185-200.
- [4] G. Farkas, A. Ortega: *The minimal resolution conjecture and rank two Brill-Noether theory*. arXiv:1010.4060. To appear in Pure and Appl. Math. Quarterly 7, no. 4 (2011).
- [5] G. Farkas and A. Ortega: *Higher rank Brill-Noether theory on sections of K3 surfaces*. arXiv:1102.0276.
- [6] A. Knutsen: *Smooth curves on projective K3-surfaces*. Math. Scandinavia 90 (2002), 215-231.
- [7] H. Lange and P. E. Newstead: *Clifford Indices for Vector Bundles on Curves*. in: A. Schmitt (Ed.) Affine Flag Manifolds and Principal Bundles. Trends in Mathematics, 165-202. Birkhäuser (2010).
- [8] H. Lange and P. E. Newstead: *Further examples of stable bundles of rank 2 with 4 sections*. arXiv:1011.0849. To appear in Pure and Appl. Math. Quarterly 7, no. 4 (2011).
- [9] H. Lange and P. E. Newstead: *Vector bundles of rank 2 computing Clifford indices*. arXiv 1012.1469.
- [10] V. Mercat: *Clifford's theorem and higher rank vector bundles*. Int. J. Math. 13 (2002), 785-796.
- [11] B. Saint-Donat: *Projective models of K3-surfaces*. Amer. J. Math. 96 (1974), 602-639.

H. LANGE, DEPARTMENT MATHEMATIK, UNIVERSITÄT ERLANGEN-NÜRNBERG,
BISMARCKSTRASSE 1 $\frac{1}{2}$, D-91054 ERLANGEN, GERMANY

E-mail address: lange@mi.uni-erlangen.de

P.E. NEWSTEAD, DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY
OF LIVERPOOL, PEACH STREET, LIVERPOOL L69 7ZL, UK

E-mail address: newstead@liv.ac.uk