

TRANSVERSAL MULTILINEAR RADON-LIKE TRANSFORMS: LOCAL AND GLOBAL ESTIMATES

JONATHAN BENNETT, NEAL BEZ, AND SUSANA GUTIÉRREZ

ABSTRACT. We prove local “ L^p -improving” estimates for a class of multilinear Radon-like transforms satisfying a strong transversality hypothesis. As a consequence, we obtain sharp multilinear convolution estimates for measures supported on fully transversal submanifolds of euclidean space of arbitrary dimension. We also prove global estimates for the same class of Radon-like transforms under a natural homogeneity assumption.

1. INTRODUCTION

The main purpose of this paper is to obtain local and global estimates for a class of multilinear Radon-like transforms satisfying a transversality hypothesis.

A popular description of a Radon-like transform is a mapping R of the form

$$Rf(x) = \int_{\mathbb{R}^d} f(y) \delta(F(y, x)) \psi(y, x) dy,$$

where $f : \mathbb{R}^d \rightarrow \mathbb{C}$ is a suitable test function, $x \in \mathbb{R}^n$ and $F : \mathbb{R}^d \times \mathbb{R}^n \rightarrow \mathbb{R}^k$ a suitably smooth function which typically satisfies some nondegeneracy condition on the support of the cutoff function ψ . Here $d, k, n \in \mathbb{N}$ and δ denotes the Dirac delta distribution on \mathbb{R}^k . If ∇F does not vanish then $\delta \circ F$ is easily seen to be a well-defined distribution. Notice that $Rf(x)$ may be interpreted as a surface integral (or “average”) of f over the submanifold

$$M_x := \{y \in \mathbb{R}^d : F(y, x) = 0\},$$

which generically has dimension $d-k$. It is natural to seek so-called “ L^p -improving” properties of such transforms; that is, given F find the exponents p and q for which R extends to a bounded mapping from $L^p(\mathbb{R}^d)$ into $L^q(\mathbb{R}^n)$. There is a considerable literature on such problems which we do not discuss here, although the interested reader should consult the paper of Tao and Wright [13].

A natural description of a *multilinear* Radon-like transform is a mapping R of the form

$$(1) \quad Rf(x) = \int_{\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m}} f_1(y_1) \cdots f_m(y_m) \delta(F(y, x)) \psi(y, x) dy,$$

where $f = (f_j)_{j=1}^m$, $f_j : \mathbb{R}^{d_j} \rightarrow \mathbb{C}$ is a suitable test function, $x \in \mathbb{R}^n$ and $F : \mathbb{R}^n \times \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m} \rightarrow \mathbb{R}^k$ is a suitably smooth function which typically satisfies

Date: 11 July 2011.

some nondegeneracy conditions on the support of the cutoff function ψ . Again it is natural to seek $L^{p_1}(\mathbb{R}^{d_1}) \times \dots \times L^{p_m}(\mathbb{R}^{d_m}) \rightarrow L^q(\mathbb{R}^n)$ estimates for R . By duality these qualities may be expressed as bounds on multilinear forms such as

$$(2) \quad \int_{\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_{m+1}}} \prod_{j=1}^{m+1} f_j(y_j) \delta(F(y)) \psi(y) \, dy \leq C \prod_{j=1}^{m+1} \|f_j\|_{L^{p_j}(\mathbb{R}^{d_j})}.$$

Estimates of the form (2) arise frequently in problems in a variety of fields including geometric and harmonic analysis, and dispersive PDE. Often these are manifested as certain multilinear singular convolution inequalities, which we now describe. In the work of Tao, Vargas and Vega [12] it was shown that whenever S_1 and S_2 are transversal compact submanifolds of \mathbb{R}^d , where $d \geq 2$, which are smooth with nonvanishing gaussian curvature, we have the estimate

$$(3) \quad \|f_1 d\sigma_1 * f_2 d\sigma_2\|_{L^2(\mathbb{R}^2)} \leq C \|f_1\|_{L^{\frac{4d}{3d-2}}(d\sigma_1)} \|f_2\|_{L^{\frac{4d}{3d-2}}(d\sigma_2)}.$$

Here, $d\sigma_j$ is the measure supported on S_j given by

$$\int_{\mathbb{R}^d} f(x) \, d\sigma_j(x) = \int_{U_j} f(\Sigma_j(x')) \, dx',$$

where $\Sigma_j : U_j \rightarrow \mathbb{R}^d$ parametrises S_j for some compact subset U_j of \mathbb{R}^{d-1} . See [8] for the case $d = 3$ on which [12] built. The estimate in (3) was obtained in [12] from a $L^{d/d-1}(\mathbb{R}^d) \rightarrow L^d(\mathbb{R}^d)$ estimate on the Radon-like transform in (1) with $m = 1$ and where F satisfies a rotational curvature condition on the support of the cutoff. By Plancherel's theorem, (3) immediately implies a bilinear adjoint Fourier restriction estimate for transversal compact subsets of surfaces given by the graph of an elliptic phase (such as a paraboloid).

At higher levels of multilinearity, in particular when the level coincides with the ambient dimension, transversality is key and additional curvature hypotheses do not increase the L^p improving nature of the singular convolution operation. It is known that if S_1, \dots, S_d are transversal $C^{1,\beta}$ codimension-one submanifolds of \mathbb{R}^d at the origin, $1 \leq q \leq \infty$ and $p'_j \leq (d-1)q'$, then

$$(4) \quad \|f_1 d\sigma_1 * \dots * f_d d\sigma_d\|_{L^q(\mathbb{R}^d)} \leq C \prod_{j=1}^d \|f_j\|_{L^{p_j}(d\sigma_j)}$$

if each $f_j \in L^{p_j}(d\sigma_j)$ has support sufficiently close to the origin. This follows from [7] when $d = 3$, and [5] for $d \geq 4$. In [4] the case $d = 3$, $q = \infty$ was considered under certain scaleable assumptions at each point of the hypersurfaces.

A major goal of this paper is to provide a certain generalisation of (4) to transversal submanifolds of *general codimension*. This will be a consequence of Theorem 1.2 below concerning local multilinear Radon-like transform estimates. To be precise, let U_j be a compact subset of \mathbb{R}^{d_j} and let $\Sigma_j : U_j \rightarrow \mathbb{R}^d$ parametrise a $C^{1,\beta}$ and d_j -dimensional submanifold S_j of \mathbb{R}^d , where $d \geq 2$ and $1 \leq j \leq m$. As above, define the associated measure $d\sigma_j$ on \mathbb{R}^d by

$$\int_{\mathbb{R}^d} f(x) \, d\sigma_j(x) = \int_{U_j} f(\Sigma_j(y)) \, dy.$$

We say that S_1, \dots, S_m are *fully transversal* at the origin if

$$(5) \quad \bigoplus_{j=1}^m \ker(d\Sigma_j(0))^* = \mathbb{R}^d.$$

Theorem 1.1. *Suppose that the submanifolds S_1, \dots, S_m are fully transversal at the origin. If $1 \leq q \leq \infty$ and $p'_j \leq (m-1)q'$, then there exists a constant C such that*

$$(6) \quad \|f_1 d\sigma_1 * \dots * f_m d\sigma_m\|_{L^q(\mathbb{R}^d)} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(d\sigma_j)}$$

for all $f_j \in L^{p_j}(d\sigma_j)$ supported in a sufficiently small neighbourhood of the origin.

The case $m = 3$ was considered by Bejenaru–Herr [2] (albeit under certain scaleable assumptions over the entire patches U_j in the spirit of [4]) in proving certain well-posedness results for the three-dimensional Zakharov system. Also, when $d_j = d-1$ for each $1 \leq j \leq m$, Theorem 1.1 was proved in [5].

In Theorem 1.1, the most interesting case is where $p'_j = (m-1)q'$ for each j since, of course, the remaining cases follow from Hölder’s inequality. We shall discuss the optimality of the estimates given by Theorem 1.1 later in Section 3, including a justification that given, $1 \leq q \leq \infty$, the exponent p satisfying $p' = (m-1)q'$ cannot be improved.

Under the hypotheses of Theorem 1.1, taking $q = 2$ and via Plancherel’s theorem, we obtain the estimate

$$\left\| \prod_{j=1}^m \widehat{f_j d\sigma_j} \right\|_{L^2(\mathbb{R}^d)} \leq C \prod_{j=1}^m \|f_j\|_{L^{(2m-2)'}(d\sigma_j)}$$

for all $f_j \in L^{(2m-2)'}(d\sigma_j)$ supported in a sufficiently small neighbourhood of the origin. This estimate is, of course, a certain multilinear adjoint Fourier restriction estimate for fully transversal submanifolds of \mathbb{R}^d , and extends previous results of this nature in [7] and [5].

Notice that for the transversality assumption (5) to hold it is necessary that

$$(7) \quad \sum_{j=1}^m d_j = (m-1)d.$$

It is also natural to assume that $m \geq 2$ and $1 \leq d_j \leq d-1$ for each $1 \leq j \leq m$. We shall now state our local result on Radon-like transforms, for which we shall assume the above restrictions on d, m and each d_j . In order to state the theorem, we need to introduce a little notation which shall be adopted throughout the paper.

Let

$$\mathcal{K}_j = \{d_1 + \dots + d_{j-1} + 1, \dots, d_1 + \dots + d_{j-1} + d_j\}$$

for $1 \leq j \leq m$, so that $(\mathcal{K}_j)_{j=1}^m$ partition $\{1, \dots, (m-1)d\}$. Note that the cardinality of each \mathcal{K}_j is d_j . In addition, it is natural to introduce some language from exterior algebra to express the nondegeneracy assumption on the mapping F in the distribution $\delta \circ F$. We use the standard notation $\Lambda^n(\mathbb{R}^d)$ for the n th exterior algebra of \mathbb{R}^d . Also, we use $\star : \Lambda^n(\mathbb{R}^d) \rightarrow \Lambda^{d-n}(\mathbb{R}^d)$ for the Hodge star operator.

For a *linear* map $F : \mathbb{R}^{d_1} \times \cdots \times \mathbb{R}^{d_m} \rightarrow \mathbb{R}^d$, let $Y_j(F) \in \Lambda^{d_j}(\mathbb{R}^d)$ be given by

$$(8) \quad Y_j(F) = \bigwedge_{k \in \mathcal{K}_j} F(e_k) \in \Lambda^{d_j}(\mathbb{R}^d),$$

where e_k denotes the k th standard basis vector in $\mathbb{R}^{(m-1)d} \cong \mathbb{R}^{d_1} \times \cdots \times \mathbb{R}^{d_m}$.

We shall write e_k for the k th standard basis vector throughout; the dimension shall be clear from the context. Also, with an index such as $k \in \mathcal{K}_j$, as in (8), we shall always mean that the operation is being performed as k increases over \mathcal{K}_j .

Theorem 1.2. *Let $\sum_{j=1}^m d_j = (m-1)d$. If $F : \mathbb{R}^{d_1} \times \cdots \times \mathbb{R}^{d_m} \rightarrow \mathbb{R}^d$ is $C^{1,\beta}$ in a neighbourhood of some point $y_* \in \mathbb{R}^{d_1} \times \cdots \times \mathbb{R}^{d_m}$ for some $\beta > 0$ and*

$$(9) \quad \star \bigwedge_{j=1}^m \star Y_j(dF(y_*)) \neq 0,$$

then there exists a neighbourhood V of y_* and a constant C such that

$$(10) \quad \int_V \prod_{j=1}^m f_j(y_j) \delta(F(y)) \, dy \leq C \prod_{j=1}^m \|f_j\|_{L^{(m-1)' }(\mathbb{R}^{d_j})}$$

for all nonnegative $f_j \in L^{(m-1)' }(\mathbb{R}^{d_j})$, $1 \leq j \leq m$.

It is important to point out the nondegeneracy hypothesis (9) need only be imposed on some neighbourhood of the zero set of F . Some further remarks concerning Theorem 1.2 are now in order.

Firstly, suppose we have a mapping $G : (\mathbb{R}^{d-1})^{d-1} \rightarrow \mathbb{R}$ which is $C^{1,\beta}$ in a neighbourhood of a point $u_* \in (\mathbb{R}^{d-1})^{d-1}$ and satisfies the nondegeneracy assumption

$$\det(\nabla_{u_1} G(u_*) \cdots \nabla_{u_{d-1}} G(u_*)) \neq 0.$$

Then there exists a neighbourhood V of u_* and a constant C such that

$$(11) \quad \int_V f_1(u_1) \cdots f_{d-1}(u_{d-1}) f_d(u_1 + \cdots + u_{d-1}) \delta(G(u)) \, du \leq C \prod_{j=1}^d \|f_j\|_{L^{(d-1)' }(\mathbb{R}^{d-1})}$$

for all nonnegative $f_j \in L^{(d-1)' }(\mathbb{R}^{d-1})$. This was proved in [7] for $d = 3$ and [5] for $d \geq 4$, and the multilinear singular convolution estimate in (4) for hypersurfaces is a consequence of (11). Observe that (11) follows from Theorem 1.2 by taking $m = d$, $d_j = d - 1$ for each j , and $F : (\mathbb{R}^{d-1})^d \rightarrow \mathbb{R}^d$ given by

$$F(y_1, \dots, y_d) = (y_d - y_{d-1} - \cdots - y_1, G(y_1, \dots, y_{d-1})).$$

An especially elegant case of Theorem 1.2 occurs when $m = d = 3$ and $d_j = 2$ for each j . In this case, the nondegeneracy assumption (9) is simply that

$$(12) \quad \det(Y_1(dF(y_*)) \ Y_2(dF(y_*)) \ Y_3(dF(y_*))) \neq 0,$$

where

$$Y_j(dF(y_*)) = \nabla_{(y_j)_1} F(y_*) \times \nabla_{(y_j)_2} F(y_*).$$

Then Theorem 1.2 tells us that there exists a neighbourhood V of y_* in $(\mathbb{R}^2)^3$ and a constant C such that

$$\int_V f_1(y_1)f_2(y_2)f_3(y_3)\delta(F(y)) \, dy \leq C\|f_1\|_{L^2(\mathbb{R}^2)}\|f_2\|_{L^2(\mathbb{R}^2)}\|f_3\|_{L^2(\mathbb{R}^2)}$$

for all nonnegative $f_1, f_2, f_3 \in L^2(\mathbb{R}^2)$.

One interpretation of inequality (10) is that it is a distributional L^p variant of the multilinear weighted L^2 estimates of Tao [11]. At the end of this section we provide another more explicit perspective from a dispersive PDE point of view.

Next, we present our *global* extension of Theorem 1.2 under the additional hypothesis that F is homogeneous of degree one. Our original motivation for considering this setting stemmed from the appearance of certain globally-defined multilinear Radon-like transforms in diffraction tomography, and we shall elaborate on this shortly. The hypothesis that F be homogeneous of degree one is of course a natural assumption since it encompasses the case of linear F . Observe that mappings F which are homogeneous of degree one are not, in general, smooth at the origin. As a consequence, the neighbourhood V obtained from Theorem 1.2 will not contain the origin and thus a trivial globalisation argument based on a direct isotropic scaling argument will not run.

Theorem 1.3. *Suppose $\sum_{j=1}^m d_j = (m-1)d$. If $F : \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m} \rightarrow \mathbb{R}^d$ is homogeneous of degree one, with regularity $C^{1,\beta}$ for some $\beta > 0$, and*

$$(13) \quad \star \bigwedge_{j=1}^m \star Y_j(dF(\omega)) \neq 0$$

for each ω in the unit sphere, then there exists a constant C such that

$$\int_{\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m}} \prod_{j=1}^m f_j(y_j)\delta(F(y)) \, dy \leq C \prod_{j=1}^m \|f_j\|_{L^{(m-1)'}(\mathbb{R}^{d_j})}$$

for all nonnegative $f_j \in L^{(m-1)' }(\mathbb{R}^{d_j})$, $1 \leq j \leq m$.

There is a superficially stronger version of Theorem 1.3 where the hypothesis (13) is made on the support of a cutoff function ψ defined on the unit sphere in $\mathbb{R}^{d_1+\dots+d_m}$, and the resulting inequality is replaced by

$$\int_{\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m}} \prod_{j=1}^m f_j(y_j)\delta(F(y))\psi(y/|y|) \, dy \leq C \prod_{j=1}^m \|f_j\|_{L^{(m-1)' }(\mathbb{R}^{d_j})}.$$

This fact, which is immediately apparent from the forthcoming proof, allows Theorem 1.3 to be applied in situations where the form in (13) has vanishing points.

Again, we emphasise that (13) only need hold for each unit vector ω which belongs to some neighbourhood of the zero set of F . We can see the importance of this in the example $F : (\mathbb{R}^2)^3 \rightarrow \mathbb{R}^3$ given by

$$F(y_1, y_2, y_3) = (y_3 - y_1 - y_2, |y_1| - |y_2|)$$

since although the modulus function ceases to be smooth at the origin, F is smooth at unit vectors ω belonging to a small neighbourhood of the zero set of F .

Similar explicit examples of globally-defined multilinear Radon-like transforms arise in diffraction tomography, and it was these specific operators that inspired Theorem 1.3. A simple example arises in the theory of obstacle scattering, and in particular in the recovery of singularities (in the scale of classical Sobolev spaces) of a potential q by its so-called Born approximation q_B . As may be seen (for example in [9]), smoothing estimates for the mapping $q \mapsto q - q_B$ may be reduced to bounds on multilinear forms comprising the associated Born series. In particular, an important trilinear term takes the form

$$(14) \quad \Lambda(\widehat{q}, \widehat{q}, \widehat{q}) := \int_{(\mathbb{R}^2)^3} \frac{\widehat{q}(y_1) \widehat{q}(y_2) \widehat{q}(y_3)}{|y_1|^a |y_2|^b |y_3|^c} \delta(F(y)) \, dy,$$

where $F : (\mathbb{R}^2)^3 \rightarrow \mathbb{R}^3$ is given by

$$F(y_1, y_2, y_3) = \left(y_3 - y_1 - y_2, \left| y_1 - \frac{y_2}{2} \right| - \left| \frac{y_2}{2} \right| \right)$$

and a, b, c are real numbers. Here, as usual, \widehat{q} denotes the Fourier transform of q . As may be seen in [9], a direct and very case-specific analysis yields Sobolev norm bounds on $\Lambda(\widehat{q}, \widehat{q}, \widehat{q})$. Although the very general Theorem 1.3 succeeds in shedding some light on the underlying geometry, it (as stated at least) does not appear to be sufficiently quantitative to readily recover these results.

Our proof of Theorem 1.2 rests on the nonlinear Brascamp–Lieb inequalities proved in [5], and proceeds via a parametrisation of the support of the distribution $\delta \circ F$. Using this approach, one is led estimates of the form

$$(15) \quad \int_U \prod_{j=1}^m f_j(B_j(x)) \, dx \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j}(\mathbb{R}^{d_j})},$$

where U is a neighbourhood of \mathbb{R}^d , and $B_j : U \rightarrow \mathbb{R}^{d_j}$ are local submersions. See Section 2 for a precise statement of the inequalities from [5] that we need in this paper. We also note that L^p -improving estimates of the type (15) have been obtained by Tao and Wright [13] for $m = 2$, and Stovall [10] for $m \geq 3$; in these works, curvature plays a more prominent role than transversality, and are restricted to the case where the fibres of the underlying mappings B_j are one-dimensional.

We provide a direct proof of Theorem 1.3 in Section 4. We note that one may also obtain Theorem 1.3 as a consequence of the global nonlinear Brascamp–Lieb inequalities in the companion paper [1]. In the setting of estimates of the form (15), our globalisation argument is especially natural.

To conclude this section, we mention an interpretation of some of our results from a dispersive PDE point of view. For $n \in \mathbb{N}$, consider solutions $u_j : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ to the dispersive PDE

$$(16) \quad \partial_t u_j(t, x) = i h_j(D) u_j(t, x)$$

for $1 \leq j \leq n + 1$. Here, D is the frequency operator $i^{-1} \nabla$ and the $h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ are the dispersion relations. If the initial data are sufficiently nice (say Schwartz functions) we may write the solutions as

$$u_j(t, x) = \int_{\mathbb{R}^n} \widehat{u_j(0)}(\xi) e^{2\pi i (t h_j(\xi) + x \cdot \xi)} \, d\xi,$$

and consequently

$$\int_{\mathbb{R} \times \mathbb{R}^n} \prod_{j=1}^{n+1} u_j(t, x) dt dx = \int_{(\mathbb{R}^n)^{n+1}} \prod_{j=1}^{n+1} \widehat{u_j(0)}(\xi_j) \delta(F(\xi)) d\xi$$

where $F : (\mathbb{R}^n)^{n+1} \rightarrow \mathbb{R}^{n+1}$ is given by

$$F(\xi) = (\xi_1 + \cdots + \xi_{n+1}, h_1(\xi_1) + \cdots + h_{n+1}(\xi_{n+1})).$$

In this case, the nondegeneracy condition (9) is equivalent to the transversality of the hypersurfaces in \mathbb{R}^{n+1} which are the graphs of the h_j . If this holds then from either Theorem 1.2 or Theorem 1.3 we obtain “interaction” estimates of the type

$$(17) \quad \left| \int_{V \subseteq \mathbb{R} \times \mathbb{R}^n} \prod_{j=1}^{n+1} u_j(t, x) dt dx \right| \leq C \prod_{j=1}^{n+1} \|\widehat{u_j(0)}\|_{L^{n'}(\mathbb{R}^n)}.$$

In the case of $n + 1$ solutions of the classical Schrödinger equation, where $h_j(\xi) = -|\xi|^2$ for each j , we have the appropriate transversality by insisting that the initial data have disjoint compact supports. When $n = 2$, (17) is a certain weak form of the sharp multilinear extension estimates associated to compact subsets of the paraboloid, where the endpoint Lebesgue space on the left-hand side is L^1 .

When each $h_j(\xi) = |\xi|$ (strictly speaking, giving rise to a pseudo-differential operator associated to the wave equation), by Theorem 1.3 we may take $V = \mathbb{R} \times \mathbb{R}^n$ if we have initial data supported in disjoint conical regions.

Finally, we mention the case where $h_j(\xi) = -v_j \cdot \xi$ for some fixed $v_j \in \mathbb{R}^n$ in which case (16) is of course a transport equation. Since h_j is linear and when v_1, \dots, v_{n+1} are non-colinear we obtain global estimates in (17). However, we know that the stronger estimate

$$\left| \int_{\mathbb{R} \times \mathbb{R}^n} \prod_{j=1}^{n+1} u_j(t, x) dt dx \right| \leq C \prod_{j=1}^{n+1} \|u_j(0)\|_{L^n(\mathbb{R}^n)}$$

is true in this case, so that (17) follows by an application of the Hausdorff–Young inequality on the right-hand side. This follows because we know that the analogous improvement to Theorem 1.2 is true when F is linear (some further remarks on this can be found in Section 3). Furthermore, for such h_j , since nonnegativity is preserved by the solution map, and the solution is given by $u_j(t, x) = u_j(0)(x - v_j t)$, we may pass modulus signs through the integral on the left-hand side of (17), and the resulting estimate may be viewed as a re-interpretation of the Loomis–Whitney inequality.

Organisation of the paper. In Section 2 we prove a certain quantitative version of Theorem 1.2. In Section 3 we deduce the multilinear singular convolution estimates in Theorem 1.1. Finally, Theorem 1.3 is proved in Section 4.

Acknowledgement. We would like to thank Juan Antonio Barceló and Alberto Ruiz for drawing our attention to the form (14), and to Juan Antonio Barceló for his significant contribution during the early stages of this work. We are also grateful for the support of the Isaac Newton Institute for Mathematical Sciences where part of this work was carried out.

2. THE LOCAL CASE

We begin a quantitative version of Theorem 1.2 which extends earlier versions in [7] and [5], and from which we deduce Theorem 1.1 in Section 3.

Theorem 2.1. *Let $\beta, \varepsilon, \kappa > 0$ be given and suppose $\sum_{j=1}^m d_j = (m-1)d$. If $F : \mathbb{R}^{d_1} \times \cdots \times \mathbb{R}^{d_m} \rightarrow \mathbb{R}^d$ is such that $\|F\|_{C^{1,\beta}} \leq \kappa$ in a neighbourhood of some point $y_* \in \mathbb{R}^{d_1} \times \cdots \times \mathbb{R}^{d_m}$ and*

$$\left| \star \bigwedge_{j=1}^m \star Y_j(dF(y_*)) \right| \geq \varepsilon$$

then there exists a neighbourhood V of y_ , depending only on $\beta, \varepsilon, \kappa$ and d , and a constant C depending only on d , such that*

$$\int_V \prod_{j=1}^m f_j(y_j) \delta(F(y)) \, dy \leq C \varepsilon^{-\frac{1}{m-1}} \prod_{j=1}^m \|f_j\|_{L^{(m-1)' }(\mathbb{R}^{d_j})}$$

for all nonnegative $f_j \in L^{(m-1)' }(\mathbb{R}^{d_j})$, $1 \leq j \leq m$.

It will be clear from the proof of Theorem 2.1 that the rate of blow-up $\varepsilon^{-1/m-1}$ is sharp.

Before entering the proof of Theorem 2.1 we offer an outline in the case explicitly presented in the Introduction, where $m = d = 3$ and each $d_j = 2$. For additional simplicity, set $y_* = 0$. The first step is to reduce to a ‘‘canonical’’ mapping F via linear changes of variables. Specifically, there exist invertible matrices M, N such that if

$$\tilde{F} := N \circ F \circ M^{-1}$$

then

$$(18) \quad d\tilde{F}(0) = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{pmatrix}.$$

To see this, let $n_1, n_2, n_3 \in \mathbb{S}^2$ be given by

$$n_1 = \frac{Y_3(dF(0))}{|Y_3(dF(0))|}, \quad n_2 = \frac{Y_2(dF(0))}{|Y_2(dF(0))|}, \quad n_3 = \frac{Y_1(dF(0))}{|Y_1(dF(0))|}$$

and let N be the matrix whose i th row is equal to n_i . Furthermore, let N_1, N_2, N_3 be the matrices given by

$$N_1 = \begin{pmatrix} n_1 \\ n_2 \end{pmatrix}, \quad N_2 = \begin{pmatrix} n_1 \\ n_3 \end{pmatrix}, \quad N_3 = \begin{pmatrix} n_2 \\ n_3 \end{pmatrix}$$

and let F_1, F_2, F_3 be the matrices given by

$$F_j = \left(\nabla_{(y_j)_1} F(0) \quad \nabla_{(y_j)_2} F(0) \right)$$

for $j = 1, 2, 3$. Finally, we let M be the block diagonal matrix given by

$$M = \begin{pmatrix} E_1 N_1 F_1 & 0 & 0 \\ 0 & E_2 N_2 F_2 & 0 \\ 0 & 0 & E_3 N_3 F_3 \end{pmatrix}$$

where E_1 and E_3 are simply the identity matrices, and E_2 is the elementary matrix which reflects in the line spanned by $(1, 1)$. It is an easy check to verify that the above construction yields (18). Observe that using the scalar quadruple product formula, one obtains

$$|\det(N_j F_j)| = |Y_j(dF(0))| |\det(N)|$$

for each $j = 1, 2, 3$, which allows one to obtain the desired quantitative control.

Note that if F were the linear map given by (18) then, by parametrising the zero set of F , one is led to linear mappings $B_j : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ whose kernels are one-dimensional and form the coordinate axes. An appropriate form of the implicit function theorem allows one to handle small nonlinear perturbations of this situation, and the required multilinear Radon transform estimate follows in this simple situation from the nonlinear Loomis–Whitney inequality in [7]. At this stage, in the general case, we rely on the following nonlinear Brascamp–Lieb inequalities.

Theorem 2.2. [5] *Suppose that for each $1 \leq j \leq m$ the mappings $B_j : \mathbb{R}^d \rightarrow \mathbb{R}^{d_j}$ are $C^{1,\beta}$ submersions in a neighbourhood of a point $x_0 \in \mathbb{R}^d$. Suppose further that*

$$(19) \quad \bigoplus_{j=1}^m \ker dB_j(x_0) = \mathbb{R}^d.$$

Then there exists a neighbourhood U of x_0 and a constant C such that

$$(20) \quad \int_U \prod_{j=1}^m f_j \circ B_j \leq C \prod_{j=1}^m \|f_j\|_{L^{m-1}(\mathbb{R}^{d_j})}$$

for all nonnegative $f_j \in L^{m-1}(\mathbb{R}^{d_j})$, $1 \leq j \leq m$.

However, for most cases, the kernels of the derivatives of the mappings B_j at the point in question do not form a direct sum decomposition. So, as a cautionary note, in general, a further “tensoring” argument is needed at this stage in order to apply Theorem 2.2. For this to proceed, we combine the B_j appropriately in “block” form, to give rise to mappings which do satisfy the direct sum hypotheses of Theorem 2.2 (see [5] for the origin of this idea).

Proof of Theorem 2.1. We have $\sum_{j=1}^m d_j = (m-1)d$ in which case we frequently identify $(y_1, \dots, y_m) \in \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m}$ with $(u_1, \dots, u_{(m-1)d}) \in \mathbb{R}^{(m-1)d}$.

We first prove that the following special case implies Theorem 2.1.

Proposition 2.3. *Let $\beta, \kappa > 0$ be given and suppose $\sum_{j=1}^m d_j = (m-1)d$. If $F : \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m} \rightarrow \mathbb{R}^d$ is such that $\|F\|_{C^{1,\beta}} \leq \kappa$ in a neighbourhood of some point $y_* \in \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m}$ and $dF(y_*) : \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m} \rightarrow \mathbb{R}^d$ is given by*

$$(21) \quad (dF(y_*)u)_i = \sum_{j=0}^{m-2} u_{i+jd}$$

for $1 \leq i \leq d$, then there exists a neighbourhood V of y_ , depending only on β, κ and d , and a constant C depending only on d , such that*

$$\int_V \prod_{j=1}^m f_j(u_j) \delta(F(u)) du \leq C \prod_{j=1}^m \|f_j\|_{L^{(m-1)' }(\mathbb{R}^{d_j})}$$

for all nonnegative $f_j \in L^{(m-1)'}(\mathbb{R}^{d_j})$, $1 \leq j \leq m$.

We remark that if F satisfies (21) then $dF(y_*)$ is the augmented matrix comprising $m - 1$ copies of the identity matrix I_d , and moreover $|\star \bigwedge_{j=1}^m \star Y_j(dF(y_*))| = 1$.

The proof that Proposition 2.3 implies Theorem 2.1 is based on the change of variables outlined above. To set this up in general, for $1 \leq j \leq m$, we let

$$\mathcal{K}_j^* = \{d - (d_1^* + \cdots + d_{j-1}^* + d_j^*) + 1, \dots, d - (d_1^* + \cdots + d_{j-1}^*)\},$$

where $d_j^* := d - d_j$. Clearly, \mathcal{K}_j^* has d_j^* elements.

Now let $\{n_k : k \in \mathcal{K}_j^*\}$ be any orthonormal basis for the orthogonal complement of the space spanned by $\{dF(y_*)(e_k) : k \in \mathcal{K}_j\}$, and let N be the $d \times d$ matrix whose i th row is equal to n_i for each $1 \leq i \leq d$.

Denote by N_j the $d_j \times d$ matrix obtained by deleting from N the rows n_k for each $k \in \mathcal{K}_j^*$, and by F_j the $d \times d_j$ matrix obtained by deleting the k th column from $dF(y_*)$ for $k \notin \mathcal{K}_j$.

From Lemma A.1, for each $1 \leq j \leq m$, we have

$$(22) \quad \star Y_j(dF(y_*)) = \|Y_j(dF(y_*))\|_{\Lambda^{d_j}(\mathbb{R}^d)} \bigwedge_{k \in \mathcal{K}_j^*} n_k.$$

We shall use (22) to demonstrate various identities involving the determinants of the matrices describing the changes of variables, from which we obtain inequalities which allow us to establish the claimed dependences in the sizes of the neighbourhood and constants arising in Theorem 2.1.

It follows from (22) that

$$\left| \star \bigwedge_{j=1}^m \star Y_j(dF(y_*)) \right| = \left| \star \bigwedge_{j=1}^m \bigwedge_{k \in \mathcal{K}_j^*} n_k \right| \prod_{j=1}^m \|Y_j(dF(y_*))\|_{\Lambda^{d_j}(\mathbb{R}^d)}$$

and therefore

$$(23) \quad \left| \star \bigwedge_{j=1}^m \star Y_j(dF(y_*)) \right| = |\det(N)| \prod_{j=1}^m \|Y_j(dF(y_*))\|_{\Lambda^{d_j}(\mathbb{R}^d)}.$$

Furthermore,

$$\begin{aligned} |\det(N_j F_j)| &= \left| \left\langle \bigwedge_{k \notin \mathcal{K}_j^*} n_k, \bigwedge_{\ell \in \mathcal{K}_j} dF(y_*)(e_\ell) \right\rangle_{\Lambda^{d_j}(\mathbb{R}^d)} \right| \\ &= \left| \left\langle \bigwedge_{k \notin \mathcal{K}_j^*} a_k, Y_j(dF(y_*)) \right\rangle_{\Lambda^{d_j}(\mathbb{R}^d)} \right| \\ &= \|Y_j(dF(y_*))\|_{\Lambda^{d_j}(\mathbb{R}^d)} \left| \left\langle \bigwedge_{k \notin \mathcal{K}_j^*} n_k, \star \bigwedge_{\ell \in \mathcal{K}_j^*} n_\ell \right\rangle_{\Lambda^{d_j}(\mathbb{R}^d)} \right| \\ &= \|Y_j(dF(y_*))\|_{\Lambda^{d_j}(\mathbb{R}^d)} |\det(N)|. \end{aligned}$$

Consequently,

$$(24) \quad \prod_{j=1}^m |\det(N_j F_j)| = |\det(N)|^{m-1} \left| \star \bigwedge_{j=1}^m \star Y_j(dF(y_*)) \right|.$$

Now define M to be the block diagonal $\sum_{j=1}^m d_j \times \sum_{j=1}^m d_j$ matrix with $d_j \times d_j$ invertible matrices M_j on the diagonal. Here M_j is given by

$$M_j = E_j^{-1} A_j F_j$$

where $E_j : \mathbb{R}^{d_j} \rightarrow \mathbb{R}^{d_j}$ is given by

$$E_j(x_1, \dots, x_{d_j}) = (x_{d'_1 + \dots + d'_{j-1} + 1}, \dots, x_{d_j}, x_1, \dots, x_{d'_1 + \dots + d'_{j-1}})$$

for $2 \leq j \leq m-1$, $E_1 = I_{d_1}$ and $E_m = I_{d_m}$. Since $\sum_{\ell=1}^m d'_\ell = d$ and each $d'_\ell \geq 1$ it follows that

$$\sum_{\ell=1}^{j-1} d_\ell < (j-1)d < \sum_{\ell=1}^j d_\ell$$

for each $2 \leq j \leq m-1$; in particular, it follows that each E_j is well-defined. By construction, we have that if $\tilde{F} : \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m} \rightarrow \mathbb{R}^d$ is the map given by

$$\tilde{F} := N \circ F \circ M^{-1}$$

then $d\tilde{F}(\tilde{y}_*) : \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m} \rightarrow \mathbb{R}^d$ is the map given by

$$(d\tilde{F}(\tilde{y}_*)u)_i = \sum_{j=0}^{m-2} u_{i+jd}$$

for $1 \leq i \leq d$, where $\tilde{y}_* := My_*$.

By changes of variables,

$$\int_V \prod_{j=1}^m f_j(y_j) \delta(F(y)) dy = \frac{|\det(N)|}{|\det(M)|} \int_{M(V)} \prod_{j=1}^m (f_j \circ M_j^{-1})(y_j) \delta(\tilde{F}(y)) dy,$$

where the neighbourhood V of y_* shall be chosen momentarily. It follows from (23) that $|\det(N)|$ is bounded below by a constant depending only on ε, κ and d . Since $|\det(M)| = \prod_{j=1}^m |\det(M_j)|$ and $|\det(M_j)| = |\det(N_j F_j)|$ it follows from (24) that

$$(25) \quad |\det(M)| = |\det(N)|^{m-1} \left| \star \bigwedge_{j=1}^m \star Y_j(dF(y_*)) \right|$$

and $|\det(M)|$ is also bounded below by a constant depending only on ε, κ and d . Therefore, the operator norm of M^{-1} and, consequently, the $C^{1,\beta}$ norm of \tilde{F} are bounded above by a constant depending only on $\beta, \varepsilon, \kappa$ and d . By Proposition 2.3 there exists a neighbourhood V , depending only on $\beta, \varepsilon, \kappa$ and d , and a constant C , depending only on d , such that

$$\int_{M(V)} \prod_{j=1}^m (f_j \circ M_j^{-1})(y_j) \delta(\tilde{F}(y)) dy \leq C \prod_{j=1}^m \|f_j \circ M_j^{-1}\|_{L^{(m-1)'(\mathbb{R}^{d_j})}}.$$

Therefore,

$$\begin{aligned} \int_V \prod_{j=1}^m f_j(u_j) \delta(F(y)) \, dy &\leq C \frac{|\det(N)|}{|\det(M)|} \left(\prod_{j=1}^m |\det(M_j)| \right)^{\frac{1}{(m-1)'}} \prod_{j=1}^m \|f_j\|_{L^{(m-1)' }(\mathbb{R}^{d_j})} \\ &\leq C \varepsilon^{-\frac{1}{m-1}} \prod_{j=1}^m \|f_j\|_{L^{(m-1)' }(\mathbb{R}^{d_j})}. \end{aligned}$$

The second inequality follows from (25). This concludes our proof that Theorem 2.1 is implied by Proposition 2.3.

Proof of Proposition 2.3. Without loss of generality we suppose that $y_* = 0$ and $F(0) = 0$. Since

$$dF(0)|_{\text{span}\{e_r : (m-2)d+1 \leq r \leq (m-1)d\}} = I_d$$

we obtain from Theorem B.1 that there exists a neighbourhood W of the origin in $\mathbb{R}^{(m-2)d}$ and a mapping $\eta : W \rightarrow \mathbb{R}^d$ such that for each

$$x = (u_1, \dots, u_{m-2}, (u_{m-1})_1, \dots, (u_{m-1})_{d_{m-1}-d'_m}) \in W$$

we have $F(x, \eta(x)) = 0$ and $\eta(0) = 0$. The neighbourhood W depends only on β, κ and d , and the mapping η satisfies $\|\eta\|_{C^{1,\beta}} \leq \tilde{\kappa}$ for some constant $\tilde{\kappa}$ which depends only on β, κ and d .

Let $B_j : W \subset \mathbb{R}^{(m-2)d} \rightarrow \mathbb{R}^{d_j}$ be the map given by

$$B_j(x) = (x_{1+\sum_{\ell=1}^{j-1} d_\ell}, x_{2+\sum_{\ell=1}^{j-1} d_\ell}, \dots, x_{\sum_{\ell=1}^j d_\ell})$$

for $1 \leq j \leq m-2$,

$$B_{m-1}(x) = (x_{1+\sum_{\ell=1}^{m-2} d_\ell}, \dots, x_{(m-2)d}, -\eta_1(x), \dots, -\eta_{d'_m}(x))$$

and

$$B_m(x) = (-\eta_{d'_m+1}(x), \dots, -\eta_d(x)).$$

Let $S^{(j)}$ denotes the $(m-2)$ -tuple obtained by deleting $j-2 \pmod{m}$ and $j-1 \pmod{m}$ from the m -tuple $(1, \dots, m)$. Then define $B_j^\oplus : W \subset \mathbb{R}^{(m-2)d} \rightarrow \mathbb{R}^{d_j^\oplus}$, where

$$d_j^\oplus = \sum_{\ell=1}^{m-2} d_{S_\ell^{(j)}},$$

by

$$B_j^\oplus(x) = (B_{S_1^{(j)}}(x), \dots, B_{S_{m-2}^{(j)}}(x)).$$

Lemma 2.4. *Suppose $3 \leq j \leq m-1$. Then*

$$\begin{aligned} \ker dB_1^\oplus(0) &= \text{span}\left\{e_r : r = 1 + \sum_{\ell=1}^{m-2} d_\ell, \dots, (m-2)d\right\}, \\ \ker dB_2^\oplus(0) &= \text{span}\{e_r : r = d'_m + 1, \dots, d_1\}, \\ \ker dB_j^\oplus(0) &= \text{span}\left\{e_r - e_{r+d} : r = 1 + \sum_{\ell=1}^{j-3} d_\ell, \dots, \left(\sum_{\ell=1}^{j-1} d_\ell\right) - d\right\}, \\ \ker dB_m^\oplus(0) &= \text{span}\left\{e_r - e_{r+d} : r = 1 + \sum_{\ell=1}^{m-3} d_\ell, \dots, (m-3)d\right\} \\ &\quad + \text{span}\{e_s : s = (m-3)d + 1, \dots, d'_m + (m-3)d\}. \end{aligned}$$

Proof. Of course, for $1 \leq j \leq m-2$, B_j is linear and therefore $dB_j(0) = B_j$. By the chain rule,

$$d\eta(0) = -dF(0)|_{\text{span}\{e_r : 1 \leq r \leq (m-2)d\}}$$

because $dF(0)|_{\text{span}\{e_r : (m-2)d+1 \leq r \leq (m-1)d\}} = I_d$. Hence,

$$dB_{m-1}(0)x = \left(x_{1+\sum_{\ell=1}^{m-2} d_\ell}, \dots, x_{(m-2)d}, -\sum_{\ell=0}^{m-3} x_{1+\ell d}, \dots, -\sum_{\ell=0}^{m-3} x_{d'+\ell d}\right)$$

and

$$dB_m(0)x = \left(-\sum_{\ell=0}^{m-3} x_{d'_m+1+\ell d}, \dots, -\sum_{\ell=0}^{m-3} x_{d+\ell d}\right).$$

The claimed expression for the kernel of each $dB_j^\oplus(0)$ now follows from the fact that

$$\ker dB_j^\oplus(0) = \bigcap_{\ell=1}^{m-2} \ker dB_{S_\ell^{(j)}}(0)$$

and straightforward considerations. \square

Corollary 2.5. *We have*

$$\bigoplus_{j=1}^m \ker dB_j^\oplus(0) = \mathbb{R}^{(m-2)d}.$$

Proof. For $3 \leq j \leq m-1$, define

$$\begin{aligned} \mathfrak{B}_1 &:= \left\{e_r : r = 1 + \sum_{\ell=1}^{m-2} d_\ell, \dots, (m-2)d\right\} \\ \mathfrak{B}_2 &:= \{e_r : r = d'_m + 1, \dots, d_1\} \\ \mathfrak{B}_j &:= \left\{e_r - e_{r+d} : r = 1 + \sum_{\ell=1}^{j-3} d_\ell, \dots, \left(\sum_{\ell=1}^{j-1} d_\ell\right) - d\right\} \\ \mathfrak{B}_m &:= \left\{e_r - e_{r+d} : r = 1 + \sum_{\ell=1}^{m-3} d_\ell, \dots, (m-3)d\right\} \\ &\quad \cup \{e_s : s = (m-3)d + 1, \dots, d'_m + (m-3)d\}, \end{aligned}$$

and define

$$\mathfrak{B} = \bigcup_{j=1}^m \mathfrak{B}_j.$$

Since $\sum_{j=1}^m |\mathfrak{B}_j| = (m-2)d$ it suffices to check that $\text{span}(\mathfrak{B}) = \mathbb{R}^{(m-2)d}$. In order to show this, we prove that $e_r \in \text{span}(\mathfrak{B})$ for each $r \in \{1, \dots, (m-2)d\}$. To help clarify the notation in the rest of the proof, note that

$$\{1, \dots, (m-2)d\} = \bigcup_{j=2}^{m-1} \left\{ 1 + \sum_{\ell=1}^{j-2} d_\ell, \dots, \sum_{\ell=1}^{j-1} d_\ell \right\} \cup \left\{ 1 + \sum_{\ell=1}^{m-2} d_\ell, \dots, (m-2)d \right\}$$

and in our considerations we split

$$\left\{ 1 + \sum_{\ell=1}^{j-2} d_\ell, \dots, \sum_{\ell=1}^{j-1} d_\ell \right\} = \left\{ 1 + \sum_{\ell=1}^{j-2} d_\ell, \dots, d'_m + (j-2)d \right\} \cup \left\{ d'_m + 1 + (j-2)d, \dots, \sum_{\ell=1}^{j-1} d_\ell \right\}.$$

Case 1: $r \in \{1 + \sum_{\ell=1}^{m-3} d_\ell, \dots, d'_m + (m-3)d\}$.

Clearly we have $e_r \in \mathfrak{B}_m \subseteq \text{span}(\mathfrak{B})$ for $r \in \{(m-3)d + 1, \dots, d'_m + (m-3)d\}$. For the remaining $r \in \{1 + \sum_{\ell=1}^{m-3} d_\ell, \dots, (m-3)d\}$ we have $r + d \in \{1 + \sum_{\ell=1}^{m-2} d_\ell, \dots, (m-2)d\}$ so

$$e_r = (e_r - e_{r+d}) + e_{r+d} \in \mathfrak{B}_m + \mathfrak{B}_1 \subseteq \text{span}(\mathfrak{B}).$$

Case 2: $r \in \{1 + \sum_{\ell=1}^{j-2} d_\ell, \dots, d'_m + (j-2)d\}$, $2 \leq j \leq m-2$.

Write

$$e_r = \sum_{\ell=1}^{m-j-1} (e_{r+(\ell-1)d} - e_{r+\ell d}) + e_{r+(m-j-1)d}.$$

We have

$$r + (\ell-1)d \geq 1 + (\ell-1)d + \sum_{k=1}^{j-2} d_k \geq 1 + \sum_{k=1}^{j+\ell-3} d_k$$

and

$$r + (\ell-1)d \leq d'_m + (j+\ell-3)d \leq \left(\sum_{k=1}^{j+\ell-1} d_k \right) - d.$$

For each $1 \leq \ell \leq m-j-1$ we have $2 \leq j \leq \ell + j - 1 \leq m-2$ so

$$e_{r+(\ell-1)d} - e_{r+\ell d} \in \mathfrak{B}_{j+\ell}.$$

Similarly, we have $r + (m-j-1)d \in \{1 + \sum_{\ell=1}^{m-3} d_\ell, \dots, d'_m + (m-3)d\}$ and therefore $e_{r+(m-j-1)d} \in \text{span}(\mathfrak{B})$ by Case 1. Hence $e_r \in \text{span}(\mathfrak{B})$ for $r \in \{1 + \sum_{\ell=1}^{j-2} d_\ell, \dots, d'_m + (j-2)d\}$, where $2 \leq j \leq m-2$.

Case 3: $r \in \{d'_m + 1, \dots, d_1\}$.

Immediately we have $e_r \in \mathfrak{B}_2 \subseteq \text{span}(\mathfrak{B})$ for such r .

Case 4: $r \in \{d'_m + 1 + (j-2)d, \dots, \sum_{\ell=1}^{j-1} d_\ell\}$, $3 \leq j \leq m-1$.
Write

$$e_r = \sum_{\ell=1}^{j-2} (e_{r-(\ell-1)d} - e_{r-\ell d}) + e_{r-(j-2)d}.$$

Since

$$r - \ell d \in \left\{ 1 + \sum_{k=1}^{j-\ell-2} d_k, \dots, \left(\sum_{k=1}^{j-\ell} d_k \right) - d \right\}$$

and $r - (j-2)d \in \{d'_m + 1, \dots, d_1\}$, we get $e_{r-(\ell-1)d} - e_{r-\ell d} \in \mathfrak{B}_{j-\ell-1}$ and, by Case 3, $e_{r-(j-2)d} \in \mathfrak{B}_2$. Hence, $e_r \in \text{span}(\mathfrak{B})$ for $r \in \{d'_m + 1 + (j-2)d, \dots, \sum_{\ell=1}^{j-1} d_\ell\}$ and each $2 \leq j \leq m-1$.

Case 5: $r \in \{1 + \sum_{\ell=1}^{m-2} d_\ell, \dots, (m-2)d\}$.

Immediately we have $e_r \in \mathfrak{B}_1$ for such r .

This completes our proof of Corollary 2.5. \square

Given Corollary 2.5 we can apply Theorem 1.3 of [5], a certain quantitative version of Theorem 2.2, to obtain a neighbourhood U of the origin in $\mathbb{R}^{(m-2)d}$, depending on β, κ and d , and a constant C , depending on d , such that

$$\int_U \prod_{j=1}^m g_j(B_j^\oplus(x)) \, dx \leq C \prod_{j=1}^m \|g_j\|_{L^{m-1}(\mathbb{R}^{d_j^\oplus})}$$

for all $g_j \in L^{m-1}(\mathbb{R}^{d_j^\oplus})$. Given nonnegative $f_j \in L^{(m-1)' }(\mathbb{R}^{d_j})$ we let

$$f_j^\otimes = \bigotimes_{\ell=1}^{m-2} f_{S_\ell^{(j)}}^{1/(m-2)}$$

so that

$$\int_U \prod_{j=1}^m f_j^\otimes(B_j^\oplus(x)) \, dx = \int_U \prod_{j=1}^m f_j(B_j(x)) \, dx$$

and

$$\prod_{j=1}^m \|f_j^\otimes\|_{L^{m-1}(\mathbb{R}^{d_j^\oplus})} = \prod_{j=1}^m \|f_j\|_{L^{(m-1)' }(\mathbb{R}^{d_j})},$$

and therefore

$$\int_U \prod_{j=1}^m f_j(B_j(x)) \, dx \leq C \prod_{j=1}^m \|f_j\|_{L^{(m-1)' }(\mathbb{R}^{d_j})}.$$

Finally, since $F(x, \eta(x)) = 0$ for $x \in W \subset \mathbb{R}^{(m-2)d}$,

$$dF(0)|_{\text{span}\{e_r : (m-2)d+1 \leq r \leq (m-1)d\}} = I_d,$$

and $\|F\|_{C^{1,\beta}} \leq \kappa$, we obtain a neighbourhood $V \subset \mathbb{R}^{(m-1)d}$, depending on β, κ and d , such that

$$\int_V \prod_{j=1}^m f_j(u_j) \delta(F(u)) \, du \leq 2 \int_U \prod_{j=1}^m f_j(B_j(x)) \, dx.$$

This completes the proof of Proposition 2.3. \square

Proposition 2.3 implies Theorem 2.1 and thus our proof of Theorem 2.1 is complete. \square

3. MULTILINEAR SINGULAR CONVOLUTION

In this section, we prove Theorem 1.1 and demonstrate that the exponents are sharp.

Proof of Theorem 1.1. Recall that it suffices to handle the case where $p'_j = (m-1)q'$ for each $1 \leq j \leq m$. Furthermore, since the operator

$$(f_1, \dots, f_m) \mapsto f_1 d\sigma_1 * \dots * f_m d\sigma_m$$

is trivially bounded as a mapping $L^1(d\sigma_1) \times \dots \times L^1(d\sigma_m) \rightarrow L^1(\mathbb{R}^d)$, using multilinear interpolation, it suffices to handle the case $(p, q) = ((m-1)', \infty)$.

If $x \in \mathbb{R}^d$ then

$$\begin{aligned} f_1 d\sigma_1 * \dots * f_m d\sigma_m(x) &= \int_{\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m}} \prod_{j=1}^m f_j(\Sigma_j(y_j)) \delta(\Sigma_1(y_1) + \dots + \Sigma_m(y_m) - x) dy \\ &= \int_{\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m}} \prod_{j=1}^m g_j(y_j) \delta(F(y)) dy, \end{aligned}$$

where, of course, $g_j = f_j \circ \Sigma_j$ and

$$F(y) = \Sigma_1(y_1) + \dots + \Sigma_m(y_m) - x.$$

As a matrix, $dF(0)$ is the augmented matrix comprised of $d\Sigma_1(0), \dots, d\Sigma_m(0)$, and therefore

$$Y_j(dF(0)) = \bigwedge_{k=1}^{d_j} d\Sigma_j(0)(e_k).$$

If we take any orthonormal basis $\{n_k : k \in \mathcal{K}_j^*\}$ for the orthogonal complement of the image of $d\Sigma_j(0)$, or equivalently for $\ker(d\Sigma_j(0))^*$, then

$$\star Y_j(dF(0)) = \|Y_j(dF(0))\|_{\Lambda^{d_j}(\mathbb{R}^d)} \bigwedge_{k \in \mathcal{K}_j^*} n_k$$

by Lemma A.1. Consequently,

$$\star \bigwedge_{j=1}^m \star Y_j(dF(0)) = \prod_{j=1}^m \|Y_j(dF(0))\|_{\Lambda^{d_j}(\mathbb{R}^d)} \det(n_1 \dots n_d)$$

and by (5) this quantity is nonzero. Since $F \in C^{1,\beta}$ uniformly in x belonging to a sufficiently small neighbourhood of the origin, by Theorem 2.1, it follows that there is some constant C such that

$$\|f_1 d\sigma_1 * \dots * f_m d\sigma_m\|_{L^\infty(\mathbb{R}^d)} \leq C \prod_{j=1}^m \|f_j\|_{L^{(m-1)'}(d\sigma_j)}$$

if the support of each $f_j \in L^{(m-1)'}(d\sigma_j)$ is sufficiently close to the origin. This completes the proof when $(p, q) = ((m-1)', \infty)$ and hence Theorem 1.1 entirely. \square

To see that the exponents are sharp, suppose for a contradiction that when $1 \leq q \leq \infty$ and for some $p' > (m-1)q'$ there exists a constant C such that

$$(26) \quad \|f_1 d\sigma_1 * \cdots * f_m d\sigma_m\|_{L^q(\mathbb{R}^d)} \leq C \prod_{j=1}^m \|f_j\|_{L^{p'}(d\sigma_j)}$$

for all $f_j \in L^{p'}(d\sigma_j)$. It follows in particular that

$$(27) \quad \|f_1 d\sigma_1 * \cdots * f_m d\sigma_m\|_{L^2(\mathbb{R}^d)} \leq C \prod_{j=1}^m \|f_j\|_{L^r(d\sigma_j)}$$

for all $f_j \in L^r(d\sigma_j)$, where $r < (2m-2)'$. To see this, interpolate (26) with the $L^1(d\sigma_1) \times \cdots \times L^1(d\sigma_m) \rightarrow L^1(\mathbb{R}^d)$ estimate if $q > 2$, or with the $L^{(m-1)'}(d\sigma_1) \times \cdots \times L^{(m-1)'}(d\sigma_m) \rightarrow L^\infty(\mathbb{R}^d)$ estimate if $q < 2$. To see that (27) is false, it suffices to consider the case where the mappings $\Sigma_j : \mathbb{R}^{d_j} \rightarrow \mathbb{R}^d$ are linear and thus

$$\bigoplus_{j=1}^m \ker \Sigma_j^* = \mathbb{R}^d.$$

Setting $g_j = f_j \circ \Sigma_j$, by Plancherel's theorem and the linearity of the Σ_j we have

$$\|f_1 d\sigma_1 * \cdots * f_m d\sigma_m\|_{L^2(\mathbb{R}^d)}^2 = \int_{\mathbb{R}^d} \prod_{j=1}^m |\widehat{g}_j|^2(\Sigma_j^* x) dx$$

and therefore (27) is equivalent to

$$(28) \quad \int_{\mathbb{R}^d} \prod_{j=1}^m |\widehat{g}_j|^2(\Sigma_j^* x) dx \leq C \prod_{j=1}^m \|g_j\|_{L^r(\mathbb{R}^{d_j})}^2$$

for all $g_j \in L^r(\mathbb{R}^{d_j})$. However, an elementary scaling argument shows that a necessary condition for (28) to hold is $r \geq (2m-2)'$, giving the desired contradiction.

We remark that the same scaling argument tells us that the exponent $m-1$ in Theorem 2.2 is optimal; this should not be too much of a surprise since the estimate (27) with linear mappings Σ_j and $r = (2m-2)'$ follows from the linear case of Theorem 2.2 combined with an application of the classical Hausdorff–Young inequality. Explicitly,

$$\begin{aligned} \|f_1 d\sigma_1 * \cdots * f_m d\sigma_m\|_{L^2(\mathbb{R}^d)} &= \left(\int_{\mathbb{R}^d} \prod_{j=1}^m |\widehat{g}_j|^2(\Sigma_j^* x) dx \right)^{\frac{1}{2}} \\ &\leq C \prod_{j=1}^m \|\widehat{g}_j\|_{L^{m-1}(\mathbb{R}^{d_j})}^{\frac{1}{2}} \\ &\leq C' \prod_{j=1}^m \|g_j\|_{L^{(2m-2)' }(\mathbb{R}^{d_j})} = C' \prod_{j=1}^m \|f_j\|_{L^{(2m-2)' } (d\sigma_j)}. \end{aligned}$$

Thus, it is clear that for $m \geq 3$, Theorem 1.1 has a stronger form with the inputs $(f_j)_{j=1}^m$ lying in certain Fourier–Lebesgue spaces, at least when the submanifolds are subspaces. This stems from an analogous improvement to Theorem 1.2, and it is reasonable to expect such an improvement in the nonlinear case. This line of investigation will be taken up in a subsequent paper.

4. THE GLOBAL CASE

In this section we prove Theorem 1.3.

Proof of Theorem 1.3. We let κ dominate the $C^{1,\beta}$ norm of F , and note that there exists $\varepsilon > 0$ such that

$$(29) \quad \left| \star \bigwedge_{j=1}^m \star Y_j(dF(\omega)) \right| \geq \varepsilon$$

for all ω in the unit sphere. Using the local result in Theorem 2.1, we obtain $0 < \delta < 1$ depending on at most β, κ and d , and a constant C depending on at most ε and d satisfying

$$(30) \quad \int_U \prod_{j=1}^m f_j(y_j) \delta(F(y)) \, dy \leq C \prod_{j=1}^m \|f_j\|_{L^{(m-1)'}}(\mathbb{R}^{d_j})$$

for all nonnegative $f_j \in L^{(m-1)'}}(\mathbb{R}^{d_j})$, $1 \leq j \leq m$, where $U = B(\omega, \delta)$. By (30), scaling and the homogeneity of F ,

$$\int_{\Gamma(\omega, \delta)} \prod_{j=1}^m f_j(y_j) \delta(F(y)) \, dy \leq C \sum_{k \in \mathbb{Z}} \prod_{j=1}^m \|f_j\|_{L^{(m-1)'}}(\lambda^k U_j).$$

Here, U_j is the subset of \mathbb{R}^{d_j} given by

$$U_j = \pi_j(U \cap F^{-1}(0)),$$

where $\lambda = 1 + c\delta$, for some appropriately chosen absolute constant $c > 0$, and $\pi_j : \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m} \rightarrow \mathbb{R}^{d_j}$ is the coordinate projection given by

$$\pi_j(y_1, \dots, y_m) = y_j.$$

Lemma 4.1. *For each ω in the unit sphere, there exists $j_1(\omega), j_2(\omega)$ which are distinct and such that, for $r = 1, 2$, the sets $\{\lambda^k U_{j_r(\omega)} : k \in \mathbb{Z}\}$ have bounded (independent of ω) overlap.*

We remark that if one drops from U_j the intersection with the zero set of F , the claimed bounded overlap property clearly ceases to hold.

Proof. We shall prove that there exists $\eta > 0$ such that for each ω in the unit sphere intersected with the zero set of F we have $|\pi_j(\omega)| \geq \eta$ for at least two $1 \leq j \leq m$.

Suppose ω is on the unit sphere with $F(\omega) = 0$, and suppose $|\pi_j(\omega)| \geq \eta$ only when $j = j_1$. Using the homogeneity of F , we have $\omega \in \ker dF(\omega)$ and so

$$\sum_{j=1}^m \pi_j(F^{(\ell)}) \cdot \pi_j(\omega) = 0$$

for each $1 \leq \ell \leq d$, where $F^{(\ell)}$ is the ℓ th row of $dF(\omega)$. Therefore

$$(31) \quad |\pi_{j_1}(F^{(\ell)}) \cdot \pi_{j_1}(\omega)| \leq C_{\kappa, m} \eta$$

for each $1 \leq \ell \leq d$. By (29) it follows that

$$\left\| \bigwedge_{k \in \mathcal{K}_{j_1}} dF(\omega)(e_k) \right\|_{\Lambda^{d_{j_1}}(\mathbb{R}^d)} \geq C_{\kappa, m} \varepsilon$$

which means that at least one of the components (with respect to the induced basis of $\Lambda^{d_{j_1}}(\mathbb{R}^d)$) of $\bigwedge_{k \in \mathcal{K}_{j_1}} dF(\omega)(e_k)$ is bounded below in magnitude by $C_{\kappa, m} \varepsilon$. This component is the determinant of a certain $d_{j_1} \times d_{j_1}$ matrix, and as a consequence, we may write

$$\pi_{j_1}(\omega) = \sum_{\ell=1}^d \alpha_\ell \pi_{j_1}(F^{(\ell)})$$

for some $\alpha_1, \dots, \alpha_d \in \mathbb{R}$ satisfying $|\alpha_\ell| \leq C_{\kappa, m} \varepsilon^{-1}$. Thus, by (31),

$$|\pi_{j_1}(\omega) \cdot \pi_{j_1}(\omega)| \leq C_{\varepsilon, \kappa, d, m} \eta$$

which means that $|\omega| = \sum_{j=1}^m |\pi_j(\omega)| \leq \frac{1}{2}$ for sufficiently small η , depending on $\varepsilon, \kappa, d, m$. From this contradiction we obtain the desired conclusion. \square

Now

$$\int_{\Gamma(\omega, \delta)} \prod_{j=1}^m f_j(y_j) \delta(F(y)) \, dy \leq C \prod_{\substack{j \neq j_1(\omega), \\ j_2(\omega)}} \|f_j\|_{L^{(m-1)' }(\mathbb{R}^{d_j})} \sum_{k \in \mathbb{Z}} \prod_{\substack{j=j_1(\omega), \\ j_2(\omega)}} \|f_j\|_{L^{(m-1)' }(\lambda^k U_j)}$$

and by Cauchy–Schwarz, the embedding $\ell^{(m-1)' }(\mathbb{Z}) \subset \ell^2(\mathbb{Z})$, and Lemma 4.1, it follows that

$$\int_{\Gamma(\omega, \delta)} \prod_{j=1}^m f_j(y_j) \delta(F(y)) \, dy \leq C \prod_{j=1}^m \|f_j\|_{L^{(m-1)' }(\mathbb{R}^{d_j})}.$$

By compactness, the fully global estimate over $\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_m}$ follows. \square

There is obviously some “slack” in the above argument where the embedding $\ell^{(m-1)' }(\mathbb{Z}) \subset \ell^2(\mathbb{Z})$ was used. This slackness was not present in the analogous argument in [1].

APPENDIX A. SOME EXTERIOR ALGEBRA

Lemma A.1. *If $\{v_1, \dots, v_r\} \subset \mathbb{R}^d$ is a linearly independent set and $\{n_1, \dots, n_{r'}\}$ is any orthonormal basis for the orthogonal complement of the space spanned by $\{v_1, \dots, v_r\}$, where $r + r' = d$, then*

$$\star \bigwedge_{k=1}^r v_k = \left\| \bigwedge_{k=1}^r v_k \right\|_{\Lambda^r(\mathbb{R}^d)} \bigwedge_{k=1}^{r'} n_k.$$

Proof. One notes that $\{v_1, \dots, v_r, n_1, \dots, n_{r'}\}$ is a basis for \mathbb{R}^d and so we have the induced basis \mathfrak{B} for $\Lambda^{r'}(\mathbb{R}^d)$. By definition of Hodge star we want to show that

$$(32) \quad \left\| \bigwedge_{k=1}^r v_k \right\|_{\Lambda^r(\mathbb{R}^d)} \left\langle \bigwedge_{k=1}^{r'} n_k, u \right\rangle_{\Lambda^{r'}(\mathbb{R}^d)} \sigma = u \wedge \bigwedge_{k=1}^r v_k$$

for each $u \in \Lambda^{n'}(\mathbb{R}^d)$, where σ is the unit basis element for $\Lambda^d(\mathbb{R}^d)$. It suffices to check (32) for $u \in \mathfrak{B}$ and we divide this task into the cases where $u = \bigwedge_{k=1}^{r'} n_k$ and $u \neq \bigwedge_{k=1}^{r'} n_k$. In the latter case is it easy to see that both sides of (32) are equal to zero.

If $u = \bigwedge_{k=1}^{r'} n_k$ then

$$\begin{aligned} \left\| \bigwedge_{k=1}^r v_k \right\|_{\Lambda^r(\mathbb{R}^d)} \left\langle \bigwedge_{k=1}^{n'} n_k, \bigwedge_{\ell=1}^{r'} n_\ell \right\rangle_{\Lambda^{r'}(\mathbb{R}^d)} &= \left\| \bigwedge_{k=1}^r v_k \right\|_{\Lambda^r(\mathbb{R}^d)} \det(n_k \cdot n_\ell)_{1 \leq k, \ell \leq r'} \\ &= \left\| \bigwedge_{k=1}^r v_k \right\|_{\Lambda^r(\mathbb{R}^d)} \end{aligned}$$

because $\{n_1, \dots, n_{r'}\}$ is orthonormal. Now

$$u \wedge \bigwedge_{k=1}^r v_k = \det(n_1 \cdots n_{r'} v_1 \cdots v_r) \sigma$$

and, using orthogonality considerations,

$$\begin{aligned} \det(n_1 \cdots n_{r'} v_1 \cdots v_r)^2 &= \det(n_1 \cdots n_{r'} v_1 \cdots v_r)^T (n_1 \cdots n_{r'} v_1 \cdots v_r) \\ &= \det(v_k \cdot v_\ell)_{1 \leq k, \ell \leq r} = \left\| \bigwedge_{k=1}^r v_k \right\|_{\Lambda^r(\mathbb{R}^d)}^2. \end{aligned}$$

Hence (32) holds in this case too, and this completes the proof of the lemma. \square

APPENDIX B. A QUANTITATIVE VERSION OF THE IMPLICIT FUNCTION THEOREM

We provide a quantitative version of the implicit function theorem for $C^{1,\beta}$ functions.

Theorem B.1. *Suppose $d, n \in \mathbb{N}$ and $\beta, \kappa > 0$ are given. Let $R_1, R_2 > 0$ be given by*

$$R_1 = \min \left\{ 1, R_2, \frac{1}{4d\kappa} R_2 \right\} \quad \text{and} \quad R_2 = \left(\frac{1}{16d\kappa} \right)^{\frac{1}{\beta}}.$$

If $F : \mathbb{R}^n \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is such that $\|F\|_{C^{1,\beta}} \leq \kappa$, $F(0, 0) = 0$ and

$$dF(0, 0)|_{\text{span}\{e_r : n+1 \leq r \leq n+d\}} = I_d,$$

then there exists a function $\eta : B(0, R_1) \subset \mathbb{R}^n \rightarrow B(0, R_2) \subset \mathbb{R}^d$ such that

$$F(x, \eta(x)) = 0 \quad \text{for each } x \text{ belonging to } B(0, R_1),$$

and a constant $\tilde{\kappa}$, depending on at most β, κ , and n , such that $\|\eta\|_{C^{1,\beta}} \leq \tilde{\kappa}$.

For $d = 1$, a proof can be found in [5]. The argument in that case extends easily to arbitrary d and we omit the details.

REFERENCES

1. J. A. Barceló, J. Bennett, N. Bez, S. Gutiérrez, *A note on global nonlinear Brascamp–Lieb inequalities*, preprint.
2. I. Bejenaru, S. Herr, *Convolutions of singular measures and applications to the Zakharov system*, J. Funct. Anal. **261** (2011), 478–506.
3. I. Bejenaru, S. Herr, J. Holmer, D. Tataru, *On the 2d Zakharov system with L^2 Schrödinger data*, Nonlinearity **22** (2009), 1063–1089.
4. I. Bejenaru, S. Herr, D. Tataru, *A convolution estimate for two-dimensional hypersurfaces*, Rev. Mat. Iberoamericana **26** (2010), 707–728.
5. J. Bennett, N. Bez, *Some nonlinear Brascamp–Lieb inequalities and applications to harmonic analysis*, J. Funct. Anal. **259** (2010), 2520–2556.
6. J. Bennett, A. Carbery, M. Christ, T. Tao, *The Brascamp–Lieb inequalities: finiteness, structure and extremals*, Geom. Funct. Anal. **17** (2007), 1343–1415.
7. J. Bennett, A. Carbery, J. Wright, *A nonlinear generalisation of the Loomis–Whitney inequality and applications*, Math. Res. Lett. **12** (2005), 443–457.
8. A. Moyua, A. Vargas and L. Vega, *Restriction theorems and maximal operators related to oscillatory integrals in \mathbb{R}^3* , Duke Math. J. **96** (1999), 547–574.
9. A. Ruiz, A. Vargas, *Partial recovery of a potential from backscattering data*, Comm. Partial Differential Equations **30** (2005), 67–96.
10. B. Stovall, *L^p improving multilinear Radon-like transforms*, preprint.
11. T. Tao, *Multilinear weighted convolution of L^2 -functions, and applications to nonlinear dispersive equations*, Amer. J. Math. **123** (2001), 839–908.
12. T. Tao, A. Vargas and L. Vega, *A bilinear approach to the restriction and Keakeya conjectures*, J. Amer. Math. Soc. **11** (1998), 967–1000.
13. T. Tao and J. Wright, *L^p improving bounds for averages along curves*, J. Amer. Math. Soc. **16** (2003), 605–638.

JONATHAN BENNETT, NEAL BEZ AND SUSANA GUTIÉRREZ, SCHOOL OF MATHEMATICS, THE WATSON BUILDING, UNIVERSITY OF BIRMINGHAM, EDGBASTON, BIRMINGHAM, B15 2TT, ENGLAND.

E-mail address: J.Bennett@bham.ac.uk

E-mail address: N.Bez@bham.ac.uk

E-mail address: S.Gutierrez@bham.ac.uk