

**EXISTENCE AND CONVERGENCE RESULTS FOR
INFINITE DIMENSIONAL NONLINEAR STOCHASTIC
EQUATIONS WITH MULTIPLICATIVE NOISE**

VIOREL BARBU, ZDZISŁAW BRZEŹNIAK, ERIKA HAUSENBLAS,
AND LUCIANO TUBARO

ABSTRACT. The solution X_n to a nonlinear stochastic differential equation of the form $dX_n(t) + A_n(t)X_n(t) dt - \frac{1}{2} \sum_{j=1}^N (B_j^n(t))^2 X_n(t) dt = \sum_{j=1}^N B_j^n(t)X_n(t)d\beta_j^n(t) + f_n(t) dt$, $X_n(0) = x$, where β_j^n is a regular approximation of a Brownian motion β_j , $B_j^n(t)$ is a family of linear continuous operators from V to H strongly convergent to $B_j(t)$, $A_n(t) \rightarrow A(t)$, $\{A_n(t)\}$ is a family of maximal monotone nonlinear operators of subgradient type from V to V' , is convergent to the solution to the stochastic differential equation $dX(t) + A(t)X(t) dt - \frac{1}{2} \sum_{j=1}^N B_j^2(t)X(t) dt = \sum_{j=1}^N B_j(t)X(t) d\beta_j(t) + f(t) dt$, $X(0) = x$. Here $V \subset H \cong H' \subset V'$ where V is a reflexive Banach space with dual V' and H is a Hilbert space. These results can be reformulated in terms of Stratonovich stochastic equation $dY(t) + A(t)Y(t) dt = \sum_{j=1}^N B_j(t)Y(t) \circ d\beta_j(t) + f(t) dt$.

[2000]60J60, 47D07, 15A36, 31C25

Stochastic differential equations, Brownian motion, progressively measurable, porous media equations.

1. INTRODUCTION

Consider the stochastic differential equation

$$\left\{ \begin{array}{l} dX(t) + A(t)X(t) dt - \frac{1}{2} \sum_{j=1}^N B_j^2(t)X(t) dt = \\ \sum_{j=1}^N B_j(t)X(t) d\beta_j(t) + f(t) dt, \quad t \in [0, T] \\ X(0) = x \end{array} \right. \quad (1.1)$$

where $A(t): V \rightarrow V'$ is a nonlinear monotone operator, $B_j(t) \in L(V, H)$, $\forall t \in [0, T]$ and β_j are independent Brownian motion in a probability space $\{\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P}\}$.

Equation (1.1) is of course equivalent with the Stratonovich stochastic differential equation

$$(1.1)' \quad dY(t) + A(t)Y(t) dt = \sum_{j=1}^N B_j(t)Y(t) \circ d\beta_j(t) + f(t) dt, \quad t \in [0, T].$$

Here V is a reflexive Banach space with dual V' such that $V \subset H \subset V'$ algebraically and topologically, where H (the *pivot* space) is a real Hilbert space. (The assumptions on $A(t)$, $B_j(t)$ will be made precise later on.)

We associate with (1.1) the random differential equation

$$\begin{cases} \frac{dy}{dt} + \Lambda(t)y = g(t), & t \in [0, T], \mathbb{P}\text{-a.s.} \\ y(0) = x, \end{cases} \quad (1.2)$$

where $g(t) = e^{\sum_{j=1}^N \beta_j(t) B_j^*(t)} f(t)$, $B_j^*(t)$ is the adjoint of $B_j(t)$ and $\Lambda(t)$ is the family of operators

$$\begin{aligned} \Lambda(t)y = & e^{-\sum_{j=1}^N \beta_j(t) B_j(t)} A(t) e^{\sum_{j=1}^N \beta_j(t) B_j(t)} y + \\ & + \sum_{j=1}^N \int_0^{\beta_j(t)} e^{-s B_j(t)} \dot{B}_j(t) e^{s B_j(t)} y ds, \quad \forall y \in V \end{aligned} \quad (1.3)$$

where \dot{B}_j is the derivative of $t \rightarrow B_j(t) \in L(V, H)$ and $(e^{s B_j(t)})_{s \in \mathbb{R}}$ is the C_0 -group generated by $B_j(t)$ on H and V .

It is well known (see e.g., [17, pag. 202], [14], [21]) that assuming that $B_j B_k(t) = B_k(t) B_j(t)$ for all j, k , at least formally equations (1.1) and (1.2) are equivalent via the transformation

$$X(t) = e^{\sum_{j=1}^N \beta_j(t) B_j(t)} y(t), \quad \mathbb{P}\text{-a.s.}, t \in [0, T], \quad (1.4)$$

and this is indeed the case if (1.2) has a strong, progressively measurable solution $y: [0, T] \times \Omega \rightarrow H$.

We consider also the family of approximating stochastic equations

$$\begin{cases} \frac{d}{dt} X_n + A_n(t) X_n = \sum_{j=1}^N B_j^n(t) X_n(t) \dot{\beta}_j^n(t) + f_n(t), & \mathbb{P}\text{-a.s.} \\ X_n(0) = x. \end{cases} \quad (1.5)$$

where $\{\beta_j^n\}$ is a sequence of smooth stochastic processes convergent to β_j , that is $\beta_j^n(t) \rightarrow \beta_j(t)$ uniformly on $[0, T]$, \mathbb{P} -a.s. and $A_n \rightarrow A$, $B_j^n \rightarrow B_j$, $f_n \rightarrow f$ as $n \rightarrow \infty$ in a sense to be made precise below.

Equation (1.5) is just an approximation of Stratonovich equation (1.1)' where β_j^n is a regularization of β_j . One must emphasize that $\{\beta_n\}$ might be adapted to a filtration different of $\{\mathcal{F}_t\}$.

Equation (1.5) reduces via (1.4), that is

$$(1.4)_n \quad X_n(t) = e^{\sum_{j=1}^N \beta_j^n(t) B_j^n(t)} y_n(t)$$

to a random differential equation of the form (1.2) that is

$$(1.2)_n \quad \begin{cases} \frac{dy_n}{dt} + \Lambda_n(t) y_n = g_n(t), & t \in [0, T], \mathbb{P}\text{-a.s.} \\ y_n(0) = x \end{cases}$$

where $g_n(t) = e^{\sum_{j=1}^N \beta_j^n(t) B_j^{n*}(t)} f_n(t)$ and Λ_n are given by

$$(1.3)_n \quad \Lambda_n(t) = e^{-\sum_{j=1}^N \beta_j^n(t) B_j^n(t)} A_n(t) e^{\sum_{j=1}^N \beta_j^n(t) B_j^n(t)} + \sum_{j=1}^N \int_0^{\beta_j^n(t)} e^{-s B_j^n(t)} \dot{B}_j^n(t) e^{s B_j^n(t)} y ds.$$

The main result (see Theorems 2.1, 2.2, 2.3 below) is that under suitable assumptions equations (1.2), (1.2)_n have unique solutions y and y_n which are progressively measurable processes and for $n \rightarrow \infty$, we have $y_n \rightarrow y$, $X_n \rightarrow X$ in a certain precise sense. In the linear case such a result was established by a different method in [14], (we refer also to [18], [21] to other results in this direction.) The variational method we use here allows to treat a general class of nonlinear equations (1.1) possibly multi-valued (On these lines see also [5, 6].)

The applications given in Sect. 4.1 refer to stochastic porous media equations and nonlinear stochastic diffusion equations of divergence type but of course the potential area of applications is much larger.

Notation. If Y is a Banach space we denote by $L^p(0, T; Y)$, $1 \leq p \leq \infty$ the space of all (equivalence classes of) Y -measurable functions $u: (0, T) \rightarrow Y$ with $\|u\|_Y \in L^p(0, T)$ (here $\|\cdot\|_Y$ is the norm of Y). Denote by $C([0, T]; Y)$ the space of all continuous Y -valued functions on $[0, T]$ and by $W^{1,p}([0, T]; Y)$ the infinite dimensional Sobolev space $\{y \in L^p(0, T; Y), \frac{dy}{dt} \in L^p(0, T; Y)\}$ where $\frac{d}{dt}$ is considered in sense of vectorial distributions. It is well known that $W^{1,p}([0, T]; Y)$ coincides with the space of absolutely continuous functions $y: [0, T] \rightarrow Y$, a.e. differentiable and with derivative $y'(t) = \frac{dy}{dt}(t)$ a.e. $t \in (0, T)$ and $\frac{dy}{dt} \in L^p(0, T; Y)$ (see e.g., [4].) If $p \in [1, \infty]$ is given we denote by p' the conjugate exponent, i.e., $\frac{1}{p} + \frac{1}{p'} = 1$.

2. THE MAIN RESULTS

We shall study here equation (1.2) under the following assumptions

(i) V is a separable real reflexive Banach space with the dual V' and H is a separable real Hilbert space such that $V \subset H \subset V'$ algebraically and topologically.

We denote by $|\cdot|$, $\|\cdot\|_V$ and $\|\cdot\|_{V'}$ the norms in H , V and V' , respectively and by $\langle \cdot, \cdot \rangle$ the duality pairing on $V \times V'$ which coincides with the scalar product (\cdot, \cdot) of H on $H \times H$.

(ii) $A(t)y = \partial\psi(t, y)$, a.e. $t \in (0, T)$, $\forall y \in V$, \mathbb{P} -a.s., where $\psi: (0, T) \times V \times \Omega \rightarrow \mathbb{R}$ is convex and lower-semicontinuous in y on V and measurable in t on $[0, T]$. There are $\alpha_i > 0$, $\gamma_i \in \mathbb{R}$, $i = 1, 2$, $1 < p_1 \leq p_2 < \infty$

$$\gamma_1 + \alpha_1 \|y\|_V^{p_1} \leq \psi(t, y) \leq \gamma_2 + \alpha_2 \|y\|_V^{p_2}, \quad \forall y \in V. \quad (2.1)$$

(iii) There are $C_1, C_2 \in \mathbb{R}^+$ such that

$$\psi(t, -y) \leq C_1 \psi(t, y) + C_2, \quad \forall y \in V, t \in (0, T). \quad (2.2)$$

(The constants C_i, γ_i, α_i are dependent on ω .)

(iv) For each $y \in V$ the stochastic process $\psi(t, y)$ is progressively measurable with respect to filtration $\{\mathcal{F}_t\}_{t \geq 0}$.

(v) $B_j(t)$ is a family of linear, closed and densely defined operators in H such that $B_j(t) = -B_j^*(t), \forall t \in [0, T]$, $B_j(t)$ generates a C_0 -group $(e^{s B_j(t)})_{s \in \mathbb{R}}$ on H and V . Moreover, $B_j \in C^1([0, T]; L(V, H))$, $B_j(t)B_k(t) = B_k(t)B_j(t)$ for all j, k .

(vi) $f: [0, T] \times \Omega \rightarrow V'$ is progressively measurable and $f \in L^{p_1}(0, T; V')$, \mathbb{P} -a.s.

We note that by (ii) $A(t, \omega): V \mapsto V'$ is, for all $t \in [0, T]$ and $\omega \in \Omega$, maximal monotone and surjective (see [4]) but in general multi-valued if ψ is not Gâteaux differentiable in y .

Theorem 2.1. *Let $y_0 \in H$. Then under assumptions (i) \sim (vi) there is for each $\omega \in \Omega$ a unique function $y = y(t, \omega)$ to equation (1.2) which satisfies*

$$y \in L^{p_1}(0, T; V) \cap C([0, T]; H) \cap W^{1, p_2}([0, T]; V'), \quad (2.3)$$

$$\begin{cases} \frac{dy}{dt}(t) + \Lambda(t)y(t) \ni g(t), & \text{a.e. } t \in (0, T), \\ y(0) = y_0. \end{cases} \quad (2.4)$$

Moreover, the process $y: [0, T] \times \Omega \rightarrow H$ is progressively measurable with respect to the filtration $\{\mathcal{F}_t\}_{t \geq 0}$.

If $A_n(t, \cdot)$ and $A_n^{-1}(t, \cdot)$ are multi-valued we mean by $A_n(t, y)$ and $A_n^{-1}(t, z)$ single valued sections.

By \xrightarrow{G} we denote the variational or Γ -convergence. This means that for each $y \in V$ and $\xi \in A(t)y$ there are y_n and $\xi_n \in A(t)y_n$ such that $y_n \rightarrow y$ strongly in V , $\xi_n \rightarrow \xi$ strongly in V' and similarly for $A_n^{-1}(t) \rightarrow A^{-1}(t)$. Assumption (2.5) implies and is equivalent to: $\psi_n(t, z) \rightarrow \psi(t, z)$, $\psi_n^*(t, \tilde{z}) \rightarrow \psi^*(t, \tilde{z})$ for all $z \in V$, $\tilde{z} \in V'$ and $t \in [0, T]$ where ψ^* is the conjugate of ψ (See e.g., [2].)

Theorem 2.2. *Assume that for each n , Λ_n, B_j^n and f_n satisfy (i) \sim (iv). Then for any $y_0 \in V$ there is a unique function $y_n = y_n(t, \omega)$ which satisfies (2.3) and equation (2.4) with Λ_n instead of Λ . Moreover, assume that for $n \rightarrow \infty$*

$$\begin{aligned} A_n(t) &\xrightarrow{G} A(t), & t \in [0, T] \\ A_n^{-1}(t) &\xrightarrow{G} A^{-1}(t), & t \in [0, T] \end{aligned} \quad (2.5)$$

$$f_n(\cdot, \omega) \rightarrow f(\cdot, \omega), \quad \text{strongly in } L^{p_2}(0, T; V'), \mathbb{P}\text{-a.s. in } \Omega. \quad (2.6)$$

$$B_j^n x \rightarrow B_j x, \quad \text{in } C^1([0, T]; H), \quad \forall x \in H. \quad (2.7)$$

Then for $n \rightarrow \infty$

$$y_n(\cdot, \omega) \rightarrow y(\cdot, \omega) \quad (2.8)$$

\mathbb{P} -a.s. weakly in $L^{p_1}(0, T; V)$, weakly-star in $L^\infty(0, T; H)$.

Assumption (2.5) implies and is equivalent to: $\psi_n(t, z) \rightarrow \psi(t, z)$, $\psi_n^*(t, \tilde{z}) \rightarrow \psi^*(t, \tilde{z})$ for all $z \in V$, $\tilde{z} \in V'$ and $t \in [0, T]$ where ψ^* is the conjugate of ψ (See e.g., [2].)

Coming back to equation (1.1) we say that the process $X : [0, T] \rightarrow H$ is a solution to (1.1), if it is progressively measurable with respect to the filtration $\{\mathcal{F}_t\}_{t \geq 0}$ induced by the Brownian motion,

$$X \in C([0, T], H) \cap L^{p_1}(0, T; V), \quad \mathbb{P}\text{-a.s.} \quad (2.9)$$

and

$$\begin{aligned} X(t) = & x - \int_0^t A(s)X(s) ds + \frac{1}{2} \sum_{j=1}^N \int_0^t B_j^2(s)X(s) ds \\ & + \sum_{j=1}^N \int_0^t B_j(s)X(s) d\beta_j(s) + \int_0^t f(s) ds, \quad \forall t \in [0, T], \quad \mathbb{P}\text{-a.s.} \end{aligned} \quad (2.10)$$

By Theorem 2.1 and Theorem 2.2 we find that

Theorem 2.3. *Under the assumptions of Theorem 2.1 there exist unique solutions X and X_n to (1.1) and (1.5) respectively given by*

$$X(t) = e^{\sum_{j=1}^N \beta_j(t) B_j(t)} y(t), \quad X_n(t) = e^{\sum_{j=1}^N \beta_j^n(t) B_j^n(t)} y_n(t), \quad (2.11)$$

where y and y_n are solutions to (1.2) and (1.2)_n. Moreover, we have

$$X, X_n \in L^{p_1}(0, T; V), \quad \mathbb{P}\text{-a.s.} \quad (2.12)$$

$X, X_n : [0, T] \rightarrow H$ are \mathbb{P} -a.s. continuous and

$$X_n \rightarrow X \quad \text{weakly in } L^{p_1}(0, T; V), \text{ weakly-star in } L^\infty(0, T; H), \quad \mathbb{P}\text{-a.s.} \quad (2.13)$$

The precise meaning of Theorem 2.2 and Theorem 2.3 is the structural stability of the Itô stochastic differential equation (1.1) and of its Stratonovich counterpart (1.1)'. As a matter of fact, as mentioned earlier, all these results can be reformulated in terms of Stratonovich equation (1.1)'.

One of the main consequences of Theorem 2.2 is that the Stratonovich stochastic equation is stable with respect to smooth approximations of the process $B(t)X d\beta(t)$. On the other hand, the general existence theory for infinite dimensional stochastic differential equations with linear multiplicative noise (see e.g., [16, 17]) is not applicable in present situation due to the fact that the noise coefficient $x \rightarrow B(t)x$ is not bounded on the basic space H . The approach we use here to treat equation (2.4) relies on Brezis-Ekeland variational principle [12], [13] which allows to reduce nonlinear evolution equations of potential type to convex optimization problems. (On these lines see also [5]-[7], [22], [24]).

The more general case of nonlinear monotone and demicontinuous operators $A(t) : V \rightarrow V'$ is ruled out from present theory and might expect however to extend the theory to this general case by using Fitzpatrick function formalism (see [22], [24]).

As in [14], see Corollary on p. 438, we can define a solution to problem (1.1)' for any **deterministic** continuous function $\beta : \mathbb{R}_+ \rightarrow \mathbb{R}^N$. The result from [14] was a generalisation of a analogous result from Sussmann's well

known paper [23], see also Doss [19]. We will formulate our result in the same fashion as in [14], i.e. the result contains implicitly a definition. Let us observe, that in this case, we prove the existence for any multidimensional continuous signal, thus a signal more general than a rough signal from the theory of rough paths. However, this is due to the assumption of the commutativity of the vector fields B_j , $j = 1, \dots, N$. In the result below, we need deterministic versions of assumptions **(ii)**, **(vi)**. Note that the assumption **(iv)** is now redundant.

(ii') $A(t)y = \partial\psi(t, y)$, a.e. $t \in (0, T)$, $\forall y \in V$, where $\psi: (0, T) \times V \rightarrow \mathbb{R}$ is convex and lower-semicontinuous in y on V and measurable in t on $[0, T]$. There exist $\alpha_i > 0$, $\gamma_i \in \mathbb{R}$, $i = 1, 2$, $1 < p_1 \leq p_2 < \infty$, such that

$$\gamma_1 + \alpha_1 \|y\|_V^{p_1} \leq \psi(t, y) \leq \gamma_2 + \alpha_2 \|y\|_V^{p_2}, \quad \forall y \in V. \quad (2.14)$$

(vi) $f \in L^{p_1}(0, T; V')$,

Theorem 2.4. *Assume that the assumptions **(i)**, **(iii)**, **(v)** as well as **(ii')** and **(vi')** are satisfied. Then for every $x \in V$ and every $\beta \in C([0, T]; \mathbb{R}^N)$, the problem*

$$\begin{cases} dX(t) + A(t)X(t) dt = \sum_{j=1}^N B_j(t)X(t) d\beta_j(t) + f(t) dt, & t \in [0, T] \\ X(0) = x \end{cases} \quad (2.15)$$

has a unique solution $X \in L^{p_1}(0, T; V) \cap C([0, T]; H)$ in the following sense.

- (i) For every $\beta \in C^1([0, T]; \mathbb{R}^N)$, the problem (2.15) has a unique solution $X \in L^{p_1}(0, T; V) \cap C([0, T]; H)$.
- (ii) If $\beta_n \in C^1([0, T]; \mathbb{R}^N)$ and $\beta_n \rightarrow \beta$ in $C([0, T]; \mathbb{R}^N)$ and $X_n \in L^{p_1}(0, T; V) \cap C([0, T]; H)$ is the (unique) solution to the problem (2.15) corresponding to β_n , then $X_n \rightarrow X$ in weakly in $L^{p_1}(0, T; V)$, weakly-star in $L^\infty(0, T; H)$, \mathbb{P} -a.s.

From Theorem 2.4 we infer that in the framework of Theorem 2.4 but with β being a Brownian motion, the problem (1.1) generates a random dynamical system on H . In an obvious way we have the following Corollary

Corollary 1. *Assume that the assumptions **(i)**, **(iii)**, **(v)** as well as **(ii')** and **(vi')** are satisfied. Assume that β in a standard canonical two-sided \mathbb{R}^N -valued Brownian motion on a filtered probability space $\{\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P}\}$, where $\Omega = \{\omega \in C(\mathbb{R}, \mathbb{R}^N) : \omega(0) = 0\}$. Let us define a map*

$$\vartheta : \mathbb{R} \times \Omega \ni (t, \omega) \mapsto \vartheta_t \omega = \omega(\cdot + t) - \omega(0) \in \Omega.$$

Then there exists a map

$$\varphi : \mathbb{R}^+ \times \Omega \times H \ni (t, \omega, x) \mapsto \varphi(t, \omega)x \in H$$

such that a pair (φ, ϑ) is a random dynamical system on H , see for instance Definition 2.1 in [15], and, for each $s \in \mathbb{R}$ and each $x \in H$, the process X , defined for $\omega \in \Omega$ and $t \geq s$ as

$$X(t, s; \omega, x) := \varphi(t - s; \vartheta_s \omega)x, \quad (2.16)$$

is a solution to problem (1.1) over the time interval $[s, \infty)$ with an initial data given at time s .

Remark 2.1. Theorem 2.4 provides a solution to problem (2.15) for every continuous path. Our main result provides a natural interpretation of this solution in the case when β is a Brownian motion. One can also provide a similar interpretation when β is a fractional, see for instance [20].

Corollary allows one to investigate the existence of random attractors, see [15].

These questions will be investigated in the future works.

3. PROOFS

Proof of Theorem 2.1

For simplicity we consider the case $N = 1$, that is $B_j = B$, $\beta_j = \beta$ for all j . We note first that though the operator $\Lambda(t) = \Lambda(t, \omega)$ is \mathbb{P} -a.e $\omega \in \Omega$, maximal monotone from V to V' the standard existence theory (see e.g., [4, pag. 177]) does not apply here. This is, however, due to the general growth condition (2.1) on $\psi(t, \cdot)$ and implicitly on $A(t)$ as well as due to the multivaluedness of $A(t)$.

So we shall use a direct approach which makes use of the variational structure of problem (1.2). (On these lines see also [5], [7, page 280]). Namely, we can write

$$\Lambda(t) = \partial\varphi(t, \cdot) + \Gamma(t), \quad \forall t \in [0, T]. \quad (3.1)$$

Here $\varphi: [0, T] \times V \rightarrow \mathbb{R}$ is given by

$$\varphi(t, y) = \psi(t, e^{\beta(t)B(t)}y) \quad (3.2)$$

and

$$\Gamma(t)y = \int_0^{\beta(t)} e^{-sB(t)} \dot{B}(t) e^{sB(t)} y ds, \quad \forall y \in H, t \in [0, T]$$

where $\dot{B} = \frac{d}{dt}B(t)$. We fix $\omega \in \Omega$.

By the conjugacy formulae (A.3) and (A.4) we now may equivalently write (1.2) (or (2.4)) as

$$\begin{cases} \varphi(t, y(t)) + \varphi^*(t, u(t)) = \langle y(t), u(t) \rangle, & \text{a.e. } t \in [0, T] \\ y'(t) + \Gamma(t)y(t) = -u(t) + g(t), & \text{a.e. } t \in [0, T] \end{cases} \quad (3.3)$$

while

$$\varphi(t, \bar{y}) + \varphi^*(t, \bar{u}) \geq \langle \bar{y}, \bar{u} \rangle$$

for all $\langle \bar{y}, \bar{u} \rangle \in L^{p_1}(0, T; V) \times L^{p_2}(0, T; V')$.

Thus following a well known idea due to Brezis and Ekeland (see e.g., [?, 12, 13]) we are lead to the optimization problem

$$\begin{aligned} \text{Min} \left\{ \int_0^T (\varphi(t, y(t)) + \varphi^*(t, u(t)) - \langle u(t), y(t) \rangle) dt : \right. \\ \left. y' + \Gamma(t)y = -u + g, \text{ a.e. } t \in [0, T]; y(0) = y_0, \right. \\ \left. y \in L^{p_1}(0, T; V), u \in L^{p_2}(0, T; V') \right\}. \quad (3.4) \end{aligned}$$

Equivalently

$$\begin{aligned} \text{Min} \left\{ \int_0^T (\varphi(t, y(t)) + \varphi^*(t, u(t)) - \langle g(t), y(t) \rangle) dt + \frac{1}{2} (|y(T)|^2 - |y_0|^2) : \right. \\ \left. y' + \Gamma(t)y = -u + g, \text{ a.e. } t \in [0, T]; y(0) = y_0, \right. \\ \left. y \in L^{p_1}(0, T; V), u \in L^{p'_2}(0, T; V') \right\}. \quad (3.5) \end{aligned}$$

Here we have used (for the moment, formally) the integration by parts formula

$$\begin{aligned} - \int_0^T \langle u(t), y(t) \rangle dt = \frac{1}{2} (|y(T)|^2 - |y_0|^2) \\ + \int_0^T \langle \Gamma(t)y(t), y(t) \rangle dt - \int_0^T \langle g(t), u(t) \rangle dt \end{aligned}$$

and hypothesis (v) which implies that $\langle \Gamma(t)y, y \rangle = 0$. Of course the equivalence of (3.4) and (3.5) is valid only if the above equality is true which is not always the case in absence of some additional properties of minimizer y to allow integration by parts in $\int_0^T \langle u(t), g(t) \rangle dt$. In the following we shall prove however that equation (3.5) has at least one solution and show consequently that it is also a solution to equation (2.4).

Lemma 3.1. *There is a solution $y^* \in L^{p_1}(0, T; V) \cap W^{1, p'_2}([0, T]; V')$ to equation (3.5).*

Proof. We note that by standard existence theory of linear evolution equations for each $u \in L^{p'_2}(0, T; V')$ there is a unique solution $y \in L^{p_2}(0, T; V) \cap W^{1, p'_2}([0, T]; V') \subset C([0, T]; H)$ to equation

$$y' + \Gamma(t)y = -u + g, \quad \text{a.e. } t \in [0, T], \quad y(0) = y_0.$$

By assumptions (2.1) and (2.2) we have

$$\tilde{\gamma}_1 + \tilde{\alpha}_1 \|y\|_V^{p_1} \leq \varphi(t, y) \leq \tilde{\gamma}_2 + \tilde{\alpha}_2 \|y\|_V^{p_2}, \quad \forall y \in V \quad (3.6)$$

and

$$\bar{\gamma}_1 + \bar{\alpha}_1 \|y\|_{V'}^{p'_2} \leq \varphi^*(t, y) \leq \bar{\gamma}_2 + \bar{\alpha}_2 \|y\|_{V'}^{p'_1}, \quad \forall y \in V' \quad (3.7)$$

where $\tilde{\gamma}_i, \bar{\gamma}_i \in \mathbb{R}$, $\frac{1}{p_i} + \frac{1}{p'_i} = 1$ and $\tilde{\alpha}_i, \bar{\alpha}_i > 0$, $i = 1, 2$. (Recall that $e^{sB(t)}$ is invertible).

Then the infimum m^* in (3.5) is $> -\infty$ and there are the sequences $\{y_j\} \subset L^{p_1}(0, T; V)$, $\{u_j\} \subset L^{p'_2}(0, T; V')$ such that for all y

$$m^* \leq \int_0^T (\langle \varphi(t, y_j) + \varphi^*(t, u_j) - \langle g, y_j \rangle) dt + \frac{1}{2} (|y_j(T)|^2 - |y_0|^2) \leq m^* + \frac{1}{j} \quad (3.8)$$

$$\begin{cases} y'_j + \Gamma(t)y_j = -u_j + g, & \text{in } [0, T] \\ y_j(0) = y_0. \end{cases} \quad (3.9)$$

Clearly $y_j \in W^{1, p'_2}([0, T]; V')$ and by assumption (2.1) and inequality (3.8) it follows that

$$\|y_j\|_{L^{p_1}(0, T; V)} + \|y'_j\|_{L^{p'_2}(0, T; V')} \leq C \quad (3.10)$$

because as easily seen by assumption (v), $|\Gamma(t)y|_H \leq C\|y\|_V, \forall y \in V$.
Hence on a subsequence, again denoted y_j , we have for $j \rightarrow \infty$

$$\begin{aligned} y_j &\rightarrow y^* \quad \text{weakly in } L^{p_1}(0, T; V) \\ u_j &\rightarrow u^* \quad \text{weakly in } L^{p'_2}(0, T; V') \\ y'_j &\rightarrow (y^*)' = g - u^* - \Gamma(t)y^* \quad \text{weakly in } L^{p'_2}(0, T; V'). \end{aligned} \quad (3.11)$$

Since the functions $y \mapsto \int_0^T \varphi(t, y) dt$ and $u \mapsto \int_0^T \varphi^*(t, u) dt$ are weakly lower-semicontinuous on $L^{p_1}(0, T; V)$ and $L^{p'_2}(0, T; V')$ respectively, letting j tend to infinity we obtain that

$$m^* = \int_0^T (\varphi(t, y^*) + \varphi^*(t, u^*) - \langle g, y^* \rangle) dt + \frac{1}{2}(|y^*(T)|^2 - |y_0|^2) \quad (3.12)$$

and

$$\begin{cases} (y^*)' + \Gamma(t)y^* = -u^* + g, & t \in [0, T] \\ y^*(0) = y_0. \end{cases} \quad (3.13)$$

Therefore (y^*, u^*) is a solution to optimization problem (3.5) as claimed. \square

Proof of Theorem 2.1 (continued). We shall show now that y^* given by Lemma 3.1 is a solution to (2.4). To this end we notice just that without any loss of generality we may assume that $y_0 = 0$. Indeed we can reduce the problem to this case by translating in problem 2.3 y in $y - y_0$.

We prove now that $m^* = 0$. For this purpose we invoke a standard duality result for infinite dimensional convex optimal control problems, essentially due to R.T. Rockafeller. Namely, one has (see [7, Thm. 4.6, pag. 287])

$$m^* + \min (3.5)' = 0 \quad (3.14)$$

where (3.5)' is the dual control problem

$$(3.5)' \quad \text{Min} \left\{ \int_0^T (\varphi(t, -p(t)) + \varphi^*(t, v(t)) + \langle g(t), p(t) \rangle) dt + \frac{1}{2}|p(T)|^2 : \right. \\ \left. p' + \Gamma(t)p = v + g, \quad t \in [0, T] \right\}.$$

If $(p^*, v^*) \in L^{p_1}(0, T; V) \times L^{p'_2}(0, T; V')$ is optimal in (3.5)', we have

$$\langle (p^*)', p^* \rangle \in L^1(0, T) \quad (3.15)$$

$$\int_0^T \langle (p^*)', p^* \rangle dt = \frac{1}{2}(|p^*(T)|^2 - |p^*(0)|^2). \quad (3.16)$$

Here is the argument. First, note that p' solves $p' + \Gamma(t)p = v + g$. We have by the identities (A.3) and (A.4) and the fact that $\langle \Gamma(t)p^*, p^* \rangle = 0$,

$$-\langle (p^*(t))', p^*(t) \rangle \leq \varphi^*(t, v^*(t)) + \varphi(t, -p^*(t)) - \langle g(t), p^*(t) \rangle, \quad \text{a.e. } t \in (0, T)$$

and

$$\langle (p^*(t))', p^*(t) \rangle \leq \varphi^*(t, v^*(t)) + \varphi(t, p^*(t)) + \langle g(t), p^*(t) \rangle, \quad \text{a.e. } t \in (0, T).$$

Since $\varphi(t, -p^*) \in L^1(0, T)$ and by assumption (2.2), $\varphi(t, p^*) \in L^1(0, T)$ too, we infer that (3.15) holds. Now since $p^* \in W^{1, p'_2}([0, T]; V') \cap L^{p_1}(0, T; V')$ we have

$$\frac{1}{2} \frac{d}{dt} |p^*(t)|^2 = \langle (p^*)'(t), p^*(t) \rangle, \quad \text{a.e. } t \in (0, T)$$

and by (3.15) we get (3.16) as claimed.

By (3.5)' and (3.16) we see that

$$\min (3.5)' = \int_0^T (\varphi(t, -p^*) + \varphi^*(t, v^*) + \langle v^*, p^* \rangle) dt + \frac{1}{2}|p^*(0)|^2 \geq 0.$$

Similarly, by (3.12), (3.13) and by

$$\frac{1}{2}(|y^*(T)|^2 - |y^*(0)|^2) = \int_0^T \langle (y^*)', y^* \rangle dt,$$

(the latter follows exactly as (3.16)) we see that

$$m^* = \int_0^T (\varphi(t, y^*) + \varphi^*(t, u^*) - \langle u^*, y^* \rangle) dt \geq 0.$$

Then by (3.14) we have that $m^* = 0$ and therefore again by (3.12) we have that

$$\begin{aligned} \varphi(t, y^*) + \varphi^*(t, u^*) &= \langle u^*, y^* \rangle, & \text{a.e. in } [0, T] \\ (y^*)' + \Gamma(t)y^* &= g - u^* & \text{a.e. in } [0, T] \end{aligned}$$

and therefore y^* is a solution to (2.4).

On the other hand, as seen earlier, we have

$$\frac{1}{2}(|y^*(t)|^2 - |y^*(s)|^2) = \int_s^t \langle (y^*)'(\tau), y^*(\tau) \rangle d\tau, \quad \forall 0 \leq s \leq t \leq T. \quad (3.17)$$

Hence $y^* \in C([0, T]; H)$. The uniqueness of y^* is immediate by (3.17). It remains to be proven that y^* is progressively measurable.

To this end we note as minimum in (3.5) the pair (y^*, u^*) is the solution to Euler-Lagrange system (see e.g., [7, page 263])

$$\begin{aligned} (y^*)' + \Gamma(t)y^* &= -u^* + g, & \text{a.e. } t \in (0, T), \quad \omega \in \Omega \\ q' - \Gamma'(t)q &= -g + A(t)y^*, & \text{a.e. } t \in (0, T) \\ u^*(t) &= A(t)q(t), & \text{a.e. } t \in (0, T), \quad \omega \in \Omega \\ y^*(0) &= y_0, \quad q(T) = -y^*(T). \end{aligned}$$

Since the latter two point boundary value problem has a unique solution (y^*, q) and is of dissipative (accretive) type it can be solved by iteration or more precisely by a gradient algorithm (see [7, page 252]). In particular, we have $y^* = \lim_{k \rightarrow \infty} y_k$, $q = \lim_{k \rightarrow \infty} q_k$ weakly in $L^{p_1}(0, T; V)$ and $u^* = \lim_{k \rightarrow \infty} u_k$ weakly in $L^{p_2}(0, T; V')$ where

$$\begin{aligned} y_k' + \Gamma(t)y_k &= -u_k + g & t \in [0, T], \\ q_k' - \Gamma'(t)q_k &= -g + A(t)y_k, & t \in [0, T], \\ u_{k+1} &= u_k - A^{-1}(t)u_k + q_k, & t \in [0, T], \\ y_k(0) &= y_0, \quad q_k(T) = 0, & k = 0, 1, 2, \dots \end{aligned}$$

Hence, if we start with a progressively measurable u_0 , we see that all u_k are progressively measurable and so are u^* and y^* . \square

Proof of Theorem 2.2. As in the previous case it follows that equation (1.2)_n has a unique solution $y_n \in W^{1,p_2'}([0, T]; V') \cap L^{p_1}(0, T; V)$ given by the minimization problem (3.5) where $g = g_n$ and $\varphi = \varphi_n$, $\varphi^* = \varphi_n^*$. Here φ_n is

given as in (3.2) where $\psi = \psi_n$ and β is replaced by β_n while φ_n^* is the conjugate of φ_n . We have, similarly, $\partial\psi_n = A_n$ and $\varphi_n(t, y) = \psi_n(t, e^{\beta_n(t)y})$

$$(y_n, u_n) = \arg \min \left\{ \int_0^T \varphi_n(t, y(t)) + \varphi_n^*(t, u(t)) - \langle g_n(t), u(t) \rangle dt + \frac{1}{2}(|y(T)|^2 - |y_0|^2); \quad y' + \Gamma_n y = -u + g_n, \quad y(0) = y_0 \right\}.$$

Here $\Gamma_n(t)y = \int_0^{\beta_n(t)} e^{-sB_n(t)} \dot{B}_n(t) e^{sB_n(t)} y ds$.

We see that

$$\|y_n\|_{L^\infty(0,T;H)} + \|y_n\|_{L^{p_1}(0,T;V)} + \left\| \frac{dy_n}{dt} \right\|_{L^{p'_2}(0,T;V')} + |y_n(T)| \leq C,$$

and this implies that on a subsequence, again denoted $\{n\}$, we have

$$\left\{ \begin{array}{l} u_n \rightarrow \tilde{u} \quad \text{weakly in } L^{p'_2}(0, T; V') \\ y_n \rightarrow \tilde{y} \quad \text{weakly in } L^{p_1}(0, T; V) \\ y_n \rightarrow \tilde{y} \quad \text{weakly-star in } L^\infty(0, T; H) \\ \frac{dy_n}{dt} \rightarrow \frac{d\tilde{y}}{dt} \quad \text{weakly in } L^{p'_2}(0, T; V') \\ y_n(T) \rightarrow \tilde{y}(T) \quad \text{weakly in } H. \end{array} \right. \quad (3.19)$$

By (2.5), (2.6) we see that $\tilde{y}' + \Gamma\tilde{y} = -\tilde{u} + g$.

Moreover, we have

$$\begin{aligned} & \int_0^T (\varphi_n(t, y_n(t)) + \varphi_n^*(t, u_n(t)) - \langle g_n(t), y_n(t) \rangle) dt + \frac{1}{2}|y_n(T)|^2 \\ & \leq \int_0^T (\varphi_n(t, y^*(t)) + \varphi_n^*(t, u^*(t)) - \langle g_n(t), u^*(t) \rangle) dt + \frac{1}{2}|y^*(T)|^2. \end{aligned}$$

where (y^*, u^*) is the solution to (3.5). Now by assumptions (2.5) and (2.6) we have

$$\begin{aligned} \varphi_n(t, y^*(t)) & \rightarrow \varphi(t, y^*(t)) \\ \varphi_n^*(t, u_n^*(t)) & \rightarrow \varphi(t, u^*(t)) \quad \forall t \in [0, T], \\ g_n(t) & \rightarrow g(t) \end{aligned}$$

and this yields

$$\limsup_{n \rightarrow \infty} \int_0^T (\varphi_n(t, y_n) + \varphi_n^*(t, u_n) - \langle g_n, y_n \rangle) dt + \frac{1}{2}|y_n^*(T)|^2 - \frac{1}{2}|y_0|^2 = 0. \quad (3.20)$$

In order to pass to limit in (3.20) we shall use (3.19) and the convergence of $\{\varphi_n\}$ and $\{\varphi_n^*\}$ mentioned above. We set $\tilde{z}(t) = e^{\beta(t)B(t)}\tilde{y}(t)$, $z_n(t) = e^{\beta_n(t)B(t)}y_n(t)$. We have

$$\psi_n(t, \tilde{z}(t)) \leq \psi_n(t, z_n(t)) + \langle A_n(t, \tilde{z}(t)), \tilde{z}(t) - z_n(t) \rangle,$$

and since $\partial\psi_n^* = A_n^{-1}$ we have also that

$$\psi_n^*(t, \theta(t)) \leq \psi_n^*(t, \theta_n(t)) + \langle A_n^{-1}(t)(t, \theta(t)), \theta(t) - \theta_n(t) \rangle, \quad \text{a.e. } t \in (0, T).$$

where $\theta(t) = g(t) - \tilde{y}'(t)$, $\theta_n(t) = g_n(t) - y_n'(t)$.

Then by assumption (2.5) and equation (3.20) we have that

$$\limsup_{n \rightarrow \infty} \int_0^T (\varphi_n(t, \tilde{y}(t)) + \varphi_n^*(t, g(t) - \tilde{y}'(t)) - \langle g(t), \tilde{y}'(t) \rangle) dt + \frac{1}{2} |\tilde{y}^*(T)|^2 - \frac{1}{2} |y_0|^2 = 0,$$

and so, since as seen earlier (2.5) implies that $\varphi_n(t, z) \rightarrow \varphi(t, z)$, $\forall z \in V$, $\varphi_n^*(t, z^*) \rightarrow \varphi^*(t, z^*)$, $\forall z^* \in V'$, by the Fatou lemma, we have

$$\int_0^T (\varphi(t, \tilde{y}) + \varphi^*(t, \tilde{u}) - \langle g, \tilde{y} \rangle) dt + \frac{1}{2} |\tilde{y}^*(T)|^2 - \frac{1}{2} |y_0|^2 \leq 0,$$

which implies as in the previous case that \tilde{y} is a solution to (3.5) and therefore to (2.4) as claimed. \square

4. EXAMPLES

The specific examples to be presented below refer to nonlinear parabolic stochastic equations which can be written in the abstract form (1.1) where $A(t)$ are subpotential monotone and continuous operators from a separable Banach space V to its dual V' .

We briefly present below a few stochastic PDE to which the above theorems apply. We use here the standard notations for spaces of integrable functions and Sobolev spaces $W_0^{1,p}(\mathcal{O})$, $W^{-1,p'}(\mathcal{O}) = (W_0^{1,p}(\mathcal{O}))'$, $H^k(\mathcal{O})$, $k = 1, 2$ on open domains $\mathcal{O} \subset \mathbb{R}^d$.

4.1. Nonlinear stochastic diffusion equations. Consider the stochastic equation

$$\begin{cases} dX_t - \operatorname{div}_\xi a(t, \nabla_\xi X_t) dt - \frac{1}{2} b(t, \xi) \cdot \nabla_\xi (b(t, \xi) \cdot \nabla_\xi X_t) dt \\ \quad \quad \quad = b(t, \xi) \cdot \nabla_\xi X_t d\beta(t), \quad \text{in } (0, T) \times \mathcal{O} \\ X_0 = x \quad \text{in } \mathcal{O} \\ X_t = 0 \quad \text{on } (0, T) \times \partial\mathcal{O} \end{cases} \quad (4.1)$$

Here $a: (0, T) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a map of gradient type, i.e.,

$$a(t, y) = \partial j(t, y), \quad \forall y \in \mathbb{R}^d, t \in (0, T)$$

where $j: (0, T) \times \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}$ is convex in y , progressively measurable in $(t, \omega) \in [0, T) \times \Omega$ and

$$\gamma_1 + \alpha_1 |y|^{p_1} \leq j(t, y) \leq \gamma_2 + \alpha_2 |y|^{p_2}, \quad \forall y \in \mathbb{R}^d, \omega \in \Omega, t \in (0, T) \quad (4.2)$$

$$j(t, -y) \leq c_1 j(t, y) + c_2, \quad \forall y \in \mathbb{R}^d, t \in (0, T). \quad (4.3)$$

It should be emphasized that the mapping $r \rightarrow a(t, r)$ might be multivalued and discontinuous. As a matter of fact if $a(t, \cdot)$ is discontinuous at $r = r_j$, but left and right continuous (as happens by monotonicity) it is replaced by a multivalued maximal monotone mapping \tilde{a} obtained by filling the jumps at $r = r_j$.

Equation (4.1) is of the form (1.1) where $H = L^2(\mathcal{O})$, $V = W_0^{1,p_1}(\mathcal{O})$, $A(t) = \partial\psi(t, \cdot)$, $2 \leq p_1 \leq p_2 < \infty$,

$$\psi(t, u) = \int_{\mathcal{O}} j(t, \nabla u) d\xi, \quad \forall u \in W_0^{1,p_1}(\mathcal{O})$$

and

$$B(t)u = b(t, \xi) \cdot \nabla_{\xi} u = \operatorname{div}_{\xi}(b(t, \xi)u), \quad \forall u \in W_0^{1,p_1}(\mathcal{O}). \quad (4.4)$$

As regards the function $b(t, r): [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ we assume that

$$b(t, \cdot), \quad \frac{\partial b}{\partial r}(t, \cdot) \in (C([0, T]; \bar{\mathcal{O}}))^d \quad (4.5)$$

$$r \rightarrow b(t, \cdot) + \alpha r \text{ is monotone for some } \alpha \geq 0, \quad (4.6)$$

$$\operatorname{div}_{\xi} b(t, \xi) = 0, \quad b(t, \xi) \cdot \nu(\xi) = 0 \quad \forall \xi \in \partial \mathcal{O} \quad (4.7)$$

where ν is the normal to $\partial \mathcal{O}$. (The boundary $\partial \mathcal{O}$ is assumed to be of class C^1 .)

Here $\operatorname{div}_{\xi} b$ is taken in the sense of distributions on \mathcal{O} .

Then (4.4) defines a linear continuous operator $B(t)$ from V to $H = L^2(\mathcal{O})$ which as early seen is densely defined skew-symmetric, that is $-B(t) \subset B^*(t) \forall t \in [0, T]$. Moreover, $B(t)$ is m -dissipative in $L^2(\mathcal{O})$, that is the range of $u \rightarrow u - B(t)u$ is all of $L^2(\mathcal{O})$. Indeed for each $f \in L^2(\mathcal{O})$ the equation $u - B(t)u = f$ has the solution

$$u(\xi) = \int_0^{\infty} e^{-s} f(Z(s, \xi)) ds, \quad \forall \xi \in \mathcal{O},$$

where $s \rightarrow Z(s, \xi)$ is the differential flow defined by equation

$$\frac{dZ}{ds} = b(t, Z), \quad s \geq 0, Z(0) = \xi. \quad (4.8)$$

(By assumptions (4.6), (4.7), it follows that $t \rightarrow Z(t, \xi)$ is well defined on $[0, \infty)$.)

Hence, for each $t \in [0, T]$, $B(t)$ generates a C_0 -group $(e^{sB(t)})_{s \in \mathbb{R}}$ on $L^2(\mathcal{O})$ which is given by

$$(e^{B(t)s} f)(\xi) = f(Z(s, \xi)), \quad \forall f \in L^2(\mathcal{O}), s \in \mathbb{R}.$$

It is also clear that $e^{B(t)s} V \subset V$ for all $s \geq 0$.

Remark 4.1. Assumptions (4.5)–(4.7) can be weakened to discontinuous multivalued mappings $\xi \rightarrow b(t, \xi)$ satisfying (4.6), (4.7) and such that the solution $Z = Z(s, \xi; t)$ to the characteristic system (4.8) is differentiable in t . The details are omitted.

The corresponding random differential equation (1.2) has the form

$$\begin{cases} \frac{\partial y}{\partial t} - e^{\beta(t)B(t)} \operatorname{div}_{\xi}(a(t, \nabla_{\xi} e^{-\beta(t)B(t)} y) \\ \quad + \int_0^{\beta(t)} e^{sB(t)} \dot{B}(t) e^{-sB(t)} y ds = 0, & \text{in } (0, T) \times \mathcal{O}, \\ y(0, \xi) = x(\xi) & \text{in } \mathcal{O}, \\ y(t, \xi) = 0 & \text{on } (0, T) \times \partial \mathcal{O}. \end{cases} \quad (4.9)$$

Then by theorem 2.1 we have

Theorem 4.1. *There exists a solution X to (4.1) such that \mathbb{P} -a.s. $X \in L^{p_1}(0, T; W_0^{1,p_1}(\mathcal{O})) \cap L^{p_2}(0, T; W^{-1,p_2}(\mathcal{O})) \cap L^{\infty}(0, T; L^2(\mathcal{O}))$.*

We also note that in line with Theorem 2.2 if X_n , $n \in \mathbb{N}$, are solutions to equations

$$\begin{cases} dX_t^n - \operatorname{div}_\xi a^n(t, \nabla_\xi X_t^n) dt - \frac{1}{2} b_n(t, \xi) \cdot \nabla (b_n(t, \xi) X_t^n) dt = \\ \quad = b_n(t, \xi) \cdot \nabla_\xi X_t^n d\beta^n(t), \quad (t, \xi) \text{ in } (0, T) \times \mathcal{O}, \\ X_0^n = x \text{ in } \mathcal{O}, \\ X_t^n = \text{ on } (0, T) \times \partial\mathcal{O}, \end{cases} \quad (4.10)$$

where $b_n \rightarrow b$ uniformly on $[0, T] \times \mathcal{O}$ and $a_n(t, y) \rightarrow a(t, y)$, $a_n^{-1}(t, y) \rightarrow a^{-1}(t, y)$, $\beta_n(t) \rightarrow \beta(t)$ for all $y \in \mathbb{R}^d$, $t \in [0, T)$, then $X^n \rightarrow X$ weakly in $L^{p_1}(0, T; W_0^{1, p_1}(\mathcal{O}))$. Standard examples refer to structural stability, PDEs as well as to homogenization type results for equation (4.1). In latter case $a_n(t, z) = a(t, nz)$ where $a(t, \cdot)$ is periodic (see e.g., [2].)

Equation (4.1) is relevant in the mathematical description of nonlinear diffusion processes perturbed by a Brownian distribution with coefficient transport term $b(t, \xi) \cdot \nabla_\xi X$.

The assumption $p_1 \geq 2$ was taken here for technical reason required by the functional framework we work in and this excludes several relevant examples. For instance, the limit case $p_1 = 1$ which corresponds to the nonlinear diffusion function $a(t, y) = \rho \frac{y}{|y|^d}$, $\rho > 0$, which is relevant in material science and image restoring techniques (see e.g. [3, 4]) is beyond our approach and requires a specific treatment (see also [9] for the treatment of a similar problem with additive and continuous noise.)

In 2-D the appropriate functional setting to treat such a problem is $V = BV(\mathcal{O})$ the space of functions with bounded variation on \mathcal{O} with the norm $\varphi(y)$ and $H = L^2(\mathcal{O})$. Here $\varphi(y) = \|Dy\| + \int_{\partial\mathcal{O}} |\gamma_0(y)| d\mathcal{H}$, $y \in V$, $\|Dy\|$ is the variation of $y \in V$, $\gamma_0(y)$ is the trace on $\partial\mathcal{O}$ and $d\mathcal{H}$ is the Hausdorff measure on $\partial\mathcal{O}$. We recall that the norm φ is just the lower semicontinuous closure of the norm of Sobolev space $W_0^{1,1}(\mathcal{O})$ (see e.g., [1, pag. 438].) Then the approach developed in section 3 can be adapted to present situation though V is not reflexive. We expect to treat this limit case in a forthcoming work. (On these lines see also [6].)

4.2. Linear diffusion equations with nonlinear Neumann boundary conditions. Consider the equation

$$\begin{cases} dX_t - \Delta X_t dt - \frac{1}{2} b(t, \xi) \cdot \nabla_\xi (b(t, \xi) \cdot \nabla_\xi X_t) dt = \\ \quad b(t, \xi) \cdot \nabla_\xi X_t d\beta(t) \quad \text{in } [0, T] \times \mathcal{O} \\ \frac{\partial}{\partial \nu} X_t + \zeta(t, X_t) \ni 0 \quad \text{on } [0, T] \times \partial\mathcal{O} \\ X_0 = x \quad \text{in } \mathcal{O} \end{cases} \quad (4.11)$$

where $\zeta(t, r) = \partial j_0(t, r)$, $\forall t \in (0, T)$, $r \in \mathbb{R}$ and $j_0(t, \cdot)$ is a lower semicontinuous convex function on \mathbb{R} such that

$$\gamma_1 + \alpha_1 |y|^2 \leq j_0(t, y) \leq \gamma_2 + \alpha_2 |y|^2, \quad \forall y \in \mathbb{R}, t \in (0, T)$$

and $\alpha_i > 0$, $\gamma_i \in \mathbb{R}$, $i = 1, 2$.

Assume also that (4.3) holds and that $b = b(t, \cdot)$ satisfies conditions (4.5)–(4.7).

Then we may apply Theorems 2.1, 2.2, and 2.3, where $V = H^1(\mathcal{O})$, $H = L^2(\mathcal{O})$ and

$$\psi(t, y) = \frac{1}{2} \int_{\mathcal{O}} |\nabla y|^2 d\xi + \int_{\partial\mathcal{O}} j(t, y) d\xi, \quad \forall y \in V.$$

It follows so the existence of a solution $X \in L^2(0, T; V) \cap W^{1,2}([0, T]; V')$ to (4.11) and also the structural stability of (4.11) with respect to b . Problems of this type arise in thermostat central. In this case

$$\zeta(t, y) = \begin{cases} (\alpha_1(t)H(y) + \alpha_2(t)H(-y))y & \text{if } y \neq 0 \\ [-\alpha_2(t), \alpha_1(t)] & \text{if } y = 0 \end{cases}$$

where $\alpha_i > 0$, $\forall t \in [0, T]$ and H is the Heaviside function.

4.3. Nonlinear stochastic porous media equation. Consider the equation

$$\begin{cases} dX_t - \Delta_\xi \phi(t, X_t) dt - \frac{1}{2} b(t, \xi) \cdot \nabla_\xi (-\Delta)^{-1} (b(t, \xi) \cdot \nabla_\xi ((-\Delta)^{-1} X_t) dt \\ \qquad \qquad \qquad = b(t, \xi) \cdot \nabla_\xi (-\Delta)^{-1} X_t d\beta(t), & \text{in } (0, T) \times \mathcal{O} \\ X_0 = x & \text{in } \mathcal{O} \\ X_t = 0 & \text{on } (0, \infty) \times \partial\mathcal{O} \end{cases} \quad (4.12)$$

Here $\mathcal{O} \subset \mathbb{R}^d$, $d = 1, 2, 3$ is a bounded open domain and $(-\Delta)^{-1}$ is the inverse of the operator $A_0 = -\Delta$, $D(A_0) = H_0^1(\mathcal{O}) \cap H^2(\mathcal{O})$. The function $\phi: (0, T) \times \mathbb{R}^d \rightarrow \mathbb{R}$ is assumed to satisfy the following conditions

(k) $\phi = \phi(t, r)$ is monotonically decreasing in r , measurable in t and its potential

$$j(t, r) = \int_0^r \phi(t, \tau) d\tau, \quad t \in (0, T)$$

satisfies the growth conditions

$$\gamma_1 + \alpha_1 |r|^{p_1} \leq j(t, r) \leq \gamma_2 + \alpha_2 |r|^{p_2}, \quad \forall r \in \mathbb{R}, \omega \in \Omega, t \in [0, T] \quad (4.13)$$

$$j(t, -r) \leq c_1 j(t, r) + c_2, \quad \forall r \in \mathbb{R}, t \in (0, T) \quad (4.14)$$

where $\frac{6}{5} \leq p_1 \leq p_2 < \infty$ if $d = 3$, $1 < p_1 \leq p_2 < \infty$ if $d = 1, 2$.

Then equation (4.12) can be written as (1.1), where $H = H^{-1}(\mathcal{O})$, $V = L^{p_1}(\mathcal{O})$ and $A(t) = \partial\psi(t, \cdot)$ where $\psi(t, \cdot): H \rightarrow \bar{\mathbb{R}}$ is defined by

$$\psi(t, y) = \begin{cases} \int_{\mathcal{O}} j(t, y) d\xi & \text{if } y \in H^{-1}(\mathcal{O}), j(t, y) \in L^1(\mathcal{O}) \\ +\infty & \text{otherwise,} \end{cases}$$

and $B(t)$, $t \in \mathbb{R}^+$ is defined by

$$B(t)u = b(t, \xi) \cdot \nabla((-\Delta)^{-1}u), \quad u \in V. \quad (4.15)$$

The space V' is in this case the dual of $V = L^{p_1}(\mathcal{O})$ with $H^{-1}(\mathcal{O})$ as *pivot* space. By the Sobolev embedding theorem it is easily seen that since $p_1 \geq \frac{6}{5}$ we have $V \subset H^{-1}(\mathcal{O})$. The scalar product on H is defined by

$$\langle u, v \rangle_{H^{-1}(\mathcal{O})} = u(z), \quad z = (-\Delta)^{-1}v.$$

It is well known that $A(t)X = -\Delta_\xi \phi(t, X)$ is indeed the subdifferential of $\psi(t, \cdot)$ in $H^{-1}(\mathcal{O})$ (see e.g., [4, pag. 68]).

As regards $b: [0, T] \times \bar{\mathcal{O}} \rightarrow \mathbb{R}^d$ we assume that conditions (4.5)–(4.7) hold. We note that for each $t \in [0, T]$, $B(t) \in L(V, H^{-1}(\mathcal{O}))$ is densely defined and skew-symmetric on $H^{-1}(\mathcal{O}) = H$. Indeed we have

$$\begin{aligned} \langle B(t)u, u \rangle &= \int_{\mathcal{O}} \operatorname{div}(b(t, \xi)(-\Delta)^{-1}u) \cdot (-\Delta)^{-1}u \, d\xi = \\ &= \frac{1}{2} \int_{\mathcal{O}} b(t, \xi) \cdot \nabla |(-\Delta)^{-1}u(\xi)|^2 \, d\xi = 0, \end{aligned}$$

because $\operatorname{div}_{\xi} b = 0$ and $b(t, \xi) \cdot \nu(\xi) = 0$ on $\partial\mathcal{O}$.

Moreover, for each $t \in [0, T]$, $B(t)$ is m -dissipative on $H^{-1}(\mathcal{O})$. Indeed for each $f \in H^{-1}(\mathcal{O})$, the equation $u - B(t)u = f$ can be equivalently written as $v = (-\Delta)^{-1}u$, where

$$\begin{aligned} -\Delta v - b(t, \cdot) \cdot \nabla v &= f \quad \text{in } \mathcal{O}, \\ v &= 0 \quad \text{on } \partial\mathcal{O}. \end{aligned}$$

By Lax-Milgram lemma the latter has a unique solution $v \in H_0^1(\mathcal{O})$ and therefore $u \in H^{-1}(\mathcal{O})$ as claimed. Moreover, if $f \in L^{p_1}(\mathcal{O})$ and $\partial\mathcal{O}$ is of class C^2 then by the Agmon-Douglis-Nirenberg theorem $v \in W^{2, p_1}(\mathcal{O}) \cap W_0^{1, p_1}(\mathcal{O})$ and so $u \in V$.

Hence, $B(t)$ generates a C_0 -group $(e^{sB(t)})_{s \in \mathbb{R}}$ on $H = H^{-1}(\mathcal{O})$ which leaves $V = L^{p_1}(\mathcal{O})$ invariant.

Then we may apply Theorem 2.1 as well as the approximation Theorem 2.2 to the present situation. We obtain

Theorem 4.2. *There is a unique solution X to (4.12) such that \mathbb{P} -a.s. $X \in L^{p_1}(0, T; L^{p_1}(\mathcal{O})) \cap L^{\infty}(0, T; H^{-1}(\mathcal{O}))$. Moreover, the solution X is a limit of approximating solutions when the Brownian motion β is approximated by a sequence of smooth processes.*

Moreover, if $\phi_n \rightarrow \phi$ and $\phi_n^* \rightarrow \phi^*$, $b_n \rightarrow b$ we find by Theorem 2.2 that the corresponding solutions X_n to (4.12) are convergent to solution X to (4.11). The details are omitted.

Existence for stochastic porous media equation of the form

$$\begin{cases} dX_t - \Delta_{\xi} \phi(X_t) \, dt = \sigma(X_t) \, dW(t) & \text{in } (0, T) \times \mathcal{O} \\ X_0 = x & \text{in } \mathcal{O} \\ X_t = 0 & \text{on } (0, T) \times \partial\mathcal{O}, \end{cases}$$

when W_t is a Wiener process of the form

$$W(t, \xi) = \sum_{k=1}^{\infty} \mu_k e_k(\xi) \beta_k(t)$$

with $\sum_{k=1}^{\infty} \mu_k^2 \lambda_k^2 < \infty$, $\Delta e_k = -\lambda_k e_k$ in \mathcal{O} , $e_k \in H_0^1(\mathcal{O})$, and $\sigma = \sigma(x)$ is a linear continuous operator, were studied in [8, 9]. Note that in this case the noise term can also be written in our form with commuting and, contrary to our paper, bounded operators B_j . Here the multiplicative term $\sigma(X_t) = b \cdot \nabla (-\Delta)^{-1} X_t$ is however discontinuous on the space $H^{-1}(\mathcal{O})$ and so Theorem 4.2 is from this point of view different and in this sense more general.

Equation (4.12) models diffusion processes and the motion of fluid flows in porous media. The case considered here ($p_1 > 1$) is that of slow diffusion.

Remark 4.2. Theorem 2.4 and Remark 2.1 are also valid in the current setup.

APPENDIX A. CONVEX FUNCTIONS

We summarize in this paragraph some facts about convex functions, which we have used in our paper.

Given a convex and lower-semi continuous function $\phi : Y \rightarrow \bar{\mathbb{R}} = (-\infty, \infty]$ we denote by $\partial\phi : Y \rightarrow Y'$ (the dual space) the *subdifferential* of ϕ , i.e.

$$\partial\phi(y) := \{z \in Y' : \phi(y) - \phi(u) \leq \langle y - u, z \rangle, \forall u \in Y\}. \quad (\text{A.1})$$

(Here $\langle \cdot, \cdot \rangle$ is the duality pairing between Y and Y'). The function $\phi^* : Y' \rightarrow Y$ defined by

$$\phi^*(z) = \sup \{\langle y, z \rangle - \phi(y) : y \in Y\}, \quad (\text{A.2})$$

is called the conjugate of ϕ and as ϕ . Similarly to it is convex lower semi continuous function on Y' . Also we notice the following key conjugacy formulae (see e.g. [7, p. 89]). If $y \in Y$, and $z \in Y'$

$$\phi(y) + \phi^*(z) \geq \langle y, z \rangle \quad (\text{A.3})$$

$$\phi(y) + \phi^*(z) = \langle y, z \rangle \quad \text{iff } z \in \partial\phi(y), \quad (\text{A.4})$$

A vector x^* is said to be a subgradient of a convex function ϕ at a point x if

$$\phi(z) \geq \phi(x) + \langle x^*, z - x \rangle. \quad (\text{A.5})$$

Moreover, straightforward calculations give

$$\phi^*(x^*) = \phi_y^*(x^*) - \langle y, x^* \rangle, \quad (\text{A.6})$$

whenever $\phi(x) = \phi_y(x + y)$.

Acknowledgements. The work of V. Barbu was supported by the grant of the Romanian National Authority for Scientific Research 1ERC/02.07.2012 and by CIRM (Fondazione Bruno Kessler). The work of E. Hausenblas was supported by the Austrian Science Fund (FWF): P20705. Moreover, the authors would like to thank the Newton Institute, where part of this work was done during the special semester on ‘‘Stochastic Partial Differential Equations’’.

REFERENCES

- [1] H. Attouch, VARIATIONAL CONVERGENCE FOR FUNCTIONS AND OPERATORS, *Applicable Mathematics Series*. Pitman (Advanced Publishing Program), Boston, 1984.
- [2] H. Attouch, G. Buttazzo, G. Michaille, VARIATIONAL ANALYSIS IN SOBOLEV AND BV SPACES: APPLICATIONS TO PDES AND OPTIMIZATION. *MPS/SIAM Series on Optimization*, Philadelphia, PA, 2006
- [3] T. Barbu, V. Barbu, V. Biga, D. Coca, A PDE variational approach to image denoising and restoration. *Nonlinear Anal. Real World Appl.* **10** (2009), no. 3, 1351-1361.
- [4] V. Barbu, NONLINEAR DIFFERENTIAL EQUATIONS OF MONOTONE TYPES IN BANACH SPACES. *Springer Monographs in Mathematics*. Springer, New York, 2010
- [5] V. Barbu, A variational approach to stochastic nonlinear parabolic problems, *J. Math. Anal. Appl.* **384** (2011), no. 1, 2-15.
- [6] V. Barbu, Optimal control approach to nonlinear diffusion equation driven by Wiener noise, *J. Optim. Theory Appl.* **153** (2012), n. 1, 1-26.

- [7] V. Barbu, Th. Precupanu. CONVEXITY AND OPTIMIZATION IN BANACH SPACES Springer Monographs in Mathematics, Springer New York, 2011
- [8] V. Barbu, M. Röckner. On a random scaled porous media equation. *J. Diff. Equations* **251** (2011), 2494-2514.
- [9] V. Barbu, G. Da Prato, M. Röckner, Stochastic nonlinear diffusion equations with singular diffusivity. *SIAM J. Math. Anal.* **41** (2009), no. 3, 1106-1120.
- [10] V. Barbu, G. Da Prato, M. Röckner, Existence of strong solutions for stochastic porous media equation under general monotonicity conditions. *Ann. Probab.* **37** (2009), no. 2, 428-452.
- [11] V. Barbu, G. Da Prato, M. Röckner, Existence and uniqueness of nonnegative solutions to the stochastic porous media equation. *Indiana Univ. Math. J.* **57** (2008), no. 1, 187-211.
- [12] H. Brezis, I. Ekeland, Un principe variationnel associé à certaines équations paraboliques. Le cas indépendant du temps. *C. R. Acad. Sci. Paris Sér. A-B* **282** (1976), no. 17, A971-A974.
- [13] H. Brezis, I. Ekeland, Un principe variationnel associé à certaines équations paraboliques. Le cas dépendant du temps. *C. R. Acad. Sci. Paris Sér. A-B* **282** (1976), no. 20, A1197-A1198.
- [14] Z. Brzeźniak, M. Capiński, F. Flandoli. A convergence result for stochastic partial differential equations. *Stochastics* **24** (1988), no. 4, 423-445.
- [15] Z. Brzeźniak, Yuhong Li, Asymptotic compactness and absorbing sets for 2D stochastic Navier-Stokes equations on some unbounded domains, *Trans. Amer. Math. Soc.* **358** (2006), no. 12, 5587-5629
- [16] G. Da Prato, KOLMOGOROV EQUATIONS, Birkhauser, 2004
- [17] G. Da Prato, J. Zabczyk, STOCHASTIC EQUATIONS IN INFINITE DIMENSIONS, Cambridge University Press, 2008
- [18] G. Da Prato, L. Tubaro, Fully nonlinear stochastic partial differential equations. *SIAM J. Math. Anal.* **27** (1996), no. 1, 40-55
- [19] H. Doss, Liens entre equations différentielles stochastiques et ordinaires, *Ann. Inst. H. Poincaré* **13** (1977), n. 2, 99-125
- [20] T.E. Duncan, B. Maslowski, B. Pasik-Duncan, Stochastic equations in Hilbert space with a multiplicative fractional Gaussian noise, *Stochastic Process. Appl.* **115** (2005), 1357-1383
- [21] F. Flandoli, H. Lisei, Stationary conjugation of flows for Parabolic SPDEs with multiplicative noise and some applications. *Stochastic Analy. Appl.* **22** (2005), no. 2, 1385-1420.
- [22] A. Răşcanu, E. Rotenstein, The Fitzpatrick Function – A Bridge between Convex Analysis and Multivalued Stochastic Differential Equations. *Jour. Convex Anal.*, **18** (2008), n. 1, 105-138
- [23] H. Sussmann, On the gap between deterministic and stochastic ordinary differential equations. *Annals of Probability* **6** (1978), n.1, 19-41
- [24] A. Visintin, Extension of the Brezis-Ekeland-Nayroles principle to monotone operators. *Sci Appl. Adv. Math.* **18** (2008), 633-650

UNIVERSITY AL. I. CUZA AND INSTITUTE OF MATHEMATICS OCTAV MAYER,,
IAŞI, ROMANIA

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YORK
YO10 5DD, UK

DEPARTMENT OF MATHEMATICS AND INFORMATIONTECHNOLOGY, MONTA-
NUNIVERSITY LEOBEN,, FRANZ JOSEFSTR. 18, 8700 LEOBEN, AUSTRIA

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TRENTO, ITALY