

On Inviscid Limits for the Stochastic Navier-Stokes Equations and Related Models

Nathan Glatt-Holtz, Vladimír Šverák, and Vlad Vicol

ABSTRACT. We study inviscid limits of invariant measures for the 2D Stochastic Navier-Stokes equations. As shown in [Kuk04] the noise scaling $\sqrt{\nu}$ is the only one which leads to non-trivial limiting measures, which are invariant for the 2D Euler equations. We show that any limiting measure μ_0 is in fact supported on bounded vorticities. Relationships of μ_0 to the long term dynamics of Euler in the L^∞ with the weak* topology are discussed. In view of the Batchelor-Kraichnan 2D turbulence theory, we also consider inviscid limits for the weakly damped stochastic Navier-Stokes equation. In this setting we show that only an order zero noise (i.e. the noise scaling ν^0) leads to a nontrivial limiting measure in the inviscid limit.

1. Introduction

We consider incompressible Euler's equations and the randomly forced incompressible Navier-Stokes equation on a two-dimensional torus $\mathbb{T}^2 = \mathbb{R}^2/\Lambda$, where $\Lambda \subset \mathbb{R}^2$ is a lattice.¹ The Euler equation will mostly be considered in the vorticity form

$$\partial_t \omega + \mathbf{u} \cdot \nabla \omega = 0, \quad (1.1)$$

where we assume that $\int_{\mathbb{T}^2} \omega(x, t) dx = 0$ and the velocity field $\mathbf{u} = (u_1, u_2)$ is determined by ω from the equations

$$\operatorname{curl} \mathbf{u} = \nabla^\perp \cdot \mathbf{u} = \partial_1 u_2 - \partial_2 u_1 = \omega, \quad \operatorname{div} \mathbf{u} = 0, \quad \int_{\mathbb{T}^2} \mathbf{u} = 0. \quad (1.2)$$

In terms of the stream function, $\mathbf{u} = \nabla^\perp \psi$, $\Delta \psi = \omega$, with the usual notation $\nabla^\perp \psi = (-\psi_{x_2}, \psi_{x_1})$. The Navier-Stokes equation will be written either in the velocity formulation

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \frac{\nabla p}{\rho} - \nu \Delta \mathbf{u} = \mathbf{f}, \quad (1.3)$$

where ρ is the (constant) density and the forcing term $\mathbf{f} = \mathbf{f}(x, t)$ satisfies $\int_{\mathbb{T}^2} \mathbf{f}(x, t) dx = 0$ for each t , or in the vorticity formulation

$$\partial_t \omega + \mathbf{u} \cdot \nabla \omega - \nu \Delta \omega = \operatorname{curl} \mathbf{f}, \quad (1.4)$$

with the relation between \mathbf{u} and ω given as above. The particular stochastic form of \mathbf{f} will be discussed below (1.10).

In this paper we study certain classes of invariant measures for (1.1) and related equations. In particular, we address regularity properties and relations to the long term dynamics of 2D Euler for a particular class of invariant measures which arise as an inviscid limit of the stochastic Navier-Stokes equations, with a suitable scaling of the noise coefficients.

Date: February 3, 2013

¹The reason why we consider general flat tori, rather than just $\mathbb{R}^2/\mathbb{Z}^2$ is that the geometry of the torus might have some influence on various predictions concerning the long-time behavior of the solution. This issue will however come up only tangentially, and it will not be important for the results proved in the paper. The reader can take $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$ or $\mathbb{T}^2 = \mathbb{R}^2/2\pi\mathbb{Z}^2$ most of the time.

To discuss topics which come up in various accounts of 2D turbulence, we will also use a linear damping operator Y defined in the Fourier coordinates² by

$$\widehat{Y\mathbf{u}}_k = \gamma_k \hat{\mathbf{u}}_k, \quad (1.6)$$

where $\gamma_k \geq 0$. Often it is assumed that $\gamma_k \neq 0$ only for a few lowest modes, but other options, such as $Y\mathbf{u} = \gamma\mathbf{u}$ for some $\gamma > 0$ (corresponding to $\gamma_k = \gamma$ for each k) are also possible. We will denote the operator Y with $\gamma_k \neq 0$ only for a few low modes by Y_{low} . Its precise form will not be important for our discussion.

1.1. Two dimensional turbulence. The standard theory of 2D turbulence conjecturally describes the behavior of solutions of

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \frac{\nabla p}{\rho} - \nu \Delta \mathbf{u} + Y_{\text{low}} \mathbf{u} = \mathbf{f}, \quad (1.7)$$

where $\mathbf{f} = \mathbf{f}(x)$ is a “sufficiently generic” smooth vector field, which is on the Fourier side supported in a few relatively low Fourier modes.³ One can expect that for many \mathbf{f} the system will become “turbulent” for sufficiently low ν , although it is important to keep in mind that there are examples where this is not the case, see [Mar87, CR07]. In the turbulent regime one expects the famous downward cascade of energy together with an upward cascade of vorticity, as conjectured by Kraichnan [Kra67] and Batchelor [Bat69]. This is why we need the operator Y_{low} . In dimension $n = 3$ the operator Y_{low} should not be needed and we expect the so-called Kolmogorov-Richardson energy cascade. The striking feature of these phenomena is that, conjecturally, as $\nu \rightarrow 0_+$, the velocity field \mathbf{u} should satisfy some bounds independent of ν , such as

$$\|\mathbf{u}\|_{L^\infty} \leq C, \quad C \text{ independent of } \nu. \quad (1.8)$$

Moreover, there are conjectures as to how energy will be distributed in the Fourier modes (see, for instance, the classical works [KM80, MWC92, Fri95, Tsi01, FJMR02, Tab02]). Rigorous treatment of these scenarios seem to be out of reach of the present-day techniques and we have nothing new to say in this direction. Note however that the following 1D model given by the Burgers equation

$$\partial_t u + uu_x - \nu u_{xx} = f(x), \quad x \in \mathbb{R}/\mathbb{Z}, \quad \int u(x, t) dx = 0, \quad \int f(x) dx = 0, \quad (1.9)$$

is treatable. The behavior of the solutions for $\nu \rightarrow 0_+$ can be studied in detail via the Cole-Hopf transformation. In particular, the bound (1.8) can be established in this case.

Instead of relying on the chaos produced by the presumably complicated dynamics of (1.7) for low ν , we can input “genericity” into the system by considering a “random” \mathbf{f} . This point of view may be traced back to Novikov [Nov65] (see also, e.g. [BT73, VKF79]). We may take for example

$$\mathbf{f}(x, t) = \alpha \sum_k b_k \mathbf{e}_k(x) \dot{W}^k(t), \quad (1.10)$$

where $\mathbf{e}_k(x)$ is given by (1.5), the sum is finite, over a few relatively low modes, α is a suitable constant of order 1. The W^k are independent copies of the standard Wiener process (Brownian motion) so that \dot{W}^k are white noise processes and hence are stationary in time. After a suitable non-dimensionalization, a representative case of (1.10) is when $\sum_k |b_k|^2$ and α are both of order unity.⁴

² The Fourier representation is set up in the following way. For scalar functions v on \mathbb{T}^2 we will write $v(x) = \sum_{k \in 2\pi\Lambda^*} \hat{v}_k e^{ikx}$, where Λ^* is the lattice dual to Λ . Our functions v will satisfy $\int_{\mathbb{T}^2} v(x) dx = 0$ which is the same as $\hat{v}_0 = 0$ and the above sum will always be taken over the non-zero elements of $2\pi\Lambda^*$. If there is no danger of confusion we will write simply $v(x) = \sum_k \hat{v}_k e^{ikx}$. Divergence-free vector fields \mathbf{f} will be written as $\mathbf{f}(x) = \sum_k \hat{f}_k \mathbf{e}_k(x)$, where

$$\mathbf{e}_k(x) = \begin{pmatrix} -ik_2 & ik_1 \\ |k| & |k| \end{pmatrix} e^{ikx}. \quad (1.5)$$

We note that in our normalization we have $\frac{1}{|\mathbb{T}^2|} \int_{\mathbb{T}^2} |v|^2 dx = \sum_k |v_k|^2$, and $\frac{1}{|\mathbb{T}^2|} \int_{\mathbb{T}^2} |\mathbf{f}|^2 dx = \sum_k |f_k|^2$.

³The assumption of “sufficient genericity” is important, as we need that the system creates enough “chaos” for low ν .

⁴If we wish to consider dimensions of the various quantities, one natural choice seems to be to take \mathbf{e}_k dimension-less, b_k of the same dimension as \mathbf{u} (which is $[\text{length}][\text{time}]^{-1}$), and α of dimension $[\text{time}]^{-\frac{1}{2}}$, so that $\alpha \dot{W}^k(t)$ has dimension $[\text{time}]^{-1}$.

With a random forcing of the form (1.10), the equation (1.7) can then be viewed as a stochastic equation. With some additional assumptions, there will exist a unique invariant measure $\mu = \mu_\nu$ for the process defined by (1.7), see e.g. [FM95, DPZ96, Mat99, Mat02, BKL01, KS01, KS02, Mat03, MP06, HM06, Kup10, HM11, Deb11, KS12] and containing references. Relations conjectured by Kraichnan's theory would then be satisfied in a suitable mean value sense. The benefit of working with the random forcing is that even though we still cannot make much progress on establishing Kraichnan's conjectures in this setup, we now at least have a quite canonical object for our analysis, the measure μ .⁵ Indeed, the above mentioned works establish ergodic or even mixing properties of μ . These properties provide some theoretical justification for the measurement of the physical quantities described in turbulence theories. In the deterministic case the measure μ should presumably be replaced by a suitable invariant measure on the attractor, see e.g. [CF88, FMRT01].

One can of course go through similar considerations in three dimensions, but in that case the lack of rigorous results concerning the basic existence and uniqueness questions for the Navier-Stokes solutions prevents obtaining rigorously the measure μ above (or its analogues on the purported attractors in the deterministic case). Note however the recent works [DPD03, FR08, Deb11] on weaker notions of solutions and associated invariant measures.

1.2. Kuksin measures. Kuksin [Kuk04], see also [KP05, Kuk06b, Kuk06a, Kuk07, Kuk08, Shi11, KS12], suggested to study of a different limiting regime and put in (1.10)

$$\alpha = c\sqrt{\nu}, \quad (1.11)$$

where c is a constant independent of ν .⁶ Assume we have this scaling and omit the terms $\mathbf{u} \cdot \nabla \mathbf{u}$, Y_{low} from the equation. Then for each Fourier mode u_k we have

$$\mathbb{E}(|u_k|^2) = \frac{|b_k|^2 \alpha^2}{|k|^2 \nu} = |b_k|^2 \frac{c^2}{|k|^2}, \quad (1.12)$$

a bound independent of ν . In the non-linear case a similar bound can be obtained, see [Kuk04, KS12] and (1.17). With scaling (1.11) the operator Y is no longer needed⁷ and the invariant measures $\mu = \mu_\nu$ of the process generated by the equation

$$d\mathbf{u} + \left(\mathbf{u} \cdot \nabla \mathbf{u} + \frac{\nabla p}{\rho} - \nu \Delta \mathbf{u} \right) dt = c\sqrt{\nu} \sum_k b_k e_k(x) dW^k(t) \quad (1.13)$$

have a meaningful limit (perhaps after passing to a suitable subsequence). At the level of the vorticity $\omega = \text{curl } u$ we have

$$d\omega + (\mathbf{u} \cdot \nabla \omega - \nu \Delta \omega) dt = c\sqrt{\nu} \sum_k g_k dW^k(t) e^{ikx}, \quad g_k = |k| b_k, \quad (1.14)$$

which is the form which we will mostly work with.⁸ A deterministic version of the situation considered by Kuksin would correspond to setting

$$f = cv\tilde{f} \quad (1.15)$$

in (1.3). This situation is relevant for Section 5.

Let μ_ν be an invariant measure on the space of the vorticity functions ω of the process defined by (1.14). Note that for a sufficiently fast decay in the b_k 's in (1.13), these measures μ_ν are supported on smooth

⁵The case of the Burgers equation (1.9) with stochastic forcing can be analyzed rigorously, see for example [EKMS00].

⁶If our quantities are not dimensionless and we use the same dimension count as in a previous footnote, then c should have dimension $[\text{length}]^{-1}$.

⁷Indeed in this scaling, (1.11), the term Y leads to a trivial limiting measure μ as we establish rigorously in Section 6 below.

⁸The term $-\nu \Delta \omega$ can be replaced by more general dissipation, such as fractional Laplacian $\nu(-\Delta)^\alpha$; see Remark 4.5 below.

functions, see e.g. [KS12]. Also, under rather general assumptions on these b_k 's such a measure is unique, cf. references above. Applying Ito's formula

$$d \int_{\mathbb{T}^2} \frac{1}{2} \omega^2 dx = \int_{\mathbb{T}^2} (\omega d\omega + \frac{1}{2} d\omega d\omega) dx \quad (1.16)$$

and taking the expectation, we obtain

$$\mathbb{E} \left(\int_{\mathbb{T}^2} |\nabla \omega|^2 dx \right) = \int \|\nabla \omega\|_{L^2}^2 d\mu_\nu(\omega) = \frac{c^2}{2} \sum_k |g_k|^2. \quad (1.17)$$

Due to this bound, as $\nu \rightarrow 0_+$, the family of measures μ_ν has a subsequence converging weakly⁹ to a limit μ_0 , which is a measure supported in the space

$$H^1 = \left\{ \omega \in H^1(\mathbb{T}^2), \int_{\mathbb{T}^2} \omega dx = 0 \right\}, \quad (1.18)$$

with

$$\mathbb{E}(\|\nabla \omega\|_{L^2}^2) = \int \|\nabla \omega\|_{L^2}^2 d\mu_0(\omega) \leq \frac{c^2}{2} \sum_k |g_k|^2. \quad (1.19)$$

Kuksin [Kuk04] proves that this measure invariant for the evolution given by Euler's equation (1.1). See also Kuksin and Shirikian [KS12] and references therein for many other interesting properties, such as the non-triviality of the measure. We will call the measures μ_0 constructed in this way *Kuksin measures*.

1.3. Main Result on Kuksin measures and the dynamical systems approach to 2D Euler. We now show that the Kuksin measures are closely related to the long-time behavior of solutions of Euler's equation. One of our main results in this paper will be the following:

Theorem 1.1 (Kuksin measures are supported on L^∞). *Let μ be a Kuksin measure as above. Then*

$$\int \|\omega\|_{L^\infty} d\mu_0(\omega) < +\infty. \quad (1.20)$$

We shall discuss the outline of the proof of Theorem 1.1 in Subsection 1.4 below. A detailed statement of the result and its proof is found in Section 4.

In particular, from (1.20) we see that μ is supported on L^∞ . This is important, as the space L^∞ is probably the most natural space (for the vorticities) in which to consider the 2D Euler equation when studying the long-time behavior of the solution. This is due to the following facts:

- (i) The initial value problem for (1.1) is well-posed for in L^∞ , a classical result by Yudovich [Jud63].
- (ii) Let $R \geq 0$ and let $X = X_R = \{\omega \in L^\infty, \int_{\mathbb{T}^2} \omega = 0, \|\omega\|_{L^\infty} \leq R\}$. Equipped with the weak* topology, the set X is a compact metric space, which we will denote by (X, w^*) . One can check that the proof of Yudovich's theorem actually give a stronger result: namely, *the Euler equation (1.1) gives a well-defined dynamical system on X (for any $R > 0$).* A proof of Yudovich's theorem which can be easily adapted to prove our statement here can be found for example in [MB02].

From Theorem 1.1 we hence see that Kuksin measures (restricted to X) give natural invariant measures for the Euler evolution on X . The functions on which the measures are supported have additional H_0^1 -regularity.

Note that one can also construct non-trivial measures on X which are invariant under the Euler equation directly: we know that the energy functional¹⁰

$$\mathcal{E}(\omega) = \frac{1}{|\mathbb{T}^2|} \int_{\mathbb{T}^2} \frac{1}{2} |u|^2 dx = \frac{1}{|\mathbb{T}^2|} \int_{\mathbb{T}^2} -\frac{1}{2} \psi \omega dx \quad (1.21)$$

is continuous on (X, w^*) . Therefore the energy level sets $X_E = X_{R,E}$ given by $\{\omega \in X, \mathcal{E}(\omega) = E\}$ are compact subsets in X which are invariant under the Euler equation (due to the energy conservation). By the classical

⁹More precisely we have that $\int_{L^2} f(\omega) d\mu_{\nu_j}(\omega)$ converges to $\int_{L^2} f(\omega) d\mu_0(\omega)$ for every continuous, bounded real valued test function f . In fact the convergence holds also in $H^{1-\varepsilon}$ for any ε positive.

¹⁰More precisely, energy per unit mass.

Kryloff-Bogoliouboff procedure, every non-empty $X_{R,E}$ supports an invariant measure. This measure cannot be trivial when $E > 0$ i.e. supported at $\omega = 0$. There are additional conserved quantities for the evolution by Euler's equation, namely the integrals

$$I_F(\omega) = \frac{1}{|\mathbb{T}^2|} \int_{\mathbb{T}^2} F(\omega) dx, \quad (1.22)$$

but these quantities are not continuous on (X, w^*) , making their implications for the dynamics on (X, w^*) more subtle.

In Section 2 we discuss various hypothesis for the long term dynamics of the (1.1) on the phase space (X, w^*) . These hypothesis may illuminate further possible structure of the support of the invariant measures μ_0 . Two extreme scenarios present themselves. On, the one hand, taking the view of statistical mechanics, we may predict long time behavior from the maximization of various notions of ‘‘entropy’’ subject to the constrains of the Eulerian dynamics. In many cases these ‘‘entropy maximizers’’ may have a fairly simple shear flow like structure. Thus in this scenario we would expect that the Kuksin measures would be supported on steady flows with a relatively simple topology. At the other extreme we might suppose that all of the solution trajectories of the Euler dynamical system are pre-compact in L^2 . In this case many of the Casimirs (1.22) must be conserved at the end-states which would suggest that μ_0 has a much richer structure.

There is some evidence for both of the above scenarios. On the one hand we do not have a single example where it is proved that an initial condition yields an orbit which is not precompact in L^2 . Moreover a recent result in [Šve12] (which we recall in Theorem 2.1 below) rigorously shows that at least some such precompact orbits must exist. On the other hand recent numerical result of [BS09] suggest that μ_0 is concentrated on certain laminar states obtained as an ‘‘Entropy maximization’’. It seems unlikely that either of these scenarios holds universally and that the structure of μ_0 is given by an intermediate situation.

1.4. Moser iteration for SPDE and applications to L^∞ estimates for stationary solutions. We now turn to discuss some aspects of the proof of Theorem 1.1. We will see that the main ingredients involve a suitable rescaling of the equations and then developing a Moser iteration scheme for SPDEs of drift diffusion type which evidences a parabolic regularization from L^2 to L^∞ . The detailed proofs are given below in Section 4.

A natural rescaling of time makes the interpretation of the measures μ_ν and the Kuksin measures μ_0 perhaps more transparent. If we replace the function $\mathbf{u}(x, t)$ by $\tilde{\mathbf{u}}(x, t) = \mathbf{u}(x, t/\nu)$ and replace $W^k(t)$ by the equivalent process $\tilde{W}(t) = \sqrt{\nu}W^k(t/\nu)$, we obtain, after dropping the tildes we obtain

$$d\omega + \left(\frac{1}{\nu} \mathbf{u} \cdot \nabla \omega - \Delta \omega \right) dt = c \sum_k g_k dW^k(t). \quad (1.23)$$

Note that the measure μ_ν is also an invariant measure for this process. See Section 2.1 below for some motivating discussion of analogous finite dimensional situations.

As $\nu \rightarrow 0$ in (1.23) the drift velocity $\nu^{-1} \mathbf{u}$ grows unboundedly. As such to obtain Theorem 1.1 need find a way estimate L^∞ norms of solutions to equations of the form

$$d\omega + (\mathbf{a} \cdot \nabla \omega - \Delta \omega) d\tau = c \sum_k g_k dW^k, \quad \nabla \cdot \mathbf{a} = 0 \quad (1.24)$$

with constants that *do not* depend on the *size* of the sufficiently smooth divergence free drift velocity \mathbf{a} .¹¹

In the deterministic case, one usually obtains such drift-independent L^∞ bounds either by appealing to maximum principle-type arguments, or by using L^p estimates, with p independent bounds, and passing $p \rightarrow \infty$. Neither of these direct approaches appear to be available in the stochastic case. The first approach seems to fail since one cannot exchange \mathbb{E} and \sup_x , and due to the lack of smoothness in time of the

¹¹Since the noise in (1.24) is additive, one could shift the equation by subtracting the solution of an Ornstein-Uhlenbeck process, but the L^∞ bounds one obtains on the resulting random PDE appear to depend essentially on the size of the drift velocity.

stochastic terms in (1.24). On the other hand, for L^p bounds, a direct application of the Itô lemma to (1.24) yields

$$d\|\omega\|_{L^p}^p = \left(p\langle \Delta\omega, \omega|\omega|^{p-2} \rangle + \frac{p(p-1)}{2} \sum_k \|g_k|\omega|^{(p-2)/2}\|_{L^2}^2 \right) dt + p \sum_k \langle g_k, \omega|\omega|^{p-2} \rangle dW^k \quad (1.25)$$

where we have used that \mathbf{a} is divergence-free. The Itô correction term in (1.25) grows quadratically in p , which is too fast. On the other hand, letting $X = \|\omega\|_{L^p}^p$, one may apply Itô's lemma to $\phi(X) = (1+X)^{2/p}/p$ (see e.g. [Kry10, Remark 5.2]) and prove using standard estimates that

$$\sup_{p \geq 2} \left(\mathbb{E} \sup_{t \in [0, T]} \frac{\|\omega(t)\|_{L^p}^2}{p} \right) \leq C \left(1 + \sup_{p \geq 2} \left(\mathbb{E} \frac{\|\omega_0\|_{L^p}^2}{p} \right) + \nu T \|\sigma\|_{L^\infty} \right)$$

where ω is the solution of (3.2). This however does not yield a bound on $\mathbb{E} \left(\sup_{t \in [0, T]} \sup_{p \geq 2} \|\omega(t)\|_{L^p}^2 / p \right)$.

Since (1.24) is a parabolic SPDE, in the spirit of the classical DeGiorgi-Nash-Moser [DG57, Nas58, Mos60] theory for deterministic parabolic PDEs, one may expect an instant regularization of the solution. The difficulty in carrying over this program lies in treating the stochastic forcing term in (1.24) and obtaining bounds which are independent of the size of drift velocity \mathbf{a} . In the deterministic case, for drift velocities that are divergence free, one obtains the L^2 to L^∞ regularization of solutions to the parabolic equation, with bounds that are *independent* of the drift using e.g. the elegant argument of Nash [Nas58]. Drift independent bounds for a deterministic analogue of (1.24) have also been obtained using Moser iteration, see, e.g. [Kuk99]. Therefore, one may expect that the same result holds for stochastic drift-diffusion equations such as (1.24).

In order to treat the stochastic term, it turns out that the iteration technique introduced by Moser [Mos60] is better suited in view of the L^p Itô formula (1.25). This fact was recently observed in the context of semilinear SPDE in [DMS05, DMS09] where the authors obtain an L^∞ maximum principle. We however cannot appeal to these results since they rely essentially on the fact that the initial data already lies in L^∞ . By (1.17) we only have ν -independent H^1 bounds on the statistically steady solutions of (1.14). To overcome this difficulty we prove in Theorem 4.1 (see also Remark 4.4 below) that the solution $\omega(t)$ of (1.24) lies in L^∞ (in x) for arbitrarily small positive time t :

$$\mathbb{E} \sup_{t \in [T, 2T]} \|\omega(t)\|_{L^\infty} \leq C \left(1 + T^{-5/4} \right) \mathbb{E} \left(\|\omega\|_{L^4([0, 2T]; L^2)} \vee c \sum_k \|g_k\|_{L^\infty} \right), \quad (1.26)$$

for $T \leq 1/8$, where the constant C is independent on \mathbf{a} . To the best of our knowledge the parabolic regularization estimate (1.26) is new in the context of SPDE. As in [DMS05, DMS09], one of the main differences between the stochastic Moser iteration (see the proof of Theorem 4.1) and the classical approach for deterministic PDE naturally arises due to the random forcing. In view of the Burkholder-Davis-Gundy inequality we need to bound quadratic variations of the Martingale on the right side of (1.25), and hence the integrability in time needs to be twice that in space in order to close the iteration scheme (cf. (4.32) below).

In view of the predictions made by Statistical Mechanics arguments regarding the “end states” of the 2D Euler dynamics, and having already established that the Kuksin measures are supported on $H^1(\mathbb{T}^2) \cap L^\infty(\mathbb{T}^2)$, we believe that:

Conjecture 1.2. *Kuksin measures are in fact supported on continuous vorticities.*

The immediate difficulty which arises in proving this conjecture is that even in the deterministic case, for the two-dimensional linear parabolic equation

$$\partial_t v + \mathbf{b}(x, t) \cdot \nabla v - \Delta v = f, \quad \nabla \cdot \mathbf{b} = 0, \quad (1.27)$$

the size of the smooth drift comes into play for the DeGiorgi-Nash-Moser proof of Hölder regularity. If the drift is rough, one may even construct solutions that are not continuous functions for all time, although they obey the L^∞ maximum principle (see, e.g. [SVZ12]).

On the other hand, one of the key ingredients of the proof of Theorem 4.1 was the (statistical) stationarity of the solution to (1.23). As a deterministic toy problem one may hence consider time-independent solutions of (1.27), with drift $\mathbf{b}(x)$. In this case, following the ideas in [SSŠZ12] and an elliptic Moser iteration we are indeed able to prove in Theorem 5.1 below that the solution obeys a drift-independent logarithmic modulus of continuity, and is hence uniformly continuous. The analogy between time-independent solutions to (1.27) and statistically stationary solutions of (1.24) is however tentative at best.

1.5. Inviscid limits for damped models; different scalings. In view of the foregoing discussion concerning the Batchelor-Kraichnan theory of 2D turbulence, it appears that when working on the periodic box the stochastic Navier-Stokes equations should be augmented (as in (1.7)) with a suitable damping term Y to prevent a pile up of energy at large scales. Note that the Y term also frequently appears in geophysical models closely related to the 2D Navier-Stokes equations to account for friction with boundaries. In these situations, if the scaling in this damping term is held fixed as $\nu \rightarrow 0$, then a different scaling must be introduced for the noise in order to obtain a non-trivial inviscid limit in the class of the associated invariant measures.

To this end, we consider operators of the form $Y = Y_{\tau,\gamma} = \tau \Lambda^{-\gamma} = \tau (-\Delta)^{-\gamma/2}$ and $\tau > 0$, $\gamma \in [0, 1)$ and study weakly damped and driven stochastic Navier-Stokes equations of the form

$$d\mathbf{u} + (Y\mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla \pi - \nu \Delta \mathbf{u}) dt = \nu^\alpha \rho dW, \quad \nabla \cdot \mathbf{u} = 0, \quad (1.28)$$

for different values of $\alpha \in \mathbb{R}$.¹² As above for the undamped case (1.17) energetic considerations allow us to deduce the correct scaling with α in (1.28). Consider a collection of invariant measures $\{\mu_\nu^\alpha\}_{\nu>0}$ for (1.28). Let \mathbf{u}^ν be stationary solutions of (1.28) corresponding to μ_ν^α and denote $\omega^\nu = \nabla^\perp \cdot \mathbf{u}^\nu$. Applying the Itô lemma to the vorticity formulation of (1.28) and using stationarity one deduces that:

$$\mathbb{E} (\nu \|\nabla \omega^\nu\|_{L^2}^2 + \|Y^{1/2} \omega^\nu\|_{L^2}^2) = \frac{\nu^{2\alpha}}{2} \|\sigma\|_{L^2}^2.$$

Making use of the above relation, we will show below in Theorem 6.1 that $\alpha = 0$ is the only relevant scaling for (1.28). Here stationary solutions of a damped stochastic Euler equation arise. See also [BF12, Bes08].

Organization of the Paper. In Section 2 we review some notions related to the time-asymptotic behavior of the 2D Euler equations. Our discussions in this section allow us to make some hypotheses regarding the structure of the support of Kuksin measures in this context. Section 3 recalls the mathematical framework for the Navier-Stokes Equations and its associated Markovian semigroup. We also review some properties of Kuksin measures established in previous works. Section 4 is devoted to the proof of the main theorem. Here we detail the Moser iteration scheme which addresses a more general class of drift-diffusion equations. Section 5 concerns a deterministic toy model for stationary solutions of the stochastic Navier-Stokes equations. We establish a modulus of continuity for this system. The final Section 6 we consider a weakly damped stochastic Navier-Stokes equation and establish inviscid limits in the appropriate scaling for this model.

2. Long term behavior of 2D Euler and related systems; connections to invariant measures

In this section we discuss some aspects of the long time dynamics in (L^∞, w^*) of solutions to 2D Euler and the relation of this behavior to possible properties of the Kuksin measures, which are now accessible due to Theorem 1.3. We begin with some motivation from finite dimensional Hamiltonian systems.

2.1. Noise scaling limits in finite dimensions. A finite dimensional situation related to the above is studied in the theory of the small random perturbations of dynamical systems. Let

$$\dot{x} = b(x) \quad (2.1)$$

¹²As explained above it is of interest to consider Y acting only at the largest scales; cf. (1.6). This situation is more delicate to analyze rigorously and would seem to require the establishment of suitable ‘‘hypocoercivity’’ properties for (1.7).

be a dynamical system in \mathbb{R}^n . Consider its stochastic perturbation

$$dx = b(x) dt + \sqrt{\varepsilon} Q dW, \quad (2.2)$$

where $W = (W^1, \dots, W^n)$ are normalized independent Wiener processes and Q is a matrix. By setting $x(t) = \tilde{x}(\varepsilon t)$, $W(t) = \frac{1}{\sqrt{\varepsilon}} \tilde{W}(\varepsilon t)$ and $\tau = \varepsilon t$ and dropping the tildes, we obtain

$$dx = \frac{1}{\varepsilon} b(x) d\tau + Q dW. \quad (2.3)$$

Such systems have been extensively studied, see e.g. [FW12]. In case of measure-preserving flows, the behavior of (2.3) as $\varepsilon \rightarrow 0_+$ can be understood from the following picture: under some assumptions the equation $\dot{x} = \frac{1}{\varepsilon} b(x)$ takes the trajectories very quickly through “ergodic components”, and hence for $\varepsilon \rightarrow 0_+$ the system (2.3) should in some sense describe a diffusion in the space of the ergodic components.

Equation (1.13) (or its rescaled version (1.23)) is of a slightly different nature than the perturbation of Hamiltonian systems considered in [FW12], in that we add not only a small noise, but also small damping. Such procedure can be illustrated by a simple example of the Langevin oscillator:

Example (The Langevin oscillator). We consider a simple 1d harmonic oscillator with damping and random forcing

$$m\ddot{q} + \gamma\dot{q} + \kappa q = \alpha \dot{w}, \quad (2.4)$$

where w is a normalized Wiener process and $m, \gamma, \kappa > 0$. Letting $p = m\dot{q}$ as usual, it is easy to check that the (unique) invariant measure of the system

$$\begin{aligned} \dot{q} &= \frac{p}{m} \\ \dot{p} &= -\kappa q - \frac{\gamma}{m} p + \alpha \dot{w} \end{aligned} \quad (2.5)$$

is given by the *Gibbs measure*

$$d\mu = \frac{1}{Z} e^{-\beta H(q,p)} dq dp, \quad (2.6)$$

where the Hamiltonian H given by

$$H = \frac{p^2}{2m} + \frac{\kappa q^2}{2}, \quad \beta = \frac{\gamma}{\alpha^2}, \quad (2.7)$$

and $Z = Z(\beta, \kappa, m)$ is a suitable normalizing constant. We see that from the point of view of Statistical Mechanics the quantity $\frac{\alpha^2}{\gamma}$ corresponds to (a multiple of) temperature. A similar calculation can be done for a general one dimensional Hamiltonian of the form

$$H = \frac{p^2}{2m} + V(q). \quad (2.8)$$

In higher dimensions one can also calculate further examples; especially when the Hamiltonian is quadratic. The invariant measure does not necessarily have to be the Gibbs canonical measure as in (2.6). If the damping and the forcing are taken to 0 with the analogue of the ratio $\frac{\alpha^2}{\gamma}$ converging to a limit, the invariant measure will converge to an invariant measure of the original Hamiltonian system. For example, in the case of a completely integrable n -dimensional system with the full system of mutually commuting integrals of motion f_1, \dots, f_n the limiting invariant measure can be expected to be of the form

$$d\mu = \frac{1}{Z} e^{-\phi(f_1, \dots, f_n)} dq_1 \dots dq_n dp_1, \dots, dp_n, \quad (2.9)$$

where the function ϕ will depend on specific choices of damping/forcing. We see that the vanishing damping/random forcing method can be viewed as a way of producing suitable statistical “ensembles”, closely related to those used in Statistical Mechanics. Considerations in this direction in the context of the KdV equation can be found in [Kuk07]. In the terms of Statistical Mechanics the ensembles produced by this method are related to “canonical ensembles”. One can also consider the “micro-canonical ensembles”,

which in the last examples would simply be given (under some “genericity assumptions”) by invariant measures on the tori

$$f_1 = c_1, f_2 = c_2, \dots, f_n = c_n. \quad (2.10)$$

Under suitable assumptions, invariant measures on these tori are the “irreducible components” of the measures (2.9). By analogy, we see that Kuksin measures should be related to the Statistical Mechanics of Euler’s equation. Their decomposition into irreducible components should give “ergodic components” of the Euler evolution. However, this analogy may break down due to infinite dimensional effects. As $\nu \rightarrow 0_+$, it is conceivable that fast Euler evolution moves enstrophy to high spatial frequencies (in the Fourier spectrum), so that “complexity” is lost (by disappearing to infinity in the Fourier space) and Kuksin measures may conceivably be supported on some relatively simple sets, perhaps even equilibria. This would be an infinite-dimensional effect,¹³ which does not have an analogy for finite dimensional or completely integrable hamiltonian systems. This is discussed in more details below, but still at a heuristic level. We were not able to obtain rigorous results in this direction.

2.2. Long-time behavior of solutions of Euler’s equation. Equation (1.23) together with some analysis of the long-time behavior of solutions of Euler’s equation seems to give some good hints about what one should expect concerning some of the properties of Kuksin measures. We recall some of the expected properties of the Euler solutions.

We consider equation (1.1), with the conventions (1.2). We also recall the obvious identity

$$\int_{\mathbb{T}^2} \omega(x, t) dx = 0, \quad t \in \mathbb{R}. \quad (2.11)$$

Let us start with some classical observations about the long-time behavior of the solutions of (1.1) starting from initial data $\omega_0 \in L^\infty$, with $\|\omega_0\|_{L^\infty} = R$. Let $X = X_R$ be the ball of radius R in L^∞ , taken with the weak* topology. In addition, we can impose the constraint $\int_{\mathbb{T}^2} \omega dx = 0$ on the functions in X . The space (X, w^*) is a metric space and, as already discussed, the Euler evolution (1.1) gives a well-defined dynamical system on X . We denote the Ω -limit sets by

$$\Omega_+ = \Omega_+(\omega_0) = \bigcap_{t>0} \overline{\{\omega(s), s \geq t, \omega(0) = \omega_0\}}^{w^*}. \quad (2.12)$$

We also introduce

$$\mathcal{O}_{\omega_0} = \{\omega_0 \circ h, h: \mathbb{T}^2 \rightarrow \mathbb{T}^2 \text{ is a volume-preserving } C^1\text{-homeomorphism}\} \quad (2.13)$$

and

$$\mathcal{O}_{\omega_0, E} = \mathcal{O}_{\omega_0} \cap \{\omega, \mathcal{E}(\omega) = E\}. \quad (2.14)$$

It is not hard to see that the weak* closure of \mathcal{O}_{ω_0} , denoted by $\overline{\mathcal{O}_{\omega_0}}^{w^*}$ is a closed convex subset of L^p for any $p \geq 1$. Letting $\mathcal{E}(\omega_0) = E$, we clearly have

$$\Omega_+ \subset \overline{\mathcal{O}_{\omega_0, E}}^{w^*}. \quad (2.15)$$

There are various conjectures concerning the long-time behavior of Euler solutions motivated by the notion of “mixing”, see [Mil90, Rob91, Shn93, Šve12]. We can think of the fluid as consisting of fluid particles, with each fluid particle having a fixed value of vorticity permanently attached to it. The fluid motion mixes these particles, with the vorticity remaining attached to each particle. The most naive conjecture could be that for large times the vorticity is everywhere mixed, corresponding to the weak* convergence of $\omega(t)$ to 0 as $t \rightarrow \infty$. This would mean¹⁴

$$\Omega_+ = \{0\}. \quad (2.16)$$

In the Fourier space this would correspond to the movement of all energy¹⁵ of the solution $\hat{\omega}_k(t)$ towards larger and larger frequencies as $t \rightarrow \infty$. If $E \neq 0$, then (2.15) provides an obstacle to this. The energy cannot

¹³The effect is closely related to Landau damping, see for example [MV11].

¹⁴This presumably happens if we consider the Burgers equation with the scaling (1.13). It should not be hard to verify that for the Burgers equation the Kuksin measures are trivial.

¹⁵It would be more precise to say *enstrophy*, but in the situation here this does not make a difference.

all move to high (spatial) frequencies, as the energy functional \mathcal{E} is weakly* continuous. We can “fix” this by trying to “move” as much as possible energy to high frequencies which is still compatible with (2.15). More specifically, we can try to minimize

$$I(\omega) = \frac{1}{|\mathbb{T}^2|} \int_{\mathbb{T}^2} |\omega|^2 dx = \sum_k |\omega_k|^2 \quad (2.17)$$

subject to the constraint

$$\omega \in \overline{\mathcal{O}}_{\omega_0, E}^{w*}. \quad (2.18)$$

Note that $I(\omega)$ is preserved during the evolution, but it can conceivably drop on the “end-states” Ω_+ , as it is not weakly* continuous.

Minimizing I (subject to (2.18)) is of course the same as maximizing $-I(\omega)$ subject to (2.18). More generally, we can take a concave function F and maximize $I_F(\omega)$ given by (1.22), subject to (2.18). The quantity $I_F(\omega)$ can be called the *entropy* of the “configuration” ω and the above principle is then nothing but the usual entropy maximization under given constraints, as well-known from Statistical Mechanics. The entropy I_F could be considered as too simple, the usual entropy in Statistical Mechanics is based on suitable “counting of states”. Closely related to the notion of entropy is A. Shnirelman’s notion of mixing in [Shn93].

There are indeed more sophisticated notions of entropy, see for example [Mil90, Rob91, Tur99, Šve12], which can be more “non-local” than the I_F above. For example, let $\omega_0 = \sum_l a_l \chi_{A_l}$, where A_l is a division of \mathbb{T}^2 into disjoint measurable sets with $|A_l| = \varkappa_l |\mathbb{T}^2|$, and $\sum_l a_l \varkappa_l = 0$. Then, the closure of \mathcal{O}_{ω_0} defined in (2.13) is

$$\overline{\mathcal{O}}_{\omega_0}^{w*} = \left\{ \omega : \omega(x) = \sum_l a_l \rho_l(x), \quad 0 \leq \rho_l \leq 1, \quad \sum_l \rho_l = 1 \right\} \quad (2.19)$$

and one can define the entropy (generated by ω_0) as

$$S(\omega) = S_{\omega_0}(\omega) = \sup \left\{ \frac{1}{|\mathbb{T}^2|} \int_{\mathbb{T}^2} \sum_l -\rho_l \log \rho_l dx : \omega(x) = \sum_l a_l \rho_l(x), \quad 0 \leq \rho_l \leq 1, \quad \sum_l \rho_l = 1 \right\}. \quad (2.20)$$

This entropy leads to the theories of Miller and Robert, [Mil90, Rob91]. When the division $\mathbb{T}^2 = \cup_l A_l$ has only two sets A_1 and A_2 , then this entropy is of the form I_F for a suitable F . For example, when we only have two sets and $a_1 = -a_2 = 1$, then

$$F(\omega) = -\left(\frac{1+\omega}{2}\right) \log\left(\frac{1+\omega}{2}\right) - \left(\frac{1-\omega}{2}\right) \log\left(\frac{1-\omega}{2}\right). \quad (2.21)$$

The maximizers of the entropy subject to given constraints are steady-state solutions of Euler’s equations given by stream functions ψ satisfying

$$\Delta \psi = H(\psi) \quad (2.22)$$

for a suitable function H depending on I_F . These should be the “end-states” of the evolution if the actual evolution and Statistical Mechanics lead to the same conclusions.

As is usually the case with predictions based on Statistical Mechanics considerations, it is difficult to decide whether the actual dynamics of the equation produces the behavior expected from entropy maximization, assuming all known the constraints have been taken into account. In fact, we do not have a single example in which it would be rigorously established that the trajectory

$$\Omega_t = \cup \{\omega(s), s \geq t\} \quad (2.23)$$

is not pre-compact in L^2 (and hence any L^p for $p \in [1, \infty)$). On the other hand, it is useful to recall the following result from [Šve12]:

Theorem 2.1 (Existence of L^2 precompact orbits [Šve12]). *The omega-limit set Ω_+ of any trajectory always contains an element $\tilde{\omega}_0$ whose trajectory is pre-compact in L^2 .*¹⁶

¹⁶The proof of this statement is very simple: maximize some entropy I_F with a strictly concave F over Ω_+ .

2.3. Possible Consequences for Kuksin Measures. In view of the above discussions concerning the long term dynamics of 2D Euler we now introduce two extreme scenarios:

Scenario A: Euler solutions weakly* approach entropy maximizers. Let us assume that our torus is $\mathbb{T}^2 = \mathbb{R}^2/a\mathbb{Z} \oplus b\mathbb{Z}$ with $0 < a < b$. Let us further assume that

- (i) All entropy maximizers for Euler solutions (with given constraints) are shear flows independent of x_1 . This is in fact not very far-fetched. It has been established rigorously for sufficiently small energies and local entropies I_F with F strictly concave. See [FŠ].
- (ii) All solutions weakly* approach these shear flows as $t \rightarrow \infty$. This would of course be a very strong statement which we do not really expect to be true. However, if (i) is correct, then the Statistical Mechanics predictions would suggest exactly this conclusion.

If this scenario holds, then one can expect the Kuksin measures to be supported on the steady-state shear flows. Indeed, from the re-scaled equation (1.23) we see that as $\nu \rightarrow 0$, the fast Euler dynamics will drive the solution towards the shear flows, whereas the term $\Delta\omega$ will be quickly damping the high frequency components of ω generated by the Euler evolution. This scenario is genuinely infinite-dimensional: all the complexity of the Euler dynamics and the initial data will disappear into the high frequencies, and will never “return”. Such behavior does not have an analogy in finite-dimensional systems or in completely integrable systems.

Scenario B: all solution orbits are pre-compact. Let us assume that the solution trajectories Ω_t in (2.23) are pre-compact in L^2 . This may be unlikely, but it has not been ruled out. In this case the weak closures of these trajectories will be the same as the strong closures and all the functionals I_F will be conserved on the “end-states”. In particular, the mixing envisaged by the statistical mechanics approach will never take place. In this case the Kuksin measures will have much richer structure. Their “irreducible components”, similar to the measures on the tori (2.10) in the example leading to (2.9), will be supported on the closures of the L^2 -compact trajectories, which will play a role somewhat similar to the ergodic components in finite-dimensional Hamiltonian systems. In this scenario many features familiar from finite dimensions or completely integrable systems will still be present.

We conjecture that neither of these scenarios is quite true, but that the real behavior will be intermediate between these two extremes: non-trivial L^2 -precompact trajectories will exist, but initial data leading to them will not be “generic”. The Kuksin measures will be supported on such trajectories. Depending on our degree of optimism, we can hope that these solutions represent a type of a weak* attractor for all Euler solutions.

3. Some results regarding invariant measures and inviscid limits

In this section we first recall some elements of the mathematical analysis of the stochastic Navier-Stokes equations and its associated ergodic properties. This allows us then to summarize some of the analytical properties enjoyed by the Kuksin measures, established in previous works (cf. [KS12] and references therein).

3.1. Mathematical setting: stochastic 2D NSE and its Markov semigroup. We consider the Stochastic Navier-Stokes Equations on a periodic box \mathbb{T}^2

$$d\mathbf{u} + (\mathbf{u} \cdot \nabla \mathbf{u} + \nabla \pi - \nu \Delta \mathbf{u}) dt = \sqrt{\nu} \rho dW = \sqrt{\nu} \sum_k \rho_k dW^k, \quad \nabla \cdot \mathbf{u} = 0, \quad \mathbf{u}(0) = \mathbf{u}_0. \quad (3.1)$$

As discussed in the Introduction, in order to consider the inviscid limit $\nu \rightarrow 0$, we use the scaling $\sqrt{\nu}$ for the noise coefficient. Typically we will use the vorticity formulation of (3.1). Taking $\omega = \nabla^\perp \cdot \mathbf{u}$ we obtain

$$d\omega + (\mathbf{u} \cdot \nabla \omega - \nu \Delta \omega) dt = \sqrt{\nu} \sigma dW = \sqrt{\nu} \sum_k \sigma_k dW^k, \quad \omega(0) = \omega_0. \quad (3.2)$$

We will assume that $\int_{\mathbb{T}^2} \omega_0 dx = 0$ and $\int_{\mathbb{T}^2} \sigma dx = 0$, which implies that the solution is always mean zero. Note that \mathbf{u} can be recovered from ω via the Biot-Savart law.

Let us now set some notation used throughout the rest of the work. We denote the Sobolev spaces

$$H^k = \left\{ \omega \in H_{per}^k : \int_{\mathbb{T}^2} \omega_0 dx = 0 \right\},$$

with $H^0 = L_{per}^2$ and take the usual norms and inner products denoted by $\|\cdot\|_k$, $(\cdot, \cdot)_k$. We will denote the L^p , $p \geq 1$ by $\|\cdot\|_{L^p}$.

To emphasize dependence on initial conditions we will write $\omega^\vee(t, \omega_0) = \omega(t, \omega_0)$ for the solution of (3.2) with initial condition ω_0 . Assuming

$$\sum_l \|\sigma_l\|_{L^2}^2 < \infty, \quad (3.3)$$

we have $\omega(\cdot, \omega_0) \in C([0, \infty); H^0) \cap L_{loc}^2([0, \infty); H^1)$ for any $\omega \in H^0$. For $k \geq 1$, assuming that

$$\sum_l \|\sigma_l\|_{H^k}^2 < \infty, \quad (3.4)$$

we also have the higher regularity properties for (3.2). If $\omega_0 \in H^0$ then, for any $t_0 > 0$, $\omega(\cdot, \omega_0) \in C([t_0, \infty); H^k) \cap L_{loc}^2([t_0, \infty); H^{k+1})$. Similarly, if $\omega_0 \in H^k$, $\omega(\cdot, \omega_0) \in C([0, \infty); H^k) \cap L_{loc}^2([0, \infty); H^{k+1})$. Note that the general well-posedness theory for the stochastic Navier-Stokes equations has been extensively developed. See e.g. [BT72, BT73, Vio76, Cru89, CG94, FG95, MR05, Bre00, BF00, BP00, MR04, GHZ09, DGHT11].

Notational Conventions for the Stochastic Terms: For brevity we will often write e.g.

$$\|\sigma\|_{L^2} := \left(\sum_l \|\sigma_l\|_{L^2}^2 \right)^{1/2} \quad (3.5)$$

when no confusion will arise from this abuse of notation. Similarly, for $2 < p < \infty$ will also take

$$\|\sigma\|_{L^p} := \left(\int_{\mathbb{T}^2} \left(\sum_l |\sigma_l(x)|^2 \right)^{p/2} dx \right)^{1/p} \quad (3.6)$$

and, for $p = \infty$,

$$\|\sigma\|_{L^\infty} := \sup_{x \in \mathbb{T}^2} \left(\sum_l |\sigma_l(x)|^2 \right)^{1/2}. \quad (3.7)$$

We next recall some aspects of Markovian framework for (3.2). Take $\mathcal{B}(H^k)$ to be the Borelian subsets of H^k . We define the transition functions

$$P_t(\omega_0, \Gamma) = \mathbb{P}(\omega(t, \omega_0) \in \Gamma) \quad (3.8)$$

for any $t \geq 0$, $\omega_0 \in H$ and $\Gamma \in \mathcal{B}(H^k)$. Let $C_b(H^k)$ and $M_b(H^k)$ be the set of all real valued bounded continuous respectively Borel measurable functions on H^k . For $t \geq 0$, define the *Markov semigroup* according to

$$P_t \phi(\omega_0) = \mathbb{E} \phi(\omega(t, \omega_0)) = \int_{H^k} \phi(\omega) P_t(\omega_0, d\omega) \quad (3.9)$$

which maps $M_b(H^k)$ into itself. Since $\omega(t, \omega_0)$ depends continuously on $\omega_0 \in H^k$, it follows that P_t is *Feller* i.e. P_t maps $C_b(H^k)$ into itself. Let $Pr(H^k)$ be the set of Borelian probability measures on H^k . Recall that $\mu \in Pr(H^k)$ is an *invariant measure* for (3.2) if

$$\int_{H^k} \phi(\omega_0) d\mu(\omega_0) = \int_{H^k} P_t \phi(\omega_0) d\mu(\omega_0), \quad \text{for every } t \geq 0. \quad (3.10)$$

For further generalities of the ergodic theory of stochastic partial differential equations see e.g. [DPZ96, KS12].

3.2. Existence and Uniqueness of invariant measures for SNSE. For each $\nu > 0$, there exists an invariant measure μ_ν in H^0 for (3.2). This can be established using the classical Kryloff-Bogoliouboff procedure, [KB37] by proving the tightness in H^0 of a sequence of time average measures starting from any convenient initial condition. Note that if (3.4) holds then it is not hard to show that μ_ν is supported on H^k , [KS12]. We will denote by $\omega_S^\nu(\cdot)$ a statistically stationary solution of (3.2) associated to μ_ν . In other words $\mu_\nu(\cdot) = \mathbb{P}(\omega_S^\nu(t) \in \cdot)$.

The invariant measures μ_ν along with the associated stationary solutions ω_S^ν satisfy the balance relation:

$$\mathbb{E}\|\omega_S^\nu\|_{L^2}^2 = \int \|\omega_0\|_{L^2}^2 d\mu_\nu(\omega_0) = \frac{1}{2} \sum_k \|\rho_k\|_{L^2}^2 \quad (3.11)$$

and

$$\mathbb{E}\|\omega_S^\nu\|_{H^1}^2 = \int \|\omega_0\|_{H^1}^2 d\mu_\nu(\omega_0) = \frac{1}{2} \sum_k \|\sigma_k\|_{L^2}^2. \quad (3.12)$$

We derive (3.11), (3.12) by applying the Itô formula to, respectively to (3.1), (3.2) for ω_S^ν . For example,

$$\mathbb{E}\|\omega(t, \omega_0)\|_{L^2}^2 + \nu \mathbb{E} \int_0^T \|\omega(s, \omega_0)\|_{H^1}^2 ds = \mathbb{E}\|\omega_0\|_{L^2}^2 + \nu T \|\sigma\|_{L^2}^2 \quad (3.13)$$

One can also use the Itô formula to prove that

$$\mathbb{E} \exp(\delta \|\omega_S^\nu\|_{L^2}^2) \leq C < \infty \quad (3.14)$$

for some $\delta > 0$ and a constant C that is independent of ν . See e.g. [KS12] and containing references. Note that

$$\mathbb{E}\|\omega_S^\nu\|_{H^{k+1}}^2 = \int \|\omega_0\|_{H^{k+1}}^2 d\mu_\nu(\omega_0) \leq C(\nu) \quad (3.15)$$

where $C(\nu)$ is finite. However, it is doubtful that we can bound this quantity $C(\nu)$ independently of ν for $k \geq 1$.

Remark 3.1 (Uniqueness of μ_ν for $\nu > 0$). For each $\nu > 0$ the uniqueness of μ_ν is a much deeper question and requires the imposition of much specific conditions on σ . One needs to establish smoothing properties of the Markov semigroup P_t (ellipticity or hypoellipticity of the Kolmogorov equation) and that a common state can be reached by the dynamics regardless of initial conditions (irreducibility). See, e.g. [FM95, DPZ96, Mat99, Mat02, BKL01, KS01, KS02, Mat03, MP06, HM06, Kup10, HM11, Deb11, KS12]. Since the results we develop here related to inviscid limits do not require μ_ν to be unique, we do not impose such additional conditions on σ .

Remark 3.2 (Some explicit stationary solutions). We can identify some very special choices for σ which allow us to obtain explicit statistically stationary solutions of (3.2). Suppose we have found any $\omega_E : \mathbb{T}^2 \rightarrow \mathbb{R}$ satisfying

$$\mathbf{u}_E \cdot \nabla \omega_E = 0, \quad -\Delta \omega_E = \lambda \omega_E. \quad (3.16)$$

Here $\lambda > 0$ and \mathbf{u}_E is obtained from ω_E via the Biot-Savart law. For example “laminar states” satisfy (3.16). Consider the process

$$\omega_S^\nu(t, x) = \omega_E(x) \sqrt{\nu} \int_{-\infty}^t \exp(-\nu \lambda(t-s)) dW_s^1. \quad (3.17)$$

Let us note that $\omega_S^\nu := \omega_E(x) z_S(t)$, where z_S is the unique stationary solution of the 1d Ornstein-Uhlenbeck (Langevin) process $dz + \nu \lambda z dt = \sqrt{\nu} dW^1$. Here, z_S is normally distributed with mean zero and variance $(2\lambda)^{-1}$, for each $\nu > 0$. Then ω_S^ν is a stationary solution of

$$d\omega + (\mathbf{u} \cdot \nabla \omega - \nu \Delta \omega) dt = \sqrt{\nu} \omega_E dW^1. \quad (3.18)$$

This may be checked, for example by using the mild formulation of (3.18). Hence, in this setting the invariant measure obtained as $\nu \rightarrow 0$ is also normally distributed around ω_E .

3.3. Previously established properties of Kuksin measures. The balance relation (3.12) implies that any collection of invariant measures $\mathcal{S} = \{\mu_\nu\}_{\nu>0}$ is tight and therefore weakly compact. We denote by μ_0 a limiting point of \mathcal{S} , and refer to these measures as *Kuksin measures*. Let us now recall some known properties of the measures μ_0 . We refer to [KS12] for the proofs of all the facts described in this subsection.

Define $\mathcal{K} = W^{1,1}(\mathbb{R}; H^0) \cap L^2_{loc}(\mathbb{R}, H^1)$ and let \mathcal{K}_E be the set of $\omega \in \mathcal{K}$ that satisfy the Euler equation

$$\partial_t \omega + \mathbf{u} \cdot \nabla \omega = 0 \quad (3.19)$$

in its vorticity form weakly for all $t \in \mathbb{R}$. Moreover if $\omega_1, \omega_2 \in \mathcal{K}_E$ and $\omega_1(t) = \omega_2(t)$ for some $t \in \mathbb{R}$ then $\omega_1 = \omega_2$. This follows from the methods of Yudovich (see [MB02]). Define $\pi : \mathcal{K} \rightarrow H$ by $\pi(\omega) = \omega(0)$, the fiber at $t = 0$. Note that, from uniqueness, it follows that π is injective on \mathcal{K} and let

$$\mathcal{X} = \pi(\mathcal{K}_E).$$

It then holds that $\mu_0(\mathcal{X}) = 1$.

The Euler equation is well defined as a dynamical system on \mathcal{X} . Indeed, for $\omega_0 \in \mathcal{X}$, there exists a unique solution $\omega(\cdot, \omega_0) \in \mathcal{K} \subset C(\mathbb{R}, H^0) \cap L^2_{loc}(\mathbb{R}, H^1)$. Indeed for $t \in \mathbb{R}$ define $S_t : \mathcal{X} \rightarrow \mathcal{X}$, via $S_t \omega_0 = \omega(t, \omega_0)$. We endow \mathcal{X} with the topology inherited from \mathcal{K} , i.e. we take

$$d_{\mathcal{X}}(\omega_0^1, \omega_0^2) = \sum_{N \geq 1} 2^{-N} \frac{\mathcal{E}_N(\omega_0^1, \omega_0^2)}{1 + \mathcal{E}_N(\omega_0^1, \omega_0^2)} \quad (3.20)$$

where

$$\mathcal{E}_N(\omega_0^1, \omega_0^2) = \sup_{t \in [-N, N]} \|\omega(\cdot, \omega_0^1) - \omega(\cdot, \omega_0^2)\|_0^2 + \int_{-N}^N \|\omega(\cdot, \omega_0^1) - \omega(\cdot, \omega_0^2)\|_1^2 dt. \quad (3.21)$$

We then have that $\{S_t\}_{t \in \mathbb{R}}$ is a group of homeomorphisms on \mathcal{X} . Moreover μ_0 is invariant for $\{S_t\}_{t \geq 0}$, i.e. $\mu_0(E) = \mu_0(S^{-1}(t)E) = \mu_0(S(-t)E)$, for all $t \in \mathbb{R}$ and any $E \in \mathcal{B}(\mathcal{X})$.

Using local time techniques it may be shown that μ_0 is “non-trivial” in the sense that it contains no atoms. In other words, for any $\omega \in \mathcal{X}$, we have that $\mu_0(\{\omega\}) = 0$. Further properties such as spacial homogeneity, and higher moment bounds, and pointwise in space moment bounds for the measures μ_0 are discussed in [KS12].

Remark 3.3 (Vortex patches and μ_0). Let us observe that there are no vortex patch solutions in the support of μ_0 . Indeed, for any open set $O \subset \mathbb{R}^2$ let χ_O be the indicator function on O . Define

$$\mathcal{P} = \{\chi_O : O \subset \subset \mathbb{T}^2, \text{ bounded, simply connected with smooth boundary } \partial O\}.$$

For any $\omega_0 \in \mathcal{P}$, according to e.g. [MB02] there exists $t_0 > 0$ such that $\omega(t, \omega_0) \in \mathcal{P}$, for all $t \in [-t_0, t_0]$. As such, since $H^1 \cap \mathcal{P} = \emptyset$, for any $\omega_0 \in \mathcal{P}$, $\omega^0(\cdot, \omega_0) \notin L^2_{loc}(\mathbb{R}, \dot{H}^1)$ and hence $\omega^0(\cdot, \omega_0) \notin \mathcal{K}_E$. Thus $\mu_0(\mathcal{P}) = 0$ since

$$\mathcal{P} \cap \mathcal{X} = \emptyset. \quad (3.22)$$

4. Invariant Measure Supported on L^∞ and Related Estimates

In this section we establish uniform in ν bounds on $\mathbb{E}\|\omega_\nu^S\|_{L^\infty}$, where ω_ν^S are stationary solutions of (3.2). Our approach makes use of the Moser iteration technique and draws on earlier works in this direction in [DMS05, DMS09]. However we obtain parabolic regularization and time decay for the initial data component, which was not addressed in the above works. Let us note that in the deterministic case L^∞ bounds may be obtained without appealing to the Moser iteration: one carries out L^p estimates, which take advantage of cancellation in the nonlinearity, and then sends $p \rightarrow \infty$. In the stochastic case, the Itô correction terms arise and cause the bounds on the L^p norm to grow unboundedly as $p \rightarrow \infty$.

Theorem 4.1 (Stochastic Moser). *For $\nu > 0$ consider an invariant measure μ_ν and an associated stationary solution ω_ν^S of (3.2), where the stochastic forcing is assumed to be sufficiently smooth, e.g. (3.4) holds. Then the following bound holds*

$$\mathbb{E}\|\omega_\nu^S\|_{L^\infty} \leq C < \infty \quad (4.1)$$

where $C = C(\sigma)$ is independent on ν .

An immediate consequence of estimate (4.1) is that any limit point μ_0 of any sequence of invariant measures $\{\mu_\nu\}_{\nu>0}$ is concentrated on $L^\infty(\mathbb{T}^2)$.

Theorem 4.2 (Invariant measure supported on L^∞). *Under the assumptions of Theorem 4.1, consider any collection of invariant measures $\{\mu_\nu\}_{\nu>0}$ of (3.2). Then there exists a subsequence and a measure μ_0 such that $\mu_{\nu_j} \rightharpoonup \mu_0$ (weakly) in $\text{Pr}(H^0)$ as $j \rightarrow \infty$ and $\mu_0(L^\infty) = 1$.*

Remark 4.3. Let us note that μ_0 is an invariant measure for the Euler equation over $L^\infty \cap X$, where X is the fiber at $t = 0$ of \mathcal{X}_E . See Section 3 for details.

We shall first give the proof of Theorem 4.2, assuming Theorem 4.1 holds, and then return and prove Theorem 4.1.

PROOF OF THEOREM 4.2. As in [Kuk04] by using (3.12) we have that $\{\mu_\nu\}$ is tight, and hence weakly compact on $\text{Pr}(H^0)$. Taking μ_0 to be a limit point of $\{\mu_\nu\}_{\nu>0}$ in the weak topology of H^0 , there exists a sequence $\nu_j \rightarrow 0$ such that

$$\lim_{j \rightarrow \infty} \int \phi(\omega_0) d\mu_{\nu_j}(\omega_0) = \int \phi(\omega_0) d\mu_0(\omega_0)$$

for each $\phi \in C_b(H^0)$.

According to Theorem 4.1 we have

$$\sup_{\nu>0} \mathbb{E}\|\omega_\nu^S\|_{L^\infty} = \sup_{\nu>0} \int \|\omega_0\|_{L^\infty} d\mu_\nu(\omega_0) \leq C < \infty. \quad (4.2)$$

We claim that this implies

$$\int \|\omega_0\|_{L^\infty} d\mu_0(\omega_0) \leq C. \quad (4.3)$$

Indeed, take ρ_ε to be a standard family of smooth mollifiers on \mathbb{R}^2 . For $R > 0$, $\varepsilon > 0$ define $\phi_{R,\varepsilon}(\omega) = \|\rho_\varepsilon * \omega\|_{L^\infty} \wedge R$. Young's inequality implies the $\phi_{R,\varepsilon} \in C_b(H^0)$ so that

$$\int \phi_{R,\varepsilon}(\omega_0) d\mu_0(\omega_0) \leq C.$$

Now, by Fatou's Lemma, we have

$$\int (\liminf_{\varepsilon>0} \|\rho_\varepsilon * \omega_0\|_{L^\infty} \wedge R) d\mu_0(\omega_0) \leq C.$$

Since $\|\omega_0\|_{L^\infty} \leq \liminf_{\varepsilon>0} \|\rho_\varepsilon * \omega_0\|_{L^\infty}$ for each $\omega_0 \in L^\infty$ then (4.3) follows, completing the proof. \square

PROOF OF THEOREM 4.1. As a first step we rescale the time in (3.2). Taking $\tilde{t} = t/\nu$ we obtain

$$d\tilde{\omega} + \left(\frac{1}{\nu} \tilde{\mathbf{u}} \cdot \nabla \tilde{\omega} - \Delta \tilde{\omega} \right) dt = \sigma d\tilde{W}, \quad \tilde{\omega}(0) = \omega_S(0), \quad (4.4)$$

where we have denoted $\tilde{\omega}(t, x) = \omega(\tilde{t}, x)$, $\tilde{\mathbf{u}}(t, x) = \mathbf{u}(\tilde{t}, x)$, and $\tilde{W}(t) = \sqrt{\nu}W(\tilde{t})$. Note that $\tilde{W}(t)$ has the same statistical properties as $W(t)$. For ease of notation for we drop the tildes until (4.41) below.

Fix $T > 0$, $\rho > 1$, and define T_k to be an increasing sequence of times with $T_0 = 0$ and T_k as $k \rightarrow \infty$. Let $I_k = [T_k, 2T]$ be a sequence of time intervals approaching $[T, 2T]$.

To analyze (4.4) we apply the L^p Itô Lemma. This yields

$$d\|\omega\|_{L^p}^p = \left(\frac{p}{\nu} T_{1,p} + p T_{2,p} + \frac{p(p-1)}{2} T_{3,p} \right) dt + p \sum_m S_{m,p} dW^m \quad (4.5)$$

where we have denoted

$$T_{1,p}(t) = - \int_{\mathbb{T}^2} \mathbf{u}(t, x) \cdot \nabla \omega(t, x) \omega(t, x) |\omega(t, x)|^{p-2} dx = 0$$

$$T_{2,p}(t) = \int_{\mathbb{T}^2} \Delta \omega(t, x) \omega(t, x) |\omega(t, x)|^{p-2} dx \quad (4.6)$$

$$T_{3,p}(t) = \sum_m \int_{\mathbb{T}^2} |\omega(t, x)|^{p-2} \sigma_m(x)^2 dx \quad (4.7)$$

$$S_{m,p}(t) = \int_{\mathbb{T}^2} \sigma_m(x) \omega(t, x) |\omega(t, x)|^{p-2} dx \quad (4.8)$$

for all $p \geq 2$. In the identity for $T_{1,p}$ we have integrated by parts in x and used that $\nabla \cdot \mathbf{u} = 0$. Since $\nu > 0$, we are dealing with spatially smooth solutions of (4.4), and hence the identity (4.5) may be justified by applying the Itô lemma pointwise in x , integrating over the torus, and using the stochastic Fubini theorem (see, e.g. [DPZ92]). Note however that (4.5) may be justified for much less spatially regular stochastic evolution equations, as recently established in [Kry10].

Let $s \in [T_k, T_{k+1}]$ and $t > s$. We start with (4.5) for $p \geq 2$, integrated from s to t :

$$\begin{aligned} \|\omega(t)\|_{L^p}^p - p \int_s^t T_{2,p}(\tau) d\tau &= \|\omega(s)\|_{L^p}^p + \frac{p(p-1)}{2} \int_s^t T_{3,p}(\tau) d\tau + p \sum_m \int_s^t S_{m,p}(\tau) dW_\tau^m \\ &= \|\omega(s)\|_{L^p}^p + \frac{p(p-1)}{2} \int_s^t T_{3,p}(\tau) d\tau + p \sum_m \int_{T_k}^t S_{m,p}(\tau) dW_\tau^m \\ &\quad - p \sum_m \int_{T_k}^s S_{m,p}(\tau) dW_\tau^m \end{aligned} \quad (4.9)$$

where $T_{2,p}, T_{3,p}, S_{m,p}$ are as defined in (4.6)–(4.8). We take the supremum of (4.9) over every $t \in I_{k+1}$, and obtain

$$\begin{aligned} \|\omega\|_{L^\infty(I_{k+1}; L^p)}^p - p \int_{I_{k+1}} T_{2,p}(\tau) d\tau \\ \leq \|\omega(s)\|_{L^p}^p + \frac{p(p-1)}{2} \int_{I_k} |T_{3,p}(\tau)| d\tau + p \sup_{t \in I_{k+1}} \left| \sum_m \int_{T_k}^t S_{m,p}(\tau) dW_\tau^m \right| \\ \quad + p \left| \sum_m \int_{T_k}^s S_{m,p}(\tau) dW_\tau^m \right| \\ \leq \|\omega(s)\|_{L^p}^p + \frac{p(p-1)}{2} \int_{I_k} |T_{3,p}(\tau)| d\tau + 2p \sup_{t \in I_k} \left| \sum_m \int_{T_k}^t S_{m,p}(\tau) dW_\tau^m \right| \end{aligned} \quad (4.10)$$

where we have used that $s \in [T_k, T_{k+1})$, and that $T_{2,p} \leq 0$ (cf. (4.18) below). This allowed us to bound from below the time integration on the left side from $[s, 2T]$ with the smaller one on $[T_{k+1}, 2T]$.

For the forthcoming computations it will be convenient to introduce the following standard notations for the stochastic (martingale) terms. For any $0 \leq r \leq t$, let

$$M_{[r,t],p} = \sum_m \int_r^t S_{m,p}(\tau) dW_\tau^m \quad (4.11)$$

and denote the running absolute maximum by

$$M_{[r,t],p}^* := \sup_{s \in [r,t]} \left| \sum_m \int_r^s S_{m,p}(\tau) dW_\tau^m \right|. \quad (4.12)$$

Finally we define $\langle M_{[r,\cdot]p} \rangle_t$ to be the quadratic variation of $M_{[r,t],p}$ and recall that (see e.g. [KS91])

$$\langle M_{[r,\cdot]p} \rangle_t = \int_r^t \sum_m S_{m,p}^2 ds. \quad (4.13)$$

Recall that by a version of the Burkholder-Davis-Gundy inequality given in [DMS05] (see also [RY99]) we have that, for any non-negative random variable Z , any $r < t$ and any $\delta > 0$

$$\mathbb{E}(M_{[r,t],p}^* \vee Z)^\delta \leq C_{BDG}(\delta) \cdot \mathbb{E}(\langle M_{[r,\cdot]p} \rangle_t^{1/2} \vee Z)^\delta. \quad (4.14)$$

Here the constant $C_{BDG}(\delta)$ is universal; it depends only on δ and is independent of the form of the Martingale $M_{[r,t],p}^*$ or Z . Also, note carefully that there exists a $\delta_0 > 0$ such that

$$C_{BDG}(\delta) \leq 2^{\delta^{1/2}} \quad \text{whenever } \delta < \delta_0. \quad (4.15)$$

This observation will be crucial below in estimates (4.33) and (4.37).

We return to (4.10) and take an average of (4.10) for $s \in [T_k, T_{k+1}]$, and obtain

$$\begin{aligned} & \|\omega\|_{L^\infty(I_{k+1}; L^p)}^p - p \int_{I_{k+1}} T_{2,p}(\tau) d\tau \\ & \leq \frac{1}{T_{k+1} - T_k} \int_{T_k}^{T_{k+1}} \|\omega\|_{L^p(I_k; L^p)}^p + \frac{p(p-1)}{2} \int_{I_k} |T_{3,p}(\tau)| d\tau + 2pM_{[T_k, 2T],p}^* \\ & \leq \frac{1}{(T_{k+1} - T_k)^{1/2}} \|\omega\|_{L^{2p}(I_k; L^p)}^p + \frac{p(p-1)}{2} \int_{I_k} |T_{3,p}(\tau)| d\tau + 2pM_{[T_k, 2T],p}^*. \end{aligned} \quad (4.16)$$

As usual in Moser iteration arguments, the lower bound on $-T_{2,p}$ is obtained by introducing $v = |\omega|^{p/2}$, so that $\|\omega\|_{L^p}^p = \|v\|_{L^2}^2$, and $|\nabla v|^2 = \frac{p^2}{4} |\nabla \omega|^2 |\omega|^{p-2}$. Then, upon integrating by parts in $T_{2,p}$ we have, pointwise in time, that

$$-pT_{2,p} = p(p-1) \int_{\mathbb{T}^2} |\nabla \omega|^2 |\omega|^{p-2} dx = 4 \frac{p-1}{p} \|\nabla v\|_{L^2}^2 \geq 2 \|\nabla v\|_{L^2}^2 \quad (4.17)$$

for all $p \geq 2$. Moreover, since we are in a two dimensional periodic box, the Sobolev embedding gives

$$\frac{1}{2C_s} \|v\|_{L^{2^*}}^2 \leq \|\nabla v\|_{L^2}^2 + \|v\|_{L^2}^2 \quad (4.18)$$

where $2^* \in [2, \infty)$ is arbitrary, and the constant $C_s > 0$ depends only on the size of the box and the choice of 2^* . Note that v is not zero mean in space and hence we need to add here a lower order term in (4.18). Let us choose $2^* = 4$ for simplicity. Then, in view of (4.17) and (4.18), the left hand side of (4.16) is bounded from below as

$$\|v\|_{L^\infty(I_{k+1}; L^2)}^2 + 2\|\nabla v\|_{L^2(I_{k+1}; L^2)}^2 \geq \|v\|_{L^\infty(I_{k+1}; L^2)}^2 - 2\|v\|_{L^2(I_{k+1}; L^2)}^2 + 2\left(\|v\|_{L^2(I_{k+1}; L^2)}^2 + \|\nabla v\|_{L^2(I_{k+1}; L^2)}^2\right). \quad (4.19)$$

By assuming that

$$4|I_{k+1}| = 4(2T - T_{k+1}) \leq 1, \quad (4.20)$$

which is automatically satisfied for all $k \geq 0$ if we ensure that

$$T \leq \frac{1}{8}, \quad (4.21)$$

we conclude from (4.19) that

$$\begin{aligned} \|v\|_{L^\infty(I_{k+1}; L^2)}^2 + 2\|\nabla v\|_{L^2(I_{k+1}; L^2)}^2 & \geq \|v\|_{L^\infty(I_{k+1}; L^2)}^2 (1 - 2|I_{k+1}|) + \frac{1}{C_s} \|v\|_{L^2(I_{k+1}; L^4)}^2 \\ & \geq \frac{1}{2} \|v\|_{L^\infty(I_{k+1}; L^2)}^2 + \frac{1}{C_s} \|v\|_{L^2(I_{k+1}; L^4)}^2. \end{aligned} \quad (4.22)$$

Let us now recall the following $L_t^p L_x^q$ interpolation inequality. Suppose we have $1 \leq p_1, p_2, q_1, q_2, r_1, r_2 \leq \infty$ and $0 \leq \gamma \leq 1$ satisfy

$$\frac{1}{r_1} = \frac{\gamma}{p_1} + \frac{1-\gamma}{q_1}, \quad \frac{1}{r_2} = \frac{\gamma}{p_2} + \frac{1-\gamma}{q_2}. \quad (4.23)$$

Then, for any $g \in L^{p_1}(I; L^{p_2}) \cap L^{q_1}(I; L^{q_2})$ we have

$$\|g\|_{L^{r_1}(I; L^{r_2})} \leq \|g\|_{L^{p_1}(I; L^{p_2})}^\gamma \|g\|_{L^{q_1}(I; L^{q_2})}^{1-\gamma} \quad (4.24)$$

with $I \subset \mathbb{R}$ being some interval. Taking $r_1 = 5$, $r_2 = 5/2$, $p_1 = \infty$, $p_2 = 2$, $q_1 = 2$, $q_2 = 4$, and $\gamma = 3/5$ in this inequality we find

$$\frac{1}{2C_S^{2/5}} \|v\|_{L^5(I_{k+1}; L^{5/2})}^2 \leq \frac{1}{2C_S^{2/5}} \|v\|_{L^\infty(I_{k+1}; L^2)}^{6/5} \|v\|_{L^2(I_{k+1}; L^4)}^{4/5} \leq \frac{1}{2} \|v\|_{L^\infty(I_{k+1}; L^2)}^2 + \frac{1}{C_S} \|v\|_{L^2(I_{k+1}; L^4)}^2 \quad (4.25)$$

by making use of the ε -Young inequality. In summary, we have shown that the left hand side of (4.16) is bounded from below by

$$\frac{1}{C'_S} \|v\|_{L^5(I_{k+1}; L^{5/2})}^2 = \frac{1}{C'_S} \|\omega\|_{L^{5p/2}(I_{k+1}; L^{5p/4})}^p = \frac{1}{C'_S} \|\omega\|_{L^{2\lambda p}(I_{k+1}; L^{\lambda p})}^p \quad (4.26)$$

as long as (4.20) holds and for any $p \geq 2$. Here we have denoted $C'_S = 2C_S^{2/5} \vee 1$, and denoted

$$\lambda = \frac{5}{4}. \quad (4.27)$$

For the term $T_{3,p}$ -term on the left side of (4.16) we simply use Hölder and obtain (pointwise in time)

$$\frac{p(p-1)}{2} T_{3,p} \leq \frac{p(p-1)}{2} \|\omega\|_{L^p}^{p-2} \|\sigma\|_{L^p}^2.$$

Integrating the above on I_k and using the Hölder inequality in time we obtain

$$\frac{p(p-1)}{2} \int_{I_k} T_{3,p}(\tau) d\tau \leq \frac{p(p-1)}{2} \|\sigma\|_{L^p}^2 |I_k|^{\frac{p+2}{2p}} \|\omega\|_{L^{2p}(I_k; L^p)}^{p-2} \quad (4.28)$$

for all $k \geq 0$. Thus from (4.16), (4.26), and (4.28), we obtain

$$\begin{aligned} \frac{1}{C'_S} \left(\|\omega\|_{L^{2\lambda p}(I_{k+1}; L^{\lambda p})} \vee \|\sigma\|_{L^\infty} \right)^p &\leq \frac{1}{C'_S} \left(\|\omega\|_{L^{2\lambda p}(I_{k+1}; L^{\lambda p})}^p + \|\sigma\|_{L^\infty}^p \right) \\ &\leq \frac{1}{(T_{k+1} - T_k)^{1/2}} \|\omega\|_{L^{2p}(I_k; L^p)}^p + |\mathbb{T}^2|^{2/p} p^2 |I_k|^{\frac{p+2}{2p}} \|\sigma\|_{L^\infty}^2 \|\omega\|_{L^{2p}(I_k; L^p)}^{p-2} \\ &\quad + \|\sigma\|_{L^\infty}^p + 2p M_{[T_k, 2T], p}^* \end{aligned} \quad (4.29)$$

with $M_{[T_k, 2T], p}^*$ as defined in (4.12), and we have used that $\|\sigma\|_{L^p} \leq |\mathbb{T}^2|^{1/p} \|\sigma\|_{L^\infty}$.

Let us now define

$$\kappa(p, T) := 4C'_S \left(\frac{1}{(T_{k+1} - T_k)^{1/2}} + |\mathbb{T}^2|^{2/p} p^2 |I_k|^{\frac{p+2}{2p}} + 1 + 2p |\mathbb{T}^2|^{1/p} |I_k|^{\frac{1}{2p}} \right) \quad (4.30)$$

After some direct manipulations starting from (4.29), taking p^{th} roots of both sides and then expectations we find that

$$\begin{aligned} &\mathbb{E} \left(\|\omega\|_{L^{2\lambda p}(I_{k+1}; L^{\lambda p})} \vee \|\sigma\|_{L^\infty} \right) \\ &\leq \kappa(p, T)^{\frac{1}{p}} \mathbb{E} \left(\left(\|\omega\|_{L^{2p}(I_k; L^p)} \vee \|\sigma\|_{L^\infty} \right)^p \vee |\mathbb{T}^2|^{-\frac{1}{p}} |I_k|^{-\frac{1}{2p}} M_{[T_k, 2T], p}^* \right)^{\frac{1}{p}} \\ &\leq \kappa(p, T)^{\frac{1}{p}} C_{BDG}(p^{-1}) \mathbb{E} \left(\left(\|\omega\|_{L^{2p}(I_k; L^p)} \vee \|\sigma\|_{L^\infty} \right)^p \vee |\mathbb{T}^2|^{-\frac{1}{p}} |I_k|^{-\frac{1}{2p}} \langle M_{[T_k, \cdot], p} \rangle_{2T}^{1/2} \right)^{\frac{1}{p}} \end{aligned} \quad (4.31)$$

which holds for all $p \geq 2$. Note that for the second inequality we used (4.14).

We next estimate the quadratic variation term, $\langle M_{[T_k, \cdot], p} \rangle_{2T}^{1/2}$ in (4.31). Starting from (4.8) and (4.13) we find

$$\begin{aligned} \langle M_{[T_k, \cdot], p} \rangle_{2T}^{1/2} &= \left(\int_{T_k}^{2T} \sum_m S_{m,p}^2 dt \right)^{\frac{1}{2}} \leq \left(\int_{I_k} \left(\int_{\mathbb{T}^2} \left(\sum_m \sigma_m^2 \right)^{1/2} |\omega|^{p-1} dx \right)^2 dt \right)^{\frac{1}{2}} \\ &\leq \left(\int_{I_k} \|\sigma\|_{L^p}^2 \|\omega\|_{L^p}^{2(p-1)} dt \right)^{\frac{1}{2}} \leq |I_k|^{\frac{1}{2p}} \|\sigma\|_{L^p} \|\omega\|_{L^{2p}(I_k; L^p)}^{p-1} \\ &\leq |\mathbb{T}^2|^{\frac{1}{p}} |I_k|^{\frac{1}{2p}} \|\sigma\|_{L^\infty} \|\omega\|_{L^{2p}(I_k; L^p)}^{p-1} \leq |\mathbb{T}^2|^{\frac{1}{p}} |I_k|^{\frac{1}{2p}} \left(\|\sigma\|_{L^\infty} \vee \|\omega\|_{L^{2p}(I_k; L^p)} \right)^p. \end{aligned} \quad (4.32)$$

Note that the second bound above makes use of the integral Minkowski inequality.

Let us now summarize the estimates obtained, by combining (4.31) with (4.32). We have

$$\mathbb{E} \left(\|\omega\|_{L^{2\lambda p}(I_{k+1}; L^{\lambda p})} \vee \|\sigma\|_{L^\infty} \right) \leq \kappa(p, T)^{\frac{1}{p}} C_{BDG}(p^{-1}) \mathbb{E} \left(\|\omega\|_{L^{2p}(I_k; L^p)} \vee \|\sigma\|_{L^\infty} \right) \quad (4.33)$$

for all $p \geq 2$. To set up a recurrence relation, it is hence natural to set $p = p_k$ in (4.33), where we define

$$p_k = 2\lambda^k$$

for all $k \geq 0$, where we recall that $\lambda = 5/4$. Let us now introduce some notation

$$A_k = \mathbb{E} \left(\|\omega\|_{L^{2p_k}(I_k, L^{p_k})} \vee \|\sigma\|_{L^\infty} \right) \quad (4.34)$$

$$a_k = \kappa(p_k, T)^{\frac{1}{p_k}} C_{BDG}(p_k^{-1}). \quad (4.35)$$

Then, (4.33) reads

$$A_{k+1} \leq a_k A_k. \quad (4.36)$$

So that

$$\mathbb{E} \sup_{t \in [T, 2T]} \|\omega(t, \cdot)\|_{L^\infty} \leq A_\infty \leq \left(\prod_{k \geq 0} a_k \right) \mathbb{E} \left(\|\omega\|_{L^4([0, 2T]; L^2)} \vee \|\sigma\|_{L^\infty} \right). \quad (4.37)$$

In view of (4.15), we have that

$$\prod_{k \geq 0} a_k \leq C \exp \left(\sum_{k \geq 0} \frac{\log \kappa(p_k, T)}{p_k} \right) \quad (4.38)$$

where C is a ν - and T -independent constant. We now set

$$T_k = T(1 - \lambda^{-k}).$$

Then, $T_{k+1} - T_k = T\lambda^{-k}(1 + \lambda^{-1}) \geq T\lambda^{-k}/2 = Tp_k^{-1}$. We recall the definition of $\kappa(p_k, T)$ from (4.30), which in view of the above choices may be bounded as

$$\kappa(p_k, T) \leq 4C'_S \left(T^{-\frac{1}{2}} p_k^{\frac{1}{2}} + |\mathbb{T}^2|^{\frac{2}{p_k}} p_k^2 (2T)^{\frac{p_k+2}{2p_k}} + 1 + 2p_k |\mathbb{T}^2|^{\frac{1}{p_k}} (2T)^{\frac{1}{2p_k}} \right) \leq Cp_k^2 \left(T^{-\frac{1}{2}} + 1 \right) \quad (4.39)$$

where we have also used that $|I_k| \leq 2T \leq 1/4$ (cf. (4.21)), and C is a sufficiently large T -independent constant. Using that $\sum_{k \geq 0} p_k^{-1} = 5/2$, and $\sum_{k \geq 0} p_k^{-1} \log p_k < \infty$ we may further obtain that

$$\prod_{k \geq 0} a_k \leq C(T^{-1/2} + 1)^{5/2} \leq C(T^{-5/4} + 1) \quad (4.40)$$

for some sufficiently large ν - and T -independent constant C .

In summary from (4.37), (4.40) and recalling that these estimates were carried out for the rescaled equation (4.4) above we have in conclusion

$$\mathbb{E} \sup_{t \in [T, 2T]} \|\tilde{\omega}(t)\|_{L^\infty} \leq C(T^{-5/4} + 1) \mathbb{E} \left(\|\tilde{\omega}\|_{L^4([0, 2T]; L^2)} \vee \|\sigma\|_{L^\infty} \right), \quad (4.41)$$

for any $T \leq 1/8$ and where C is independent of T and ν . Rescaling to the original variable $\omega(t) = \tilde{\omega}(\nu t)$ then with (4.41) we infer

$$\mathbb{E} \sup_{t \in [T/\nu, 2T/\nu]} \|\omega(t)\|_{L^\infty} \leq C(T^{-5/4} + 1) \left(\left(\int_0^{T/\nu} \nu \mathbb{E}(\|\omega(s)\|_{L^2}^4) ds \right)^{1/4} + \|\sigma\|_{L^\infty} \right) \quad (4.42)$$

for any $T \leq 1/8$.

We can now obtain the desired conclusion by taking ω to be ω_S^ν a stationary solution of (3.2) corresponding to μ_ν . Recalling (3.14) and taking, for example $T = 1/8$ we have that

$$\begin{aligned} \mathbb{E} \|\omega_S^\nu\|_{L^\infty} &\leq \mathbb{E} \sup_{t \in [1/(8\nu), 1/(4\nu)]} \|\omega_S^\nu(t)\|_{L^\infty} \\ &\leq C \left(\left(\int_0^{1/(8\nu)} \nu \mathbb{E}(\|\omega_S^\nu(s)\|_{L^2}^4) ds \right)^{1/4} + \|\sigma\|_{L^\infty} \right) \\ &\leq C \left(\left(\int_0^{1/(8\nu)} \frac{2\nu}{\delta^2} \mathbb{E} \exp(\delta \|\omega_S^\nu(s)\|_{L^2}^2) ds \right)^{1/4} + \|\sigma\|_{L^\infty} \right) \leq C_\sigma \end{aligned} \quad (4.43)$$

for a constant C_σ independent of ν . This gives (4.1) concluding the proof of Theorem 4.1. \square

Remark 4.4 (Linear drift-diffusion). Note that the L^∞ bounds obtained in Theorem 4.1 can also be shown to hold for any drift-diffusion equation

$$d\omega + (\mathbf{a} \cdot \nabla \omega - \Delta \omega) dt = \sigma dW, \quad \nabla \cdot \mathbf{a} = 0,$$

with sufficiently regular drift \mathbf{a} and stochastic forcing σ . Indeed as in (4.5) one may write the evolution of the L^p -norm of ω with the analogous drift-term vanishing since \mathbf{a} is divergence free. The rest of the proof follows without any change and one obtains that

$$\mathbb{E} \sup_{t \in [T, 2T]} \|\omega(t)\|_{L^\infty} \leq C(1 + T^{-5/4}) \mathbb{E}(\|\omega\|_{L^4([0, 2T]; L^2)} \vee \|\sigma\|_{L^\infty}),$$

for any $0 < T \leq 1/8$ and most importantly C is independent \mathbf{a} . Note also that this estimate corresponds to the usual parabolic regularization in the deterministic case: L^2 weak-solutions are instantaneously in L^∞ .

Remark 4.5 (Fractional Navier-Stokes). Note that the Moser iteration technique used to prove Theorem 4.1 may be used to obtain drift-independent L^∞ bounds for stationary solutions of the fractional drift-diffusion equation

$$d\omega + (\mathbf{a} \cdot \nabla \omega + (-\Delta)^{\gamma/2} \omega) dt = \sigma dW,$$

for any power $\gamma \in (0, 2)$, where as in Remark 4.4 the drift \mathbf{a} is divergence-free and sufficiently smooth. To see this, we recall the L^p lower bound on the fractional Laplacian given in [CC04]

$$p \int \omega |\omega|^{p-2} (-\Delta)^{\gamma/2} \omega dx \geq \int |(-\Delta)^{\gamma/4} (|\omega|^{p/2})|^2 dx$$

which holds for any $p \geq 2$. Using the 2D Sobolev embedding $H^{\gamma/2} \subset L^{4/(2-\gamma)}$, one may repeat the argument given above in (4.17)–(4.27), and obtain estimate (4.26) with $\lambda_\gamma = 1 + \gamma/4$. Since for any $\gamma \in (0, 2)$ we have $\lambda_\gamma^n \rightarrow \infty$ as $n \rightarrow \infty$ the Moser iteration scheme may be completed mutatis-mutandis. In particular, setting $\mathbf{a} = \frac{1}{\nu} \mathbf{u}$, which is divergence-free, in view of (4.4) one may use the above argument to study inviscid limits of the stochastic fractionally-dissipative Navier-Stokes equation.

5. Modulus of continuity for the deterministic stationary problem

In Section 4 we have proven that the stationary solution ω_S^ν of (4.4) obeys ν -independent bounds in L^∞ , that is $\mathbb{E} \|\omega_S^\nu\|_{L^\infty}$ is uniformly bounded in ν . The key ingredients used in this argument were

- *Two-dimensionality*: this ensures that the nonlinear term, whose size blows up (in comparison to the viscosity) as $\nu \rightarrow 0$, vanishes altogether in L^p estimates for the vorticity. To put it differently, there is no vorticity stretching term.

- *Stationarity*: this enables us to measure the L^∞ norm of the solution whose initial data is ω_S^ν at time $T_\nu \approx \nu^{-1}$, and hence obtain bounds on ω_S^ν itself.

Once we wish to estimate ω_S^ν in more regular spaces, for example H^s with $s > 1$, or C^γ with $\gamma > 0$, the nonlinear term does not vanish anymore, and since it's relative size becomes prohibitively large as $\nu \rightarrow 0$, we do not seem to be able to obtain ν -independent bounds on ω_S^ν , in spaces that are better than L^∞ (averaged over the probability space).

In this section we exhibit an *drift-independent* bound, in a better norm than L^∞ , of solutions to the stationary drift-diffusion equation

$$Lv = -\Delta v + b \cdot \nabla v = f \quad (5.1)$$

for $x \in \mathbb{T}^2$, where $b = b(x)$ is a divergence free-vector field, but on which we have *no bounds*. The force is assumed to be in L^∞ , with zero-mean, and we consider solutions v such that $\int_{\mathbb{T}^2} v dx = 0$. We view equation (5.1) as a *deterministic* toy-model describing the *stationary* solutions of (4.4) – the analogy is given by letting $b = \nu^{-1} \mathbf{u}$, $v = \omega$, and noting that $\omega(0)$ equals $\omega(t)$ in law, for all $t \geq 0$.

Following the ideas in [SSŠZ12] in the spirit of [Leb07] we show that v obeys a logarithmic modulus of continuity which does not depend on the size of the drift b . In particular v is a uniformly continuous function.

Theorem 5.1 (Modulus of continuity for deterministic stationary equation). *Let b be divergence free and smooth, and v be a zero mean weak solution of (5.1), that is, $v \in H^1$ and satisfies (5.1) in the sense of distributions. Then v obeys a drift-independent logarithmic modulus of continuity*

$$\sup_{|x-y| \leq r} |v(x) - v(y)| \leq \frac{C(\|f\|_{L^\infty})}{\sqrt{\log 1/r}} \quad (5.2)$$

for some $C > 0$ that is independent of b and all $r \in (0, r_*]$, for some universal constant r_* . In particular, v is uniformly continuous.

PROOF OF THEOREM 5.1. Taking the inner product of (5.1), using the Poincaré inequality and the fact that $\nabla \cdot b = 0$, we obtain

$$\frac{1}{C} \|v\|_{L^2}^2 + \frac{1}{2} \|\nabla v\|_{L^2}^2 \leq \|f\|_{L^2} \|v\|_{L^2}$$

which implies

$$\|\nabla v\|_{L^2}^2 \leq C_1 \|f\|_{L^2}^2 \quad (5.3)$$

for some $C_1 > 0$ that is independent of b . We as usual denote $\text{osc}_K v = \max_K v - \min_K v$ to be the oscillation of v over the set K . Fix $x_0 = 0$, $r_0 = \text{diam}(\mathbb{T}^2)/4 \wedge 1/2$, and let $B_r = B_r(x_0)$ for any $r \leq r_0$. Upon integrating in polar coordinates, dropping the normal derivatives, and using that by the 1D Sobolev embedding $H^1(\partial B_r) \subset C^\alpha(\partial B_r)$ for $\alpha \in (0, 1/2)$ (see [SSŠZ12, Theorem 4.2]), we obtain

$$C_1 \|f\|_{L^2}^2 \geq \int_r^{\sqrt{r}} \frac{(\text{osc}_{\partial B_\rho} v)^2}{\rho} d\rho, \quad (5.4)$$

for any $r \in (0, r_0^2]$.

If we were able to establish that v is monotone in the sense of Lebesgue, i.e. to show that $\text{osc}_{\partial B_\rho} v$ is a monotone function of ρ , the proof of the lemma would directly follow from (5.4). Instead we prove that v is *almost* monotone in the sense of Lebesgue, that is, up to an error of size r^2 . Let h solve

$$Lh = -f \text{ in } B_r, \quad h = 0 \text{ on } \partial B_r$$

so that $L(v+h) = 0$ in B_r , and hence by the maximum principle

$$\text{osc}_{B_r}(v+h) = \text{osc}_{\partial B_r}(v+h) = \text{osc}_{\partial B_r} v. \quad (5.5)$$

We claim that

$$\text{osc}_{B_r} h \leq C \|f\|_{L^\infty(B_r)} r^2 \quad (5.6)$$

for some constant $C > 0$ that is independent of b . To prove (5.6), we rescale the problem to the unit ball by letting

$$x = ry, \tilde{h}(y) = h(ry), \tilde{b}(y) = b(x), \tilde{f}(y) = f(x).$$

It follows that

$$-\Delta_y \tilde{h} + r \tilde{b} \cdot \nabla_y \tilde{h} = -r^2 \tilde{f} \text{ in } B_1, \quad \tilde{h} = 0 \text{ on } \partial B_1. \quad (5.7)$$

We obtain the desired estimate by Moser iteration. Multiplying (5.7) by $\tilde{h}|\tilde{h}|^{p-2}$ and integrating over B_1 we obtain that for any $p \geq 2$

$$\frac{4(p-1)}{p^2} \int_{B_1} |\nabla(|\tilde{h}|^{p/2})|^2 dx = (p-1) \int_{B_1} |\nabla \tilde{h}|^2 |\tilde{h}|^{p-2} dx \leq r^2 |B_1|^{1/p} \|\tilde{f}\|_{L^\infty} \|\tilde{h}\|_{L^p}^{p-1}$$

by using that $\nabla \cdot \tilde{b} = 0$, and that \tilde{h} vanishes on ∂B_1 . Using the Sobolev embedding $H^1 \subset L^4$ in 2D, we get

$$\|\tilde{h}\|_{L^{2p}}^p \leq Cp |B_1|^{1/p} r^2 \|\tilde{f}\|_{L^\infty} \|\tilde{h}\|_{L^p}^{p-1} \quad (5.8)$$

for some $C > 0$ which is independent of p . Without loss of generality we take this constant C sufficiently large so that $2C|B_1|^{1/p} \geq 1$ for any $p \geq 2$. Let

$$p_k = 2^k \text{ and } a_k = \max\{\|\tilde{h}\|_{L^{p_k}}, r^2 \|\tilde{f}\|_{L^\infty}\}.$$

It follows from (5.8) that

$$a_{k+1} \leq (Cp_k |B_1|^{1/p_k})^{1/p_k} a_k$$

and thus

$$\|\tilde{h}\|_{L^\infty(B_1)} \leq C (r^2 \|\tilde{f}\|_{L^\infty(B_1)} + \|\tilde{h}\|_{L^2(B_1)}) \leq Cr^2 \|\tilde{f}\|_{L^\infty(B_1)} \quad (5.9)$$

for some $C > 0$. The last inequality above follows by setting $p = 2$ in (5.8), and using the Hölder inequality. Upon rescaling back to x -variables, (5.9) implies (5.6).

Combining (5.5) with (5.6) gives that for any $r \leq \rho < r_0$ we have

$$\text{osc}_{\partial B_\rho} v \geq \text{osc}_{B_\rho} v - C \|f\|_{L^\infty} \rho^2 \geq \text{osc}_{B_r} v - C_2 \|f\|_{L^\infty} \rho^2 \quad (5.10)$$

for some $C_2 > 0$ that is independent of b . Inserting this bound in (5.4) yields

$$C_1 \|f\|_{L^2}^2 \geq \int_r^{\sqrt{r}} \frac{1}{\rho} (\text{osc}_{B_r} v - C_2 \|f\|_{L^\infty} \rho^2)^2 d\rho. \quad (5.11)$$

We distinguish two cases, based on whether $\text{osc}_{B_r} v$ is larger or smaller than $2C_2 \|f\|_{L^\infty} r$. When $\text{osc}_{B_r} v \geq 2C_2 \|f\|_{L^\infty} r \geq 2C_2 \|f\|_{L^\infty} \rho^2$, then (5.6) implies that

$$C_1 \|f\|_{L^2}^2 \geq \int_r^{\sqrt{r}} \frac{1}{\rho} \left(\frac{\text{osc}_{B_r} v}{2}\right)^2 d\rho = \frac{(\text{osc}_{B_r} v)^2}{8} \log \frac{1}{r},$$

which implies

$$\text{osc}_{B_r} v \leq \frac{4\sqrt{C_1} \|f\|_{L^2}}{\sqrt{\log 1/r}}.$$

On the other hand

$$\text{osc}_{B_r} v \leq 2C_2 \|f\|_{L^\infty} r \leq \frac{2C_2 \|f\|_{L^\infty}}{\sqrt{\log 1/r}}$$

for any $r \leq r_0^2 \leq 1/2$. The above two estimates imply (5.2). One may repeat this argument with x_0 being any point in \mathbb{T}^2 , not just the origin, by periodically extending v and f to one more periodic cell, thereby concluding the proof. \square

In contrast, the parabolic case is more delicate. If we consider the linear problem

$$\partial_t v + b(x, t) \cdot \nabla v - \Delta v = f,$$

even if b is divergence-free, one may construct solutions that are not continuous functions for all time, although they obey the L^∞ maximum principle. See e.g. [SVZ12] for an example with rough drift. For the nonlinear problem

$$\partial_t \omega + v^{-1} \mathbf{u} \cdot \nabla \omega - \Delta \omega = f,$$

one may hope to prove that in some average sense, the functions on the attractor remain continuous as $v \rightarrow 0$. At the moment we do not know how to prove this.

6. The Damped and Driven Navier-Stokes Equations and Other Scaling

In this section we consider the weakly damped and driven stochastic Navier-Stokes equations

$$d\mathbf{u} + (Y\mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla \pi - v\Delta \mathbf{u})dt = v^\alpha \rho dW = v^\alpha \sum_k \rho_k dW^k, \quad \nabla \cdot \mathbf{u} = 0, \quad (6.1)$$

where $Y = Y_{\tau, \gamma} = \tau \Lambda^{-\gamma} = \tau(-\Delta)^{-\gamma/2}$ and $\tau > 0, \gamma \in [0, 1)$. As above for (3.1) it follows immediately from the Kryloff-Bogoliouboff procedure that there exists an invariant measure μ_v^α for each $v > 0, \alpha \in \mathbb{R}$.

We now prove that $\alpha = 0$ is the only scaling of v in (6.1) which gives a nontrivial inviscid limit.

Theorem 6.1 (Inviscid limits in different scalings). *For $\alpha \in \mathbb{R}$ consider a collection of invariant measures $\{\mu_v^\alpha\}_{v>0}$ of (6.1). Depending on the choice of α we have one of the following three scenarios:*

- (i) *If $\alpha > 0$, then for any $v_j \rightarrow 0$, we have that $\mu_{v_j}^\alpha \rightarrow \delta_0$, i.e. weakly in $Pr(H^0)$, where δ_0 is the Dirac measure concentrated at 0.*
- (ii) *If $\alpha < 0$, then for any $v_n \rightarrow 0$ such that the $\mu_{v_n}^\alpha \rightarrow \mu_0$ then*

$$\int_{H^0} \|\mathbf{u}\|_{L^2}^2 d\mu_0(\mathbf{u}) = \infty. \quad (6.2)$$

- (iii) *If $\alpha = 0$, then there exists a sequence $v_n \rightarrow 0$ and a stationary martingale solution μ_0 of*

$$d\mathbf{u} + (Y\mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla \pi)dt = \rho dW, \quad \nabla \cdot \mathbf{u} = 0, \quad (6.3)$$

such that $\mu_{v_n}^0 \rightarrow \mu_0$. By a stationary Martingale solution of (6.3) corresponding to μ_0 we mean that there exists a stochastic basis $\mathcal{S} := (\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P}, W)$ and a predictable process

$$\mathbf{u} \in C_w([0, T]; H^0) \cap L^2([0, T], H^{1-\gamma/2}) \quad (6.4)$$

which satisfies (6.3) and is stationary, i.e. the law $\mathbb{P}(\mathbf{u}(t) \in A), A \in \mathcal{B}(H^0)$, is independent of t and identically equal to μ_0 .

PROOF. Let \mathbf{u}^v be stationary solutions of (6.1) corresponding to μ_v^α , and define $\omega^v = \nabla^\perp \cdot \mathbf{u}^v$. Applying the Itô lemma to (6.1) and using stationarity, we obtain:

$$\mathbb{E} (v \|\nabla \mathbf{u}^v\|_{L^2}^2 + \|Y^{1/2} \mathbf{u}^v\|_{L^2}^2) = \frac{v^{2\alpha}}{2} \|\rho\|_{L^2}^2. \quad (6.5)$$

Additionally, by making use of the vorticity formulation of (6.1),

$$d\omega + (Y\omega + \mathbf{u} \cdot \nabla \omega - v\Delta \omega)dt = v^\alpha \sigma dW, \quad \sigma = \nabla^\perp \rho, \quad (6.6)$$

we also obtain, again with the Itô lemma and stationarity

$$\mathbb{E} (v \|\nabla \omega^v\|_{L^2}^2 + \|Y^{1/2} \omega^v\|_{L^2}^2) = \frac{v^{2\alpha}}{2} \|\sigma\|_{L^2}^2. \quad (6.7)$$

PROOF OF (i). We begin with the case $\alpha > 0$. From (6.7) we have

$$\mathbb{E} \|\mathbf{u}^v\|_{H^{1-\gamma/2}}^2 \leq \frac{v^{2\alpha}}{2\tau} \|\sigma\|_{L^2}^2. \quad (6.8)$$

Using Chebyshev and compact embedding we infer that μ_ν^α is tight in $Pr(H^0)$.

Consider any weakly convergent subsequence $\mu_\nu^\alpha \rightarrow \mu_0^\alpha$. By the Skhorohod embedding theorem we may find a new probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ and a sequence of H^0 valued random variables $\tilde{\mathbf{u}}^\nu$ such that $\tilde{\mathbf{u}}^\nu$ is equal in law to μ_ν^α and

$$\tilde{\mathbf{u}}^\nu \rightarrow \tilde{\mathbf{u}}^0 \quad \text{a.s. in } H^0,$$

with $\tilde{\mathbf{u}}^0$ equal in law to μ_0^α . Now with (6.8) and Fatou's Lemma we infer

$$\tilde{\mathbb{E}} \|\tilde{\mathbf{u}}^0\|_{L^2}^2 \leq \liminf_{\nu > 0} \tilde{\mathbb{E}} \|\tilde{\mathbf{u}}^\nu\|_{L^2}^2 \leq \liminf_{\nu > 0} \frac{\nu^{2\alpha}}{2\tau} \|\sigma\|_{L^2}^2 = 0.$$

Hence $\tilde{\mathbf{u}}^0 = 0$ a.s. and therefore $\mu_0^\alpha = \delta_0$. This proves the first item.

PROOF OF (ii). Now we consider the case $\alpha < 0$. For every $\nu > 0$ we obtain

$$\begin{aligned} \nu \mathbb{E} \|\nabla \mathbf{u}^\nu\|_{L^2}^2 &= \nu \mathbb{E} \|\omega\|_{L^2}^2 \leq C \nu \mathbb{E} \left(\|Y^{1/2} \omega^\nu\|_{L^2}^{\frac{4}{2+\gamma}} \|\nabla \omega^\nu\|_{L^2}^{\frac{2\gamma}{2+\gamma}} \right) \\ &\leq C \nu^{\frac{2}{2+\gamma}} \mathbb{E} (\nu \|\nabla \omega^\nu\|_{L^2}^2 + \|Y^{1/2} \omega^\nu\|_{L^2}^2) \leq \nu^{\frac{2}{2+\gamma} + 2\alpha} C \|\sigma\|_{L^2}^2, \end{aligned}$$

where we have used that $Y = \tau \Lambda^{-\gamma}$, interpolation and the above balance relation (6.7). Combining (6.5) with the above estimate we obtain

$$\frac{\nu^{2\alpha}}{2} \|\rho\|_{L^2}^2 \leq \mathbb{E} \|Y^{1/2} \mathbf{u}^\nu\|_{L^2}^2 + \nu^{\frac{2}{2+\gamma} + 2\alpha} C \|\sigma\|_{L^2}^2,$$

for constant $C > 0$ which is independent of ν . This implies

$$\nu^{2\alpha} (\|\rho\|_{L^2}^2 / 2 - \nu^{\frac{2}{2+\gamma}} C \|\sigma\|_{L^2}^2) \leq \mathbb{E} \|Y^{1/2} \mathbf{u}^\nu\|_{L^2}^2 \leq C \mathbb{E} \|\mathbf{u}^\nu\|_{L^2}^2. \quad (6.9)$$

By assumption the assumption that $\mu_\nu^\alpha \rightarrow \mu_0^\alpha$ as in (i), the Skhorohod embedding theorem yields a new probability space and $\tilde{\mathbf{u}}^\nu$ which converges almost surely to $\tilde{\mathbf{u}}^0$, with the same laws as the original sequence. We infer

$$\limsup_{\nu \rightarrow 0} \nu^{2\alpha} (\|\rho\|_{L^2}^2 / 2 - \nu^{\frac{2}{2+\gamma}} C \|\sigma\|_{L^2}^2) \leq C \limsup_{\nu \rightarrow 0} \tilde{\mathbb{E}} \|\tilde{\mathbf{u}}^\nu\|_{L^2}^2 \leq C \tilde{\mathbb{E}} \|\tilde{\mathbf{u}}^0\|_{L^2}^2,$$

where we have used the Fatou lemma in the last estimate. This proves (6.2).

PROOF OF (iii). Lastly we treat the case $\alpha = 0$. By applying the Itô lemma to (6.6) we have that

$$d \|\omega^\nu\|_{L^2}^2 + (2 \|Y \omega^\nu\|_{L^2}^2 + 2\nu \|\omega^\nu\|_{H^1}^2) dt = \|\rho\|_{L^2}^2 dt + \langle \rho, \omega^\nu \rangle dW$$

Using stationarity we immediately obtain that

$$\mathbf{u}^\nu \text{ is uniformly bounded in } L^2(\Omega; L^2([0, T], H^{1-\gamma/2})) \quad (6.10)$$

and moreover that

$$\mathbf{u}^\nu(0) \text{ is uniformly bounded in } L^2(\Omega; H^{1-\gamma/2}) \text{ which implies that } \{\mu_\nu\}_{\nu > 0} \text{ is tight on } H^0. \quad (6.11)$$

Returning to (6.1) and again making use of the Itô formula,

$$d \|\mathbf{u}^\nu\|_{L^2}^2 + (2 \|Y \mathbf{u}^\nu\|_{L^2}^2 + 2\nu \|\mathbf{u}^\nu\|_{H^1}^2) dt = \|\sigma\|_{L^2}^2 dt + \langle \sigma, \mathbf{u}^\nu \rangle dW,$$

we infer with (6.11) that

$$\mathbf{u}^\nu \text{ is uniformly bounded in } L^2(\Omega; L^\infty([0, T], H^0)). \quad (6.12)$$

In order to obtain a suitable compactness required to pass to the limit we need some additional uniform estimates on fractional the time derivatives of \mathbf{u}^ν . We will apply the Aubin-Lions type compact embedding

$$L^2([0, T], H^{1-\gamma/2}) \cap W^{1/3,2}([0, T]; H^{-3}) \subset\subset L^2([0, T], H^0) \quad (6.13)$$

and the Arzela-Ascoli type compact embedding

$$W^{1/3,4}([0, T]; H^{-3}) \subset\subset C([0, T]; H^{-4}), \quad (6.14)$$

(see [FG95]). Define a sequence of measures $\{\mu_\nu\}_{\nu>0}$ on the path space $C([0, T]; H^{-4})$ associated to $\{\mathbf{u}^\nu\}_{\nu>0}$ by

$$\mu_\nu(A) = \mathbb{P}(\mathbf{u}^\nu \in A), \quad A \in \mathcal{B}(C[0, T]; H^{-4}). \quad (6.15)$$

Using the embeddings, (6.13), (6.14), and suitable estimates we will next show that

$$\{\mu_\nu\}_{\nu>0} \text{ is tight in } L^2([0, T]; H^0) \cap C([0, T]; H^{-3}). \quad (6.16)$$

For this propose we write (6.1) in its integral form

$$\mathbf{u}^\nu(t) = \left(\mathbf{u}^\nu(0) - \int_0^t (Y\mathbf{u} + P(\mathbf{u} \cdot \nabla \mathbf{u}) - \nu \Delta \mathbf{u}) ds \right) + \sigma W(t) := I_D(t) + I_S(t), \quad (6.17)$$

where P is the Leray projection operator onto L^2 divergence-free vector fields. Observe that,

$$\begin{aligned} \|I_D(t)\|_{W^{1/3,4}([0,T];H^{-3})}^4 &\leq C \int_0^T (\|\mathbf{u}^\nu(0)\|_{H^{-3}}^4 + \|Y\mathbf{u}^\nu\|_{H^{-3}}^4 + \|\nu \Delta \mathbf{u}^\nu\|_{H^{-3}}^4 + \|P(\mathbf{u}^\nu \cdot \nabla \mathbf{u}^\nu)\|_{H^{-3}}^4) dt \\ &\leq C \sup_{t \in [0, T]} (1 + \|\mathbf{u}^\nu\|_{L^2}^4)^2, \end{aligned} \quad (6.18)$$

where C is independent of $1 \geq \nu > 0$, but may depend on T . By making use of a suitable version of the Burkholder-Davis-Gundy (see e.g [FG95]) inequality we have

$$\mathbb{E} \|I_S(t)\|_{W^{1/3,4}([0,T];H^{-3})}^4 \leq C \|\sigma\|_{L^2}^4, \quad (6.19)$$

where again, by the assumption $\alpha = 0$, C is independent of ν but depends on T . Similar estimates yield

$$\|I_D(t)\|_{W^{1/3,2}([0,T];H^{-3})}^2 \leq C \sup_{t \in [0, T]} (1 + \|\mathbf{u}^\nu\|_{L^2}^4), \quad \mathbb{E} \|I_S(t)\|_{W^{1/3,2}([0,T];H^{-3})}^2 \leq C \|\sigma\|_{L^2}^2. \quad (6.20)$$

To establish the first part of the tightness bound, (6.16) we consider the sets

$$B_R^1 := \{\mathbf{u} : \|\mathbf{u}\|_{L^2([0,T],H^{1-\gamma/2})} \leq R\} \cap \{\mathbf{u} : \|\mathbf{u}\|_{W^{1/3,2}([0,T];H^{-3})} \leq R\}$$

so that, according to (6.13) B_R^1 is compact in $L^2([0, T], H^0)$ for every $R > 1$. Observe that, with an appropriate application of Chebyshev's inequality and in view of (6.10), (6.12), (6.17), (6.20)

$$\begin{aligned} \mu^\nu((B_R^1)^c) &\leq \frac{1}{R^2} \mathbb{E} \|\mathbf{u}^\nu\|_{L^2([0,T],H^{1-\gamma/2})}^2 + \frac{2}{R} \mathbb{E} \|I_D(t)\|_{W^{1/3,2}([0,T];H^{-3})} + \frac{4}{R^2} \mathbb{E} \|I_S(t)\|_{W^{1/3,2}([0,T];H^{-3})}^2 \\ &\leq \frac{C}{R} \mathbb{E} \left(\|\mathbf{u}^\nu\|_{L^2([0,T],H^{1-\gamma/2})}^2 + \|\mathbf{u}^\nu\|_{L^\infty([0,T],H^0)}^2 + \|\sigma\|_{L^2}^2 \right) \leq \frac{C}{R}, \end{aligned} \quad (6.21)$$

where C is independent of $\nu > 0$ and $R > 0$. For the second half of the tightness bound (6.16) we define

$$B_R^2 := \{\mathbf{u} : \|\mathbf{u}\|_{W^{1/3,4}([0,T];H^{-3})} \leq R\},$$

and observe with (6.18), (6.17) (6.19) that

$$\begin{aligned} \mu^\nu((B_R^2)^c) &\leq \frac{2}{R} \mathbb{E} \|I_D(t)\|_{W^{1/3,4}([0,T];H^{-3})} + \frac{8}{R^4} \mathbb{E} \|I_S(t)\|_{W^{1/3,4}([0,T];H^{-3})}^4 \\ &\leq \frac{C}{R} \mathbb{E} \left(\|\mathbf{u}^\nu\|_{L^\infty([0,T],H^0)}^2 + \|\sigma\|_{L^2}^4 \right) \leq \frac{C}{R}, \end{aligned} \quad (6.22)$$

where, once again, C is independent of $\nu > 0$ and $R > 0$. With (6.21), (6.22) we may now infer (6.16).

With (6.16) in hand we now invoke the Skorokhod theorem obtain a sequence of processes $(\tilde{\mathbf{u}}^\nu, \tilde{W}^\nu)$ defined on a new probability space, $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ such that

$$\tilde{\mathbf{u}}^\nu \rightarrow \tilde{\mathbf{u}} \quad \text{almost surely in } C([0, T]; H^{-3}) \cap L^2([0, T]; H^0), \quad (6.23)$$

$$\tilde{W}^\nu \rightarrow \tilde{W} \quad \text{almost surely in } C([0, T]; \mathfrak{U}_0), \quad (6.24)$$

$\widetilde{\mathbf{u}}^\nu$ is equal in law to \mathbf{u}^ν and each \widetilde{W}^ν is a cylindrical Brownian motion relative to the filtration $\widetilde{\mathcal{F}}_t^\nu$ given by the completion of $\sigma((\widetilde{W}^\nu(s), \widetilde{\mathbf{u}}^\nu(s)) : s \leq t)$. Note that the sequence $\widetilde{\mathbf{u}}^\nu$ maintains the same uniform bounds as in (6.10), (6.12) and it follows that

$$\widetilde{\mathbf{u}}^\nu \rightharpoonup \widetilde{\mathbf{u}} \quad \text{weakly in } L^2(\Omega; L^2([0, T]; H^{1-\gamma/2})), \quad (6.25)$$

$$\widetilde{\mathbf{u}}^\nu \rightharpoonup^* \widetilde{\mathbf{u}} \quad \text{weakly* in } L^\infty(\Omega; L^\infty([0, T]; H^0)). \quad (6.26)$$

Finally, as in [FG95] that $\widetilde{\mathbf{u}}$ is a stationary process on H^0 with stationary distribution μ_0 .

An argument from [Ben95] may now be employed to show that, for each $\nu > 0$, $\widetilde{\mathbf{u}}^\nu$ solves (6.1), but relative to the new stochastic basis $\widetilde{\mathcal{S}} = (\widetilde{\Omega}, \widetilde{\mathcal{F}}, \widetilde{\mathbb{P}}, \widetilde{\mathcal{F}}_t^\nu, \widetilde{W}^\nu)$. Now by using the convergences (6.23)–(6.26) we may pass to the limit $\nu \rightarrow 0$ in (6.17) (with \mathbf{u}^ν, W appropriately replaced with $\widetilde{\mathbf{u}}^\nu, \widetilde{W}^\nu$) and establish that $(\widetilde{\mathbf{u}}, \widetilde{W})$ satisfies (6.3) along with the required regularity. As in [FG95] it follows from the stationarity of $\widetilde{\mathbf{u}}^\nu$ and (6.23) that $\widetilde{\mathbf{u}}$ is stationary. □

Acknowledgments. We would like to thank Sergei Kuksin and Armen Shirikyan for stimulating discussions and helpful feedback on this work. The authors gratefully acknowledge the support of the Institute for Mathematics and its Applications (IMA), at the University of Minnesota where this work was initiated. VS is partially supported by NSF Grants DMS-1159376, DMS-1101428. VV is supported in part by NSF Grant DMS-1211828.

References

- [Bat69] G.K. Batchelor. Computation of the energy spectrum in homogeneous two-dimensional turbulence. *Physics of fluids*, 12(Suppl. II):233–239, 1969.
- [Ben95] A. Bensoussan. Stochastic Navier-Stokes equations. *Acta Appl. Math.*, 38(3):267–304, 1995.
- [Bes08] H. Bessaih. Stationary solutions for the 2D stochastic dissipative Euler equation. In *Seminar on Stochastic Analysis, Random Fields and Applications V*, volume 59 of *Progr. Probab.*, pages 23–36. Birkhäuser, Basel, 2008.
- [BF00] A. Bensoussan and J. Frehse. Local solutions for stochastic Navier Stokes equations. *M2AN Math. Model. Numer. Anal.*, 34(2):241–273, 2000. Special issue for R. Temam’s 60th birthday.
- [BF12] H. Bessaih and B. Ferrario. Inviscid limit of stochastic damped 2d navier-stokes equations. 12 2012.
- [BKL01] J. Bricmont, A. Kupiainen, and R. Lefevere. Ergodicity of the 2D Navier-Stokes equations with random forcing. *Comm. Math. Phys.*, 224(1):65–81, 2001. Dedicated to Joel L. Lebowitz.
- [BP00] Z. Brzeźniak and S. Peszat. Strong local and global solutions for stochastic Navier-Stokes equations. In *Infinite dimensional stochastic analysis (Amsterdam, 1999)*, volume 52 of *Verh. Afd. Natuurkd. I. Reeks. K. Ned. Akad. Wet.*, pages 85–98. R. Neth. Acad. Arts Sci., Amsterdam, 2000.
- [Bre00] H. Breckner. Galerkin approximation and the strong solution of the Navier-Stokes equation. *J. Appl. Math. Stochastic Anal.*, 13(3):239–259, 2000.
- [BS09] F. Bouchet and E. Simonnet. Random changes of flow topology in two-dimensional and geophysical turbulence. *Physical review letters*, 102(9):94504, 2009.
- [BT72] A. Bensoussan and R. Temam. Équations aux dérivées partielles stochastiques non linéaires. I. *Israel J. Math.*, 11:95–129, 1972.
- [BT73] A. Bensoussan and R. Temam. Équations stochastiques du type Navier-Stokes. *J. Functional Analysis*, 13:195–222, 1973.
- [CC04] A. Córdoba and D. Córdoba. A maximum principle applied to quasi-geostrophic equations. *Comm. Math. Phys.*, 249(3):511–528, 2004.
- [CF88] P. Constantin and C. Foias. *Navier-Stokes equations*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1988.
- [CG94] M. Capiński and D. Gatarek. Stochastic equations in Hilbert space with application to Navier-Stokes equations in any dimension. *J. Funct. Anal.*, 126(1):26–35, 1994.
- [CR07] P. Constantin and F. Ramos. Inviscid limit for damped and driven incompressible Navier-Stokes equations in \mathbb{R}^2 . *Comm. Math. Phys.*, 275(2):529–551, 2007.
- [Cru89] A. B. Cruzeiro. Solutions et mesures invariantes pour des équations d’évolution stochastiques du type Navier-Stokes. *Exposition. Math.*, 7(1):73–82, 1989.
- [Deb11] A. Debussche. Ergodicity results for the stochastic navier-stokes equations: an introduction. *Preprint*, 2011. (to appear).
- [DG57] E. De Giorgi. Sulla differenziabilità e l’analiticità delle estremali degli integrali multipli regolari. *Mem. Accad. Sci. Torino. Cl. Sci. Fis. Mat. Nat. (3)*, 3:25–43, 1957.

- [DGHT11] A. Debussche, N. Glatt-Holtz, and R. Temam. Local martingale and pathwise solutions for an abstract fluids model. *Physica D*, 240(14-15):1123–1144, 2011.
- [DMS05] L. Denis, A. Matoussi, and L. Stoica. L^p estimates for the uniform norm of solutions of quasilinear SPDE's. *Probab. Theory Related Fields*, 133(4):437–463, 2005.
- [DMS09] L. Denis, A. Matoussi, and L. Stoica. Maximum principle and comparison theorem for quasi-linear stochastic PDE's. *Electron. J. Probab.*, 14:no. 19, 500–530, 2009.
- [DPD03] G. Da Prato and A. Debussche. Ergodicity for the 3D stochastic Navier-Stokes equations. *J. Math. Pures Appl.* (9), 82(8):877–947, 2003.
- [DPZ92] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, volume 44 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1992.
- [DPZ96] G. Da Prato and J. Zabczyk. *Ergodicity for infinite-dimensional systems*, volume 229 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1996.
- [EKMS00] W. E, K. Khanin, A. Mazel, and Ya. Sinai. Invariant measures for Burgers equation with stochastic forcing. *Ann. of Math.* (2), 151(3):877–960, 2000.
- [FG95] F. Flandoli and D. Gatarek. Martingale and stationary solutions for stochastic Navier-Stokes equations. *Probab. Theory Related Fields*, 102(3):367–391, 1995.
- [FJMR02] C. Foias, M. S. Jolly, O. P. Manley, and R. Rosa. Statistical estimates for the Navier-Stokes equations and the Kraichnan theory of 2-D fully developed turbulence. *J. Statist. Phys.*, 108(3-4):591–645, 2002.
- [FM95] F. Flandoli and B. Maslowski. Ergodicity of the 2-D Navier-Stokes equation under random perturbations. *Comm. Math. Phys.*, 172(1):119–141, 1995.
- [FMRT01] C. Foias, O. Manley, R. Rosa, and R. Temam. *Navier-Stokes equations and turbulence*, volume 83 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2001.
- [FR08] F. Flandoli and M. Romito. Markov selections for the 3D stochastic Navier-Stokes equations. *Probab. Theory Related Fields*, 140(3-4):407–458, 2008.
- [Fri95] U. Frisch. *Turbulence*. Cambridge University Press, Cambridge, 1995. The legacy of A. N. Kolmogorov.
- [FŠ] J. Foldes and V. Šverák. *In preparation*.
- [FW12] M.I. Freidlin and A.D. Wentzell. *Random perturbations of dynamical systems*, volume 260 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Heidelberg, third edition, 2012. Translated from the 1979 Russian original by Joseph Szücs.
- [GHZ09] N. Glatt-Holtz and M. Ziane. Strong pathwise solutions of the stochastic Navier-Stokes system. *Advances in Differential Equations*, 14(5-6):567–600, 2009.
- [HM06] M. Hairer and J. C. Mattingly. Ergodicity of the 2D Navier-Stokes equations with degenerate stochastic forcing. *Ann. of Math.* (2), 164(3):993–1032, 2006.
- [HM11] M. Hairer and J. C. Mattingly. A theory of hypoellipticity and unique ergodicity for semilinear stochastic pdes. *Electron. J. Probab.*, 16(23):658–738, 2011.
- [Jud63] V.I. Judovič. Non-stationary flows of an ideal incompressible fluid. *Ž. Vyčisl. Mat. i Mat. Fiz.*, 3:1032–1066, 1963.
- [KB37] N. Kryloff and N. Bogoliouboff. La théorie générale de la mesure dans son application à l'étude des systèmes dynamiques de la mécanique non linéaire. *Ann. of Math.* (2), 38(1):65–113, 1937.
- [KM80] R. H. Kraichnan and D. Montgomery. Two-dimensional turbulence. *Rep. Prog. Phys.*, 43:574–619, 1980.
- [KP05] S. Kuksin and O. Penrose. A family of balance relations for the two-dimensional Navier-Stokes equations with random forcing. *J. Stat. Phys.*, 118(3-4):437–449, 2005.
- [Kra67] R. H. Kraichnan. Inertial ranges in two-dimensional turbulence. *Phys. Fluids*, 10(7):1417–1423, 1967.
- [Kry10] N. V. Krylov. Itô's formula for the L_p -norm of stochastic W_p^1 -valued processes. *Probab. Theory Related Fields*, 147(3-4):583–605, 2010.
- [KS91] I. Karatzas and S. E. Shreve. *Brownian motion and stochastic calculus*, volume 113 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1991.
- [KS01] S. Kuksin and A. Shirikyan. A coupling approach to randomly forced nonlinear PDE's. I. *Comm. Math. Phys.*, 221(2):351–366, 2001.
- [KS02] S. Kuksin and A. Shirikyan. Coupling approach to white-forced nonlinear PDEs. *J. Math. Pures Appl.* (9), 81(6):567–602, 2002.
- [KS12] S. Kuksin and A. Shirikyan. *Mathematics of Two-Dimensional Turbulence*. Number 194 in Cambridge Tracts in Mathematics. Cambridge University Press, 2012.
- [Kuk99] I. Kukavica. On the dissipative scale for the Navier-Stokes equation. *Indiana Univ. Math. J.*, 48(3):1057–1081, 1999.
- [Kuk04] S. B. Kuksin. The Eulerian limit for 2D statistical hydrodynamics. *J. Statist. Phys.*, 115(1-2):469–492, 2004.
- [Kuk06a] S. B. Kuksin. Remarks on the balance relations for the two-dimensional Navier-Stokes equation with random forcing. *J. Stat. Phys.*, 122(1):101–114, 2006.
- [Kuk06b] S. B. Kuksin. *Randomly forced nonlinear PDEs and statistical hydrodynamics in 2 space dimensions*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2006.

- [Kuk07] S. B. Kuksin. Eulerian limit for 2D Navier-Stokes equation and damped/driven KdV equation as its model. *Tr. Mat. Inst. Steklova*, 259(Anal. i Osob. Ch. 2):134–142, 2007.
- [Kuk08] S. B. Kuksin. On distribution of energy and vorticity for solutions of 2D Navier-Stokes equation with small viscosity. *Comm. Math. Phys.*, 284(2):407–424, 2008.
- [Kup10] A. Kupiainen. Ergodicity of two dimensional turbulence. *arXiv:1005.0587v1 [math-ph]*, 05 2010.
- [Leb07] H. Lebesgue. Sur le problème de dirichlet. *Rend. Circ. Mat. Palermo*, 1(24):371–402, 1907.
- [Mar87] C. Marchioro. An example of absence of turbulence for any Reynolds number. II. *Comm. Math. Phys.*, 108(4):647–651, 1987.
- [Mat99] J. C. Mattingly. Ergodicity of 2D Navier-Stokes equations with random forcing and large viscosity. *Comm. Math. Phys.*, 206(2):273–288, 1999.
- [Mat02] J. C. Mattingly. Exponential convergence for the stochastically forced Navier-Stokes equations and other partially dissipative dynamics. *Comm. Math. Phys.*, 230(3):421–462, 2002.
- [Mat03] J. C. Mattingly. On recent progress for the stochastic Navier Stokes equations. In *Journées “Équations aux Dérivées Partielles”*, pages Exp. No. XI, 52. Univ. Nantes, Nantes, 2003.
- [MB02] A.J. Majda and A.L. Bertozzi. *Vorticity and incompressible flow*, volume 27 of *Cambridge Texts in Applied Mathematics*. Cambridge University Press, Cambridge, 2002.
- [Mil90] J. Miller. Statistical mechanics of Euler equations in two dimensions. *Phys. Rev. Lett.*, 65(17):2137–2140, 1990.
- [Mos60] J. Moser. A new proof of De Giorgi’s theorem concerning the regularity problem for elliptic differential equations. *Comm. Pure Appl. Math.*, 13:457–468, 1960.
- [MP06] J. C. Mattingly and É. Pardoux. Malliavin calculus for the stochastic 2D Navier-Stokes equation. *Comm. Pure Appl. Math.*, 59(12):1742–1790, 2006.
- [MR04] R. Mikulevičius and B. L. Rozovskii. Stochastic Navier-Stokes equations for turbulent flows. *SIAM J. Math. Anal.*, 35(5):1250–1310, 2004.
- [MR05] R. Mikulevičius and B. L. Rozovskii. Global L_2 -solutions of stochastic Navier-Stokes equations. *Ann. Probab.*, 33(1):137–176, 2005.
- [MV11] C. Mouhot and C. Villani. On Landau damping. *Acta Math.*, 207(1):29–201, 2011.
- [MWC92] J. Miller, P.B. Weichman, and M.C. Cross. Statistical mechanics, euler’s equation, and jupiter’s red spot. *Physical Review A*, 45(4):2328–2359, 1992.
- [Nas58] J. Nash. Continuity of solutions of parabolic and elliptic equations. *Amer. J. Math.*, 80:931–954, 1958.
- [Nov65] E. A. Novikov. Functionals and the random-force method in turbulence theory. *Soviet Physics JETP*, 20:1290–1294, 1965.
- [Rob91] R. Robert. A maximum-entropy principle for two-dimensional perfect fluid dynamics. *J. Statist. Phys.*, 65(3-4):531–553, 1991.
- [RY99] D. Revuz and M. Yor. *Continuous martingales and Brownian motion*, volume 293 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, third edition, 1999.
- [Shi11] A. Shirikyan. Local times for solutions of the complex Ginzburg-Landau equation and the inviscid limit. *J. Math. Anal. Appl.*, 384(1):130–137, 2011.
- [Shn93] A. I. Shnirelman. Lattice theory and flows of ideal incompressible fluid. *Russian J. Math. Phys.*, 1(1):105–114, 1993.
- [SSŠZ12] G. Seregin, L. Silvestre, V. Šverák, and A. Zlatoš. On divergence-free drifts. *J. Differential Equations*, 252(1):505–540, 2012.
- [Šve12] V. Šverák. Course notes. <http://math.umn.edu/sverak/course-notes2011.pdf>, 2011/2012.
- [SVZ12] L. Silvestre, V. Vicol, and A. Zlatoš. On the loss of continuity for super-critical drift-diffusion equations. *Archive for Rational Mechanics and Analysis*, 2012.
- [Tab02] P. Tabeling. Two-dimensional turbulence: a physicist approach. *Physics Reports*, 362(1):1–62, 2002.
- [Tsi01] A. Tsinober. *An informal introduction to turbulence*, volume 63. Springer, 2001.
- [Tur99] B. Turkington. Statistical equilibrium measures and coherent states in two-dimensional turbulence. *Comm. Pure Appl. Math.*, 52(7):781–809, 1999.
- [Vio76] M. Viot. *Solutions faibles d’équations aux dérivées partielles non linéaires*. 1976. Thèse, Université Pierre et Marie Curie, Paris.
- [VKF79] M. I. Vishik, A. I. Komech, and A. V. Fursikov. Some mathematical problems of statistical hydromechanics. *Uspekhi Mat. Nauk*, 34(5(209)):135–210, 256, 1979.

INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN 55455 AND
DEPARTMENT OF MATHEMATICS, VIRGINIA TECH, BLACKSBURG, VA 24061
E-mail address: negh@ima.umn.edu, negh@vt.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN 55455
E-mail address: sverak@math.umn.edu

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08544
E-mail address: vvicol@math.princeton.edu