

JOINT CONVERGENCE OF RANDOM QUADRANGULATIONS AND THEIR CORES

LOUIGI ADDARIO-BERRY AND YUTING WEN

ABSTRACT. We show that a uniform quadrangulation, its largest 2-connected block, and its largest simple block jointly converge to the same Brownian map in distribution for the Gromov-Hausdorff-Prokhorov topology. We start by deriving a local limit theorem for the asymptotics of maximal block sizes, extending the result in [3]. The resulting diameter bounds for pendant submaps of random quadrangulations straightforwardly lead to Gromov-Hausdorff convergence. To extend the convergence to the Gromov-Hausdorff-Prokhorov topology, we show that exchangeable “uniformly asymptotically negligible” attachments of mass simply yield, in the limit, a deterministic scaling of the mass measure.

1. INTRODUCTION

Much work has been devoted to understanding the asymptotic properties of large random planar maps. It is conjectured and known in several cases, that after rescaling the graph distance properly, planar maps from many families converge to the same universal metric space, the Brownian map, in the Gromov-Hausdorff-Prokhorov sense. Recently Le Gall [11] and Miermont [14] independently proved that the Brownian map is the scaling limit of several important families of planar maps, and Addario-Berry & Albenque [1] proved that simple triangulations and simple quadrangulations also rescale to the same limit object.

Further results, the aim of this paper is to show that random quadrangulations and their cores jointly converge to the same limit object, even after conditioning on their sizes. Before making this more precise, we state one corollary (Theorem 1.1) of our main result: the Brownian map is again the scaling limit of random 2-connected quadrangulations.

For the duration of the paper, all maps are embedded in the sphere \mathbb{S}^2 and are considered up to orientation preserving homeomorphism. A *rooted map* is a pair $M = (M, uv)$ where M is a map and uv is an oriented edge of M , i.e., $\{u, v\} \in e(M)$. A *quadrangulation* is a map in which every face has degree 4. A quadrangulation is *2-connected* if the removal of any vertex does not disconnect the map. It is *simple* if it contains no multiple edges. Write \mathcal{Q}, \mathcal{R} and \mathcal{S} for the set of rooted connected, 2-connected, and simple quadrangulations, respectively. It is easy to verify that simple quadrangulations are 2-connected, so $\mathcal{S} \subset \mathcal{R} \subset \mathcal{Q}$. It is technically convenient to view a single edge as a 2-connected, simple quadrangulation, and we do this.

Given a finite set \mathcal{G} , the notation $G \in_u \mathcal{G}$ means that G is chosen uniformly at random from \mathcal{G} . Given a finite rooted or unrooted map G write μ_G for the uniform probability measure on the vertex set $v(G)$, and for $c > 0$, write cG for the measured metric space $(v(G), c \cdot d_G, \mu_G)$, where d_G denotes graph distance. Given a set \mathcal{G} of maps and $n \in \mathbb{N}$, write $\mathcal{G}_n = \{G \in \mathcal{G} : |v(G)| = n\}$. Finally, write $\mathbf{M} = (\mathcal{M}, d, \mu)$ for the measured Brownian map. (See [11] for a definition of \mathbf{M} .)

Date: March 23, 2015.

Key words and phrases. Brownian map, Gromov-Hausdorff-Prokhorov convergence, singularity analysis.

Theorem 1.1. *Let $\mathbf{R}_r \in_u \mathcal{R}_r$, then as $r \rightarrow \infty$,*

$$\left(\frac{21}{40r}\right)^{1/4} \mathbf{R}_r \xrightarrow{d} \mathbf{M}$$

in distribution for the Gromov-Hausdorff-Prokhorov topology.

A brief overview of the Gromov-Hausdorff-Prokhorov (GHP) distance appears in Section 2.2.




FIGURE 1. $(0, 1)$ is the root edge of M . For the total order \leq_M we have, e.g., $(0, 1) < (0, 2)$, $(2, 12) < (12, 2) < (12, 111) < (111, 12)$. Also, of the two copies of edge $(11, 111)$, the one succeeding $(11, 2)$ in the clockwise order is smaller for $<$. The simple block $S(M)$, highlighted in red, has vertices $0, 1, 2, 11, 12, 111$.

clear.) Note that M is nearly simple precisely if M° is simple – in this case we call M° the *simple nerve* of M .

For $A \subset v(M)$, write $M[A]$ for the submap of M induced by A . For any edge $e = \{x, y\} \in e(M)$, let $B_e \subset v(M)$ be maximal subject to the constraints that $\{x, y\} \subset B_e$, and that $M[B_e]$ is 2-connected. We call $M[B_e]^\circ$ a *2-connected block* of M . In particular, write $R^\bullet = R^\bullet(M) = (M[B_{uv}]^\circ, uv)$ and call R^\bullet the *2-connected root block* of M . Likewise, for any edge $e = \{x, y\} \in e(M)$, let $B'_e \subset v(M)$ be maximal such that $\{x, y\} \subset B'_e$, that $M[B'_e]$ is nearly simple, and, subject to these constraints, with vertex set B'_e minimal with respect to $<_M$. We call $M[B'_e]^\circ$ a *simple block* of M . Write $S^\bullet = S^\bullet(M) = (M[B'_{uv}]^\circ, uv)$ and call S^\bullet the *simple root block* of M .

Write $R(M)$ (resp. $S(M)$) for the largest 2-connected (resp. simple) block of M , rooted at its $<_M$ -minimal edge, and write $b(M) = |v(R(M))|$ and $sb(M) = |v(S(M))|$. If there are multiple 2-connected blocks with size $b(M)$, among these blocks we take $R(M)$ to be the one whose root edge $u_i u_j$ is $<_M$ -minimal, and use the same convention for $S(M)$. The next theorem states that a

To state our main results, a little more terminology is needed. Given a rooted map $M = (M, uv)$, we may define a canonical total order $<_M$ on $v(M)$ as follows. List the vertices of M as $u_1 = u, u_2 = v, \dots, u_{|v(M)|}$ according to their order of discovery by a breadth-first search which starts from the root edge uv and uses the clockwise of edges around each vertex starting from the explored edge to determine exploration priority. (See [7] for a definition of breadth-first search.) We also define a total order $<_M$ on the set of oriented edges of M as follows. Let $u_i u_j <_M u_{i'} u_{j'}$ precisely if either (a) u_i was discovered before $u_{i'}$ or (b) $i = i'$ and $u_i u_j$ has higher priority than $u_i u_{j'}$.

Fix a bipartite map $M = (M, uv)$. A cycle C in a map M is *nearly facial* if at least one connected component of $S^2 \setminus C$ contains no vertices of M (it may contain edges). We say M is *nearly simple* if every cycle in M with length two is nearly facial. Write $M^\circ = (M^\circ, uv)$ for the map obtained by collapsing each nearly facial 2-cycle into an edge. (This is a slight abuse of notation as the edge $uv \in e(M)$ may be collapsed with other edges in forming M° , but the meaning should be




FIGURE 2. The 2-connected blocks of M are $M[\{a, b, c, d\}]^\circ$ and $M[\{d, e, f, g, h, i\}]^\circ$. The simple blocks of M are $M[\{a, b, c, d\}]^\circ$, $M[\{d, e, f, i\}]^\circ$, and $M[\{d, g, h, i\}]^\circ$.

uniform quadrangulation, its largest 2-connected block, and its largest simple block jointly converge to the same Brownian map.

Theorem 1.2. *Let $\mathbf{Q}_q \in_u \mathcal{Q}_q$ and write $\mathbf{R}_q = \mathbf{R}(\mathbf{Q}_q)$, $\mathbf{S}_q = \mathbf{S}(\mathbf{Q}_q)$. Then as $q \rightarrow \infty$,*

$$\left(\left(\frac{9}{8q} \right)^{1/4} \mathbf{Q}_q, \left(\frac{9}{8q} \right)^{1/4} \mathbf{R}_q, \left(\frac{9}{8q} \right)^{1/4} \mathbf{S}_q \right) \xrightarrow{d} (\mathbf{M}, \mathbf{M}, \mathbf{M})$$

in distribution for the Gromov-Hausdorff-Prokhorov topology.

The convergence of the first coordinate in Theorem 1.2 was proved independently by Le Gall [11] and by Miermont [14]. The convergence of the third coordinate is related to a result by Addario-Berry & Albenque [1] who show that if \mathbf{S}_q is a uniform simple quadrangulation for all q , then $(3/(8|v(\mathbf{S}_q)|))^{1/4} \mathbf{S}_q \xrightarrow{d} \mathbf{M}$. It is known [3, 9] that $|v(\mathbf{S}_q)|/q \rightarrow 1/3$ in probability, so in the third coordinate the scaling factor $(9/(8q))^{1/4}$ may be replaced by $(3/(8|v(\mathbf{S}_q)|))^{1/4}$, in agreement with [1]. The second coordinate is related to Theorem 1.1, but indicates the convergence of 2-connected quadrangulations with *random* sizes.

Theorem 1.2 and Theorem 1.1 follow from a stronger “local invariance principle”, in which the sizes of the largest 2-connected block and largest simple block are fixed rather than random. Given integers $q \geq r \geq s \geq 1$, let

$$\begin{aligned} \mathcal{Q}_{q,r,s} &= \{Q \in \mathcal{Q}_q : b(Q) = r, \text{sb}(Q) = s\}, \\ \mathcal{R}_{r,s} &= \{Q \in \mathcal{R}_r : \text{sb}(Q) = s\}. \end{aligned}$$

Theorem 1.3. *Let $(r(q), q \geq 1)$ and $(s(q), q \geq 1)$ be such that $r(q) = 7q/15 + O(q^{2/3})$ and $s(q) = q/3 + O(q^{2/3})$ as $q \rightarrow \infty$. Let $\mathbf{Q}_q \in_u \mathcal{Q}_{q,r(q),s(q)}$ and write $\mathbf{R}_q = \mathbf{R}(\mathbf{Q}_q)$, $\mathbf{S}_q = \mathbf{S}(\mathbf{Q}_q)$. Then as $q \rightarrow \infty$,*

$$\left(\left(\frac{9}{8q} \right)^{1/4} \mathbf{Q}_q, \left(\frac{9}{8q} \right)^{1/4} \mathbf{R}_q, \left(\frac{9}{8q} \right)^{1/4} \mathbf{S}_q \right) \xrightarrow{d} (\mathbf{M}, \mathbf{M}, \mathbf{M})$$

in distribution for the Gromov-Hausdorff-Prokhorov topology.

We provide an outline of the proof of Theorem 1.3 (our main result) in Section 1.2.

Now and for the remainder of the paper, fix $C > 0$ and let $(r(q) : q \geq 1)$ and $(s(q) : q \geq 1)$ be such that $|r(q) - 7q/15| < Cq^{2/3}$ and $|s(q) - q/3| < Cq^{2/3}$ for all q sufficiently large. The scaling of $r(q)$ and $s(q)$ in Theorem 1.3 is explained by the following local limit theorem for the asymptotics of maximal block sizes.

Theorem 1.4. *Let $\mathbf{Q}_q \in_u \mathcal{Q}_q$, and write $\delta_r(q) = \frac{r(q) - 7q/15}{q^{2/3}}$, $\delta_s(q) = \frac{s(q) - q/3}{q^{2/3}}$. Then*

$$\mathbb{P}(b(\mathbf{Q}_q) = r(q), \text{sb}(\mathbf{Q}_q) = s(q)) = \frac{\beta \mathcal{A}(\beta \delta_s(q))}{r(q)^{2/3}} \frac{\beta' \mathcal{A}(\beta' \delta_r(q))}{q^{2/3}} (1 + o(1)),$$

where β and β' are positive constants given in Propositions 4.2 and 4.1 respectively, $\mathcal{A} : \mathbb{R} \rightarrow [0, 1]$ is a density.

Here $o(1)$ denotes a function tending to zero whose decay may depend on C , but we omit this dependence from the notation. We prove Theorem 1.4 using the machinery developed by Banderier, Flajolet, Schaeffer & Soria [3], based on singularity analysis of generating functions, in Section 4. Theorem 1.2 follows from Theorem 1.3, Theorem 1.4, and an easy averaging argument. We similarly deduce Theorem 1.1 by averaging over the second coordinate in the next proposition.

Proposition 1.5. *Let $\mathbf{R}_r \in_u \mathcal{R}_{r,s(r)}$ and write $\mathbf{S}_r = \mathbf{S}(\mathbf{R}_r)$. Then as $r \rightarrow \infty$,*

$$\left(\left(\frac{21}{40r} \right)^{1/4} \mathbf{R}_r, \left(\frac{21}{40r} \right)^{1/4} \mathbf{S}_r \right) \xrightarrow{d} (\mathbf{M}, \mathbf{M})$$

in distribution for the Gromov-Hausdorff-Prokhorov topology.

Remarks.

- (1) The proof of Proposition 1.5, given in Section 7, uses the convergence of simple quadrangulations, proved in [1], to deduce convergence of 2-connected quadrangulations, as a stepping stone to proving the joint convergence of Theorem 1.3. The results of [1] in turn use the “rerooting invariance trick” introduced by Le Gall [11], together with the convergence of uniform quadrangulations to the Brownian map [11, 14], to deduce convergence for uniform simple quadrangulations. We mention this to emphasize that the results of this paper do not constitute an independent proof of convergence for uniform quadrangulations.
- (2) In [1] it is also shown that simple triangulations converge to the Brownian map. Using this, the arguments of the current paper could be modified to show joint convergence of uniformly random triangulations and their largest loopless and simple blocks.

Before sketching our proof, we first describe the combinatorial relations between \mathcal{Q} , $\mathbf{R}^\bullet(\mathcal{Q})$ and $\mathbf{S}^\bullet(\mathcal{Q})$, on which our proofs rely.

1.1. Bijections for \mathcal{Q} , \mathbf{R} and \mathbf{S} . Suppose we are given only $\mathbf{R}^\bullet = \mathbf{R}^\bullet(\mathcal{Q})$. What additional information is required to reconstruct \mathcal{Q} ? Similarly, what do we require in addition to $\mathbf{S}^\bullet = \mathbf{S}^\bullet(\mathbf{R})$ in order to reconstruct \mathbf{R}^\bullet ? In each case, the reconstruction requires augmenting the edges with additional data. When reconstructing \mathbf{R}^\bullet from \mathbf{S}^\bullet , this data consists of a 2-connected quadrangulation for each edge of \mathbf{S}^\bullet . When reconstructing \mathcal{Q} from \mathbf{R}^\bullet , we require a *sequence* of quadrangulations for each edge of \mathbf{R}^\bullet , together with a second, binary sequence whose entries specify how to attach the quadrangulations in the sequence. In both cases, the root edge must be treated slightly differently from the others (in brief, for the root edge we must specify data twice, once for each side of the edge). We now turn to details.

A *quadrangulation of a 2-gon* is a rooted map whose unbounded face has degree 2, with all other faces of degree 4, rooted such that the unbounded face lies to the left of the root edge. Temporarily write \mathcal{T} for the set of quadrangulations of 2-gons. Given a map in \mathcal{T} , merge the two edges incident to the unbounded face to obtain a map in \mathcal{Q} ; we call this the natural bijection between \mathcal{T} and \mathcal{Q} . For $n \geq 3$, it in fact restricts to a bijection between \mathcal{T}_n and \mathcal{Q}_n . Also, \mathcal{T}_2 contains only one element: the map with one edge and two vertices. Recalling that we also view a single edge as a 2-connected quadrangulation, it follows that $\mathcal{T}_2 = \mathcal{Q}_2$, and it is convenient to view the natural bijection as associating these two sets with one another.

Let $S = (S, uv)$ be a simple quadrangulation. List the vertices of S in breadth-first order as u_1, \dots, u_n and list the edges of S as $uv = e_1, \dots, e_m$, oriented so that the tail precedes the head in breadth-first order. To build a 2-connected quadrangulation with simple root block S , proceed as follows (see Figure 3).

- (1) Create a second copy e_0 of the edge uv so that e_0 lies to the left of e_1 .
- (2) For $0 \leq i \leq m$ let M_i be a 2-connected quadrangulation, and let $M'_i = (M_i, u_i v_i)$ be the quadrangulation of a 2-gon associated to M_i by the natural bijection.
- (3) For each $0 \leq i \leq m$, identify the edge e_i with the root edge $u_i v_i$ of M'_i . The resulting map has a single facial 2-cycle (lying between M_0 and M_1), with vertices u and v ; collapse it and root at the resulting edge uv .




FIGURE 3. (a) A simple quadrangulation. (b) “Decorations” for the edges. (c) After attaching the decorations. (d) The map R .

Call the resulting map R . Then R is a 2-connected quadrangulation with $S^\bullet(R) = S$. We note that

$$|e(R)| = |e(S)| + \sum_{i=0}^{|e(S)|} |e(M_i)| \mathbf{1}_{[|e(M_i)| \neq 1]} = -1 + \sum_{i=0}^{|e(S)|} (1 + |e(M_i)| \mathbf{1}_{[|e(M_i)| \neq 1]}). \quad (1.1)$$

Proposition 1.6. *The above procedure induces a bijection φ between \mathcal{R} and the set*

$$\{(S, \Theta) : S \in \mathcal{S}, \Theta \in \mathcal{R}^{|e(S)|+1}\}.$$

Proof. Given a 2-connected quadrangulation of a 2-gon, collapsing the unbounded face to form a single edge (which is equivalent to taking the simple nerve), then rooting at this edge, yields a 2-connected quadrangulation. This operation is easily seen to be a bijection. In view of the fact that the quadrangulation $R \in \mathcal{R}$ in the above construction has $S^\bullet(R) = S$, the result follows. \square

Next let $R = (R, uv)$ be a 2-connected quadrangulation and list $e(R)$ as e_1, \dots, e_m , as above. For each $1 \leq i \leq m$, write e_i^+ and e_i^- for the head and the tail of e_i respectively. To build a quadrangulation with 2-connected root block R , proceed as follows (see Figure 4).

- (1) Create a second copy e_0 of the edge uv so that e_0 lies to the left of e_1 .
- (2) For $0 \leq i \leq m$ fix $\ell_i \in \mathbb{N}_{\geq 0}$ and sequences $L_i = (M_{i,j}, 1 \leq j \leq \ell_i) \in \mathcal{Q}^{\ell_i}$, $b_i = (b_{i,j}, 1 \leq j \leq \ell_i) \in \{0, 1\}^{\ell_i}$.
- (3) For each $1 \leq i \leq m$, add an additional ℓ_i copies of e_i ; label the resulting $\ell_i + 1$ copies of e_i as $e_{i,1}, \dots, e_{i,\ell_i+1}$ in clockwise order around e_i^- .
- (4) For $0 \leq i \leq m$ and $1 \leq j \leq \ell_i$, let $M'_{i,j}$ be the quadrangulation of a 2-gon associated to $M_{i,j}$ by the natural bijection.
- (5) Attach $M'_{i,j} = (M_{i,j}, u_{i,j}v_{i,j})$ inside the 2-cycle formed by $e_{i,j}$ and $e_{i,j+1}$ by identifying $u_{i,j}$ with e_i^- (if $b_{i,j} = 0$) or e_i^+ (if $b_{i,j} = 1$). The resulting map has a single facial 2-cycle, with edges e_{0,ℓ_0+1} and $e_{1,1}$; collapse it and root at the resulting edge uv .

Call the resulting map Q . Then Q is a connected quadrangulation with $R^\bullet(Q) = R$. We note that

$$|e(Q)| = |e(R)| + \sum_{i=0}^{|e(R)|} \sum_{j=1}^{\ell_i} (|e(M_{i,j})| + 1 + \mathbf{1}_{[|e(M_{i,j})| \neq 1]})$$




FIGURE 4. The quadrangulation in (d) can be reconstructed from its 2-connected core in (a) with the decoration $((L_i, b_i) : 0 \leq i \leq r)$ where $L_0 = (M_{0,1})$, $b_0 = (1)$, $L_1 = (M_{1,1})$, $b_1 = (0)$, $L_2 = L_3 = \emptyset$, $b_2 = b_3 = \emptyset$, $L_4 = (M_{4,1}, M_{4,2})$, $b_4 = (0, 1)$.

$$= -1 + \sum_{i=0}^{|\mathbf{e}(\mathbf{R})|} \left(1 + \sum_{j=1}^{\ell_i} (|e(M_{i,j})| + 1 + \mathbf{1}_{[|e(M_{i,j})| \neq 1]}) \right) \quad (1.2)$$

In the following proposition we write $(\mathcal{Q} \times \{0, 1\})^* = \{\emptyset\} \cup \bigcup_{n \geq 1} (\mathcal{Q} \times \{0, 1\})^n$.

Proposition 1.7. *The above procedure induces a bijection ψ between \mathcal{Q} and the set*

$$\left\{ (\mathbf{R}, \Gamma) : \mathbf{R} \in \mathcal{R}, \Gamma \in ((\mathcal{Q} \times \{0, 1\})^*)^{|\mathbf{e}(\mathbf{R})|+1} \right\}.$$

Proof. This is immediate from the fact that the above construction has $\mathbf{R}^\bullet(\mathcal{Q}) = \mathcal{R}$. \square

For both decompositions, we refer informally to the maps in the vectors Θ and Γ as *decorations* or as *pendant submaps*.

1.2. Proof sketch for Theorem 1.3. In this subsection, we assume familiarity with the Gromov-Hausdorff and Gromov-Hausdorff-Prokhorov distances. The relevant definitions appear in Section 2. We begin by stating (and sketching the proof of) a joint convergence result for a 2-connected quadrangulation and its largest simple block; the proof of this result contains most of the key ideas for the proof of Theorem 1.3.

Given $\mathbf{R}_r = (R_r, e_r) \in \mathcal{R}_{r, s(r)}$, it is easily seen that $\mathbf{S}_r = \mathbf{S}(\mathbf{R}_r)$ is uniformly distributed over $\mathcal{S}_{s(r)}$. Then by [1, Theorem 1], $(3/8s(r))^{1/4} \mathbf{S}_r \xrightarrow{d} \mathbf{M}$ as $s(r) \rightarrow \infty$. Also, the definition of $s(r)$ guarantees that $(\frac{3}{8s(r)})^{1/4} \cdot (\frac{21}{40r})^{-1/4} \rightarrow 1$ as $r \rightarrow \infty$.

Let e' be the $\langle \mathbf{R}_r \rangle$ -minimal oriented edge of \mathbf{S}_r . If $e_r \in e(\mathbf{S}_r)$ then $\mathbf{S}_r = \mathbf{S}^\bullet(\mathbf{R}_r)$. Write $\mathbf{R}'_r = (R_r, e')$. By Proposition 1.6, \mathbf{R}'_r uniquely decomposes as $\varphi(\mathbf{R}'_r) = (S, \Theta) \in \mathcal{S}_{s(r)} \times \mathcal{R}^{|\mathbf{e}(S)|+1}$, and our choice of e' guarantees that $S = \mathbf{S}_r$. Write $\Theta = (\Theta_i, 0 \leq i \leq 2s(r) - 4)$, and

$$L(\mathbf{R}_r) = \max (|v(\Theta_i)| : 0 \leq i \leq 2s(r) - 4) ,$$

$$D(\mathbf{R}_r) = \max(\text{diam}(\Theta_i) : 0 \leq i \leq 2s(r) - 4) .$$

In words, $L(\mathbf{R}_r)$ and $D(\mathbf{R}_r)$ are the greatest number of vertices and the greatest diameter, respectively, of any submap pendant to the biggest simple block of \mathbf{R}_r . The identification of \mathbf{S}_r as a submap of \mathbf{R}_r gives the bound $d_{\text{GH}}(\mathbf{R}_r, \mathbf{S}_r) \leq \frac{1}{2}D(\mathbf{R}_r)$. To prove that $(\frac{21}{40r})^{1/4} d_{\text{GH}}(\mathbf{R}_r, \mathbf{S}_r) = o(1)$ in probability, it thus suffices to show that $(\frac{21}{40r})^{1/4} D(\mathbf{R}_r) = o(1)$ in probability. (Note that here we have the Gromov-Hausdorff rather than Gromov-Hausdorff-Prokhorov distance!)

To accomplish this, we use the methodology developed by Banderier, Flajolet, Schaeffer & Soria [3], which allows one to describe the largest block size of a map whenever the map may be described by a recursive decomposition into rooted blocks, using a suitable composition schema; this is explained in greater detail in Section 4. We thereby obtain the following distributional result for $|v(\mathbf{S}_r)|$.

Proposition 1.8. *Let $\mathbf{R}_r \in_u \mathcal{R}_r$, then for any $A > 0$, uniformly over $x \in [-A, A]$,*

$$\mathbb{P}\left(\text{sb}(\mathbf{R}_r) = \lfloor 5r/7 + xr^{2/3} \rfloor\right) = \frac{\beta \mathcal{A}(\beta x)}{r^{2/3}}(1 + o(1)) ,$$

where β is given in Proposition 4.2.

The proof of Proposition 1.8 appears in Section 4. The range of values for r in the above local limit theorem is what leads to our choice for the range of $s(r)$ in Theorem 1.3 and Theorem 1.4. The following proposition bounds the size of the largest simple block of a random 2-connected quadrangulation.

Proposition 1.9. *For any $A > 0$, there exist positive constants c_1 and c_2 such that for all $r \in \mathbb{N}$ and for integer $k \in (5r/7 + Ar^{2/3}, r]$, if $\mathbf{R}_r \in_u \mathcal{R}_r$,*

$$\mathbb{P}(\text{sb}(\mathbf{R}_r) = k) \leq c_1 \exp\left(-c_2 r (k/r - 5/7)^3\right) .$$

This proposition is a slight extension of [3, Theorem 1] (which proves similar bounds but requires that $(r - k)/r^{2/3} \rightarrow \infty$). We use Proposition 1.9 in proving stretched exponential tail bounds for the size of the largest pendant submap in a random 2-connected quadrangulation.

Proposition 1.10. *For all $\varepsilon \in (0, 1/3)$, there exist positive constants c_1, c_2 , and $c_3 = c_3(\varepsilon)$ such that for all $r \in \mathbb{N}$, if $\mathbf{R}_r \in_u \mathcal{R}_{r,s(r)}$,*

$$\mathbb{P}\left(L(\mathbf{R}_r) \geq r^{2/3+\varepsilon}\right) \leq c_1 \exp(-c_2 r^{c_3}) .$$

Proofs for Proposition 1.9 and Proposition 1.10 are given in Section 5.

Next we deduce a bound for $D(\mathbf{R}_r)$ through extending a result by Chassaing & Schaeffer [6]. The following proposition follows straightforwardly from [6, Proposition 4].

Proposition 1.11. ([6]) *There exist positive constants y_0, C_1 , and C_2 such that for all $y > y_0$ and $q \in \mathbb{N}$, if $\mathbf{Q}_q \in_u \mathcal{Q}_q$,*

$$\mathbb{P}\left(\text{diam}(\mathbf{Q}_q) > yq^{1/4}\right) \leq C_1 \exp(-C_2 y) .$$

This bound is for connected quadrangulations rather than 2-connected ones. However, at the cost of polynomial corrections, we are able to transfer the result to the latter family of quadrangulations, as shown in Section 5.1. This in particular yields the following bound.

Proposition 1.12. *Let $\mathbf{R}_r \in_u \mathcal{R}_{r,s(r)}$, then there exist positive constants C_1, C_2 , and C_3 such that*

$$\mathbb{P}\left(D(\mathbf{R}_r) \geq r^{5/24}\right) \leq C_1 \exp(-C_2 r^{C_3}) .$$

The above results immediately give rise to Gromov-Hausdorff convergence of $(\mathbf{R}_r, \mathbf{S}_r)$ after rescaling, as shown in Proposition 5.2 in the end of Section 5.1. However, to deduce GHP convergence, the above propositions are insufficient, as they do not guarantee that the uniform measures on $v(\mathbf{R}_r)$ and $v(\mathbf{S}_r)$ are close in the Prokhorov sense. Here is an example of the sort of issue that may *a priori* still occur. For all $s \in \mathbb{N}$, let $\mathbf{S}_s \in_u \mathcal{S}_s$ have root edge e_s . Let P_s be the quadrangulation of a 2-gon with $2\lfloor s/5 \rfloor + 2$ vertices composed of parallel alternating 1-paths and 3-paths, and write e'_s for one of the boundary edges of P_s . Then identify e_s with e'_s , embed P_s in the face of \mathbf{S}_s to the left of e_s , and write \mathbf{R}'_s for the resulting quadrangulation; see Figure 5. Recall that $\mathbf{M} = (M, d, \mu)$ is the Brownian map. Then it is not hard to see that $(\mathbf{R}'_s, \mathbf{S}_s)$ converges after rescaling to $(\mathbf{M}', \mathbf{M})$, where $\mathbf{M}' = (M, d, \mu')$ has the geometry of the Brownian map but has mass measure $\frac{5}{7}\mu + \frac{2}{7}\delta_\rho$, where ρ is a point of M with law μ .




FIGURE 5. Parallel alternating 1-paths and 3-paths attached to the root edge e_s .

To prevent the masses of “pendant submaps” from concentrating on small regions in this manner, we use that they attach to exchangeable random locations on the simple block and that each of them has asymptotically negligible size. The first follows from the details of the construction of a 2-connected quadrangulation from its simple root block, explained in Section 1.1; the second is a consequence of Proposition 1.10.

In order to show that the facts from the preceding paragraph suffice to imply joint convergence, we prove a general result on the preservation of Gromov-Hausdorff-Prokhorov convergence under small random modifications; our result relies on results of Aldous on concentration for partial sums of exchangeable random variables. Details for this part of the proof appear in Sections 6 and 7.

We conclude the proof sketch by explaining how we strengthen Proposition 1.5 to prove Theorem 1.3. First, with $\mathbf{Q}_q \in_u \mathcal{Q}_{q,r(q),s(q)}$, we show that $\mathbf{R}(\mathbf{Q}_q)$ contains $\mathbf{S}(\mathbf{Q}_q)$ with high probability. The joint convergence of the second and third coordinates in Theorem 1.3 then follows from Proposition 1.5.

The convergence of the first coordinate does not follow from the existing result by Le Gall [11] or Miermont [14], because \mathbf{Q}_q here is not uniformly distributed over \mathcal{Q}_q , but conditioned on $b(\mathbf{Q}_q) = r(q)$ and $sb(\mathbf{Q}_q) = s(q)$. To deal with this, we require versions of Propositions 1.10 and 1.12 that apply to uniform quadrangulations sampled from \mathcal{Q}_q and $\mathcal{Q}_{q,r(q),s(r(q))}$. These follows straightforwardly once we show that with high probability, $\mathbf{S}(\mathbf{R}(\mathbf{Q}_q)) = \mathbf{S}(\mathbf{Q}_q)$. We postpone the details.

A reprise of the argument for Proposition 1.5 then shows that if $\mathbf{Q}_q \in_u \mathcal{Q}_{q,r(q),s(q)}$, then $(\mathbf{Q}_q, \mathbf{R}(\mathbf{Q}_q)) \xrightarrow{d} (\mathbf{M}, \mathbf{M})$ as $q \rightarrow \infty$. Since we also know $(\mathbf{R}(\mathbf{Q}_q), \mathbf{S}(\mathbf{Q}_q)) \xrightarrow{d} (\mathbf{M}, \mathbf{M})$ as $q \rightarrow \infty$, Theorem 1.3 follows immediately.

2. PRELIMINARIES

2.1. Hausdorff and Prokhorov distances. Let (V, d) be a compact metric space with its Borel σ -algebra $\mathcal{B}(V)$. Given $A \subset V$, the ε -neighborhood of A is defined as

$$A^\varepsilon = \{x \in V : \exists y \in A, d(x, y) < \varepsilon\} .$$

The *Hausdorff distance* $d_H : V^2 \rightarrow [0, \infty)$ between two non-empty subsets X, Y of (V, d) is defined as

$$d_H(X, Y) = \inf \{ \varepsilon > 0 : X \subset Y^\varepsilon, Y \subset X^\varepsilon \} .$$

Denote by $\mathcal{P}(V)$ the collection of all probability measures on the measurable space $(V, \mathcal{B}(V))$. The *Prokhorov distance* $d_P : \mathcal{P}(V)^2 \rightarrow [0, \infty)$ between two Borel probability measures μ and ν on V is given by

$$d_P(\mu, \nu) = \inf (\varepsilon > 0 : \mu(A) \leq \nu(A^\varepsilon) + \varepsilon \text{ and } \nu(A) \leq \mu(A^\varepsilon) + \varepsilon, \forall A \in \mathcal{B}(V)) .$$

2.2. Gromov-Hausdorff(-Prokhorov) distance. We refer the reader to [5] and [11, 14] for more details on the Gromov-Hausdorff and Gromov-Hausdorff-Prokhorov distances and the topologies they induce. Let (V, d) and (V', d') be two compact metric spaces. A *correspondence* between V and V' is a set $C \subset V \times V'$ such that for every $x \in V$, there is $x' \in V'$ with $(x, x') \in C$, and vice versa. We write $C(V, V')$ for the set of correspondences between V and V' . The *distortion* of any set $C \subset V \times V'$ with respect to d and d' is given by

$$\text{dis}(C; d, d') = \sup (|d(x, y) - d'(x', y')| : (x, x') \in C, (y, y') \in C) .$$

The *Gromov-Hausdorff distance* between (V, d) and (V', d') is defined as

$$d_{\text{GH}}((V, d), (V', d')) = \inf (\varepsilon > 0 : \exists C \in C(V, V'), \text{dis}(C; d, d') \leq 2\varepsilon) .$$

Next, suppose μ and μ' are non-negative Borel measures on (V, d) and (V', d') , respectively. The *Gromov-Hausdorff-Prokhorov distance* between $V = (V, d, \mu)$ and $V' = (V', d', \mu')$ is given by

$$d_{\text{GHP}}(V, V') = \inf (\max \{d_{\text{H}}(\phi(V), \phi'(V')), d_P(\phi_*\mu, \phi'_*\mu')\})$$

where the infimum is taken over all isometries ϕ, ϕ' from V, V' into a metric space (Z, δ) (see Miermont [13, Section 6.2]). Writing \mathbb{K} for the set of all isometry classes of compact metric spaces, $(\mathbb{K}, d_{\text{GHP}})$ is a Polish space; when we refer to GHP convergence we mean convergence in this space.

2.3. The Airy distribution. The *Airy distribution* is the probability distribution whose density is

$$\begin{aligned} \mathcal{A}(x) &= 2e^{-2x^3/3} (x\text{Ai}(x^2) - \text{Ai}'(x^2)) \\ &= \frac{1}{\pi x} \sum_{n \geq 1} (-x3^{2/3})^n \frac{\Gamma(2n/3 + 1)}{n!} \sin(-2n\pi/3) , \end{aligned}$$

where the Airy function Ai is given by

$$\begin{aligned} \text{Ai}(z) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(z t + t^3/3)} dt \\ &= \frac{1}{\pi 3^{2/3}} \sum_{n \geq 0} (3^{1/3} z)^n \frac{\Gamma((n+1)/3)}{n!} \sin(2(n+1)\pi/3) . \end{aligned}$$

3. COMPOSITION SCHEMATA

Let $F(z) = \sum_{n \geq 0} F_n z^n$ be a generating function with positive radius of convergence $r = r_F$. We say F is *singular with exponent 3/2* if the following properties hold.

- There exists $\varepsilon > 0$ such that F is continuable in $\Delta = \{z : |z| < r + \varepsilon, z \notin [r, r + \varepsilon)\}$.
- There exist positive constants $a = a_F, b = b_F, c = c_F$ such that $F(z) = a - b(1 - z/r) + c(1 - z/r)^{3/2} + O((1 - z/r)^2)$ as $z \rightarrow r$ in Δ .

Gao and Wormald [9] derived asymptotics for the coefficients of F under the above conditions.

Proposition 3.1 ([9], Theorem 1 (iii)). *Let F be singular with exponent 3/2, let r and c be as above. Then*

$$F_n \sim \frac{3c}{4\pi^{1/2}} \frac{r^{-n}}{n^{5/2}} .$$

Next, let C and H be generating functions with positive coefficients, and define a bivariate generating function M by $M(z, u) = C(uH(z))$; Banderier, Flajolet, Schaeffer & Soria [3] call this a *composition schema*. We generically write $C_k = [z^k]C(z)$ and $M_n = [z^n]M(z, 1)$, and for $n \geq 1$ let X_n be a real random variable with law given by

$$\mathbb{P}(X_n = k) = \frac{C_k}{M_n} [z^n]H(z)^k.$$

We quote from [3]:

Combinatorially, this corresponds to a composition $\mathcal{M} = \mathcal{C} \circ \mathcal{H}$ between classes of [rooted] objects, where objects of type \mathcal{H} are substituted freely at individual “atoms” (e.e., nodes, edges, or faces) of elements of \mathcal{C} ... $[z^n u^k]M(z, u)$ gives the number of \mathcal{M} -objects of total size n whose \mathcal{C} -component (the “core”) has size k , and X_n is the corresponding random variable describing core-size in this general context.

More precisely, X_n is the law of the size of the \mathcal{C} -component containing the root, in an object chosen uniformly at random from among all \mathcal{M} -objects of size n . The connection with the bijections for quadrangulations described in Section 1.1 should be clear. We say the triple (M, C, H) is a *map schema* if C and H are both singular with exponent $3/2$ and additionally $H(r_H) = r_C$.¹ The latter condition heuristically states that the singularity of $M(z) = M(z, 1)$ closest to the origin is caused by the singularity of C , rather than by a failure of the implicit function theorem/the Lagrange inversion theorem. The following results are all from [3].

Proposition 3.2 ([3], Theorems 1 and 5). *Suppose (M, C, H) is a map schema with*

$$C(z) = c_0 - c_1(1 - z/r_C) + c_{3/2}(1 - z/r_C)^{3/2} + O((1 - z/r_C)^2),$$

$$H(z) = h_0 - h_1(1 - z/r_H) + h_{3/2}(1 - z/r_H)^{3/2} + O((1 - z/r_H)^2).$$

Let $\alpha = \alpha_{(M, C, H)}$, $\beta = \beta_{(M, C, H)}$ and $\gamma = \gamma_{(M, C, H)}$ are defined by

$$\alpha = \frac{c_1 h_{3/2} h_0^{1/2} + c_{3/2} h_1^{3/2}}{h_0^{3/2}}, \beta = \frac{h_1^{5/3}}{(3h_{3/2})^{2/3} h_0}, \gamma = \beta \cdot \frac{c_{3/2} h_1^{3/2}}{\alpha \cdot h_0^{3/2}}.$$

Then

$$[z^n]M(z, 1) \sim \frac{3\alpha}{4\pi^{1/2}} \frac{r_H^{-n}}{n^{5/2}}.$$

Furthermore, writing $\alpha_0 = \alpha_{0, (M, C, H)} = h_0/h_1$, for any $A > 0$,

$$\lim_{n \rightarrow \infty} \sup_{x \in [-A, A]} |n^{2/3} \mathbb{P}(X_n = \lfloor \alpha_0 n + xn^{2/3} \rfloor) - \gamma \mathcal{A}(\beta x)| = 0. \quad (3.1)$$

Finally, there exist continuous functions $f : [\alpha_0, 1] \rightarrow (0, \infty)$ and $g : [\alpha_0, 1] \rightarrow (0, \infty)$ such that for any function $\lambda : \mathbb{N} \rightarrow \mathbb{N}$ with $\lambda(n) \rightarrow \infty$,

$$\mathbb{P}(X_n = k) = (1 + o(1)) f(k/n) \frac{(k/n - \alpha_0)^{1/2}}{n^{1/2} (1 - k/n)^{3/2}} e^{-n(k/n - \alpha_0)^3 g(k/n)}, \quad (3.2)$$

the preceding asymptotic holding uniformly in $\alpha_0 n + n^{2/3} \lambda(n) < k < n - n^{2/3} \lambda(n)$.

Remark. In [3], schema of the form $M(z, u) = C(uH(z)) + D(z)$ are also considered. Replacing $M(z, u)$ by $M(z, u) - D(z)$ turns this into a compositional schema; if the latter is a map schema then Proposition 3.2 applies. The equation involving D is convenient when considering map families in which the core may have size zero; such families should be counted by $[u^0]M(z, u)$, which is identically zero in $C(uH(z))$.

¹In [3], this is called a *critical composition schema of singular type* $(\frac{3}{2} \circ \frac{3}{2})$. We shorten this to “map schema” as such schemata seem to primarily arise in the study of maps.

Corollary 3.3. *Suppose (M, C, H) is a map schema, and let $\alpha_0 = \alpha_{0,(M,C,H)}$, f and g be as in Proposition 3.2. Then for any function $\lambda : \mathbb{N} \rightarrow \mathbb{N}$ with $\lambda(n) \rightarrow \infty$ and any $a > 0$,*

$$\mathbb{P}(X_n = k) = \Theta(1) \cdot \frac{(k/n - \alpha_0)^{1/2}}{n^{1/2}(1 - k/n)^{3/2}} e^{-n(k/n - \alpha_0)^3 g(k/n)},$$

uniformly over integers $k \in [\alpha_0 n + an^{2/3}, n - \lambda(n)n^{2/3}]$.

Proof. Note that if $k = \alpha_0 n + cn^{2/3}$ then

$$\frac{(k/n - \alpha_0)^{1/2}}{n^{1/2}(1 - k/n)^{3/2}} e^{-n(k/n - \alpha_0)^3 g(k/n)} = \frac{c^{1/2}}{n^{2/3}(1 - k/n)^{3/2}} e^{-c^3 g(\alpha_0 + c/n^{1/3})}.$$

For $|k - \alpha_0 n| = O(n^{2/3})$, the latter is $\Theta(n^{-2/3})$. By (3.1), we indeed have $\mathbb{P}(X_n = k) = \Theta(n^{-2/3})$ for such k .

If the claim of the corollary fails then there exists a sequence $(n_i, i \geq 1)$ and $k_i \in [\alpha_0 n_i + an_i^{2/3}, n_i - \lambda(n_i)n_i^{2/3}]$ along which the ratio of $\mathbb{P}(X_{n_i} = k_i)$ and

$$\frac{(k_i/n_i - \alpha_0)^{1/2}}{n_i^{1/2}(1 - k_i/n_i)^{3/2}} e^{-n_i(k_i/n_i - \alpha_0)^3 g(k_i/n_i)}$$

either diverges or tends to zero. By passing to a subsequence if necessary, we may assume that either $k_i - \alpha_0 n_i = O(n_i^{2/3})$ or $(k_i - \alpha_0 n_i)/n_i^{2/3} \rightarrow \infty$. In view of the above computation, the first possibility is in contradiction with (3.1). The second possibility is in contradiction with (3.2); thus neither can occur. \square

Corollary 3.4 ([3], Theorem 7). *Suppose (M, C, H) is a map schema with $\alpha_0 = \alpha_{0,(M,C,H)}$ and $\beta = \beta_{(M,C,H)}$ defined in Proposition 3.2. Let X_n^* be the size of the largest C -component in a random M -map of size n with uniform distribution. Then*

$$\mathbb{P}\left(X_n^* = \lfloor \alpha_0 n + xn^{2/3} \rfloor\right) = \frac{\beta \mathcal{A}(\beta x)}{n^{2/3}} (1 + o(1)),$$

uniformly for x in any bounded interval.

Let $M(z), C(z), B(z)$ be the generating functions of rooted connected, 2-connected, and simple quadrangulations respectively. More precisely, we take $[z^n]M(z) = |\mathcal{Q}_{n+2}|$, $[z^n]C(z) = |\mathcal{R}_{n+2}|$, and $[z^n]B(z) = |\mathcal{S}_{n+2}|$ for $n \geq 1$, and $[z^n]M(z) = [z^n]C(z) = [z^n]B(z) = 0$ for $n = 0$. (The latter is slightly at odds with our convention of viewing a single edge as a 2-connected quadrangulation, but is algebraically convenient below.) Define

$$H(z) = z \left(\frac{1}{1 - 2z(1 + M(z))} \right)^2, \quad (3.3)$$

$$U(z) = z(1 + C(z))^2. \quad (3.4)$$

The following two lemmas follow immediately from Propositions 1.6 and 1.7 respectively.

Lemma 3.5. *We have the following substitution relation between $M(z)$ and $C(z)$:*

$$M(z) = C(H(z)) \cdot \frac{1}{1 - 2z(1 + M(z))} + \frac{2z(1 + M(z))}{1 - 2z(1 + M(z))}. \quad (3.5)$$

Equivalently,

$$M(z) = C(H(z)) + 2z(1 + M(z))^2. \quad (3.6)$$

(3.6) is obtained by multiplying both sides of (3.5) by $1 - 2z(1 + M(z))$ and then rearranging elements. To see that (3.6) gives a composition schema, we can rewrite it as $\hat{M}(z) = C(H(z))$ with $\hat{M}(z) = M(z) - 2z(1 + M(z))^2$.

We now take a closer look at equation (3.3), which describes the “ M -decorations” of an edge of a C -object (i.e. of a 2-connected map). This is best understood with

the bijection from Proposition 1.6 at hand. In the term $2z(1 + M(z))$, the multiplier 2 counts the choice of extremity at which the decoration is attached; $M(z)$ counts the case when attachment is a quadrangulation with at least 3 vertices (recalling that z marks the number of vertices less two, and the lowest power term of $M(z)$ is $2z$); the additive term 1 counts the case when attachment is a single edge; the multiplier z adjusts the counting of extra vertices resulting from the attachment (we multiply by z instead of z^2 because the attachment vertex is already counted in the core). Taking the reciprocal of $1 - 2z(1 + M(z))$ accounts for the fact that we can attach a sequence of submaps (each two separated by an edge). Squaring the reciprocal accounts for the fact that in a quadrangulation Q we have $|e(Q)| = 2(|v(Q)| - 2)$.

In equation (3.5), the term $\frac{2z(1+M(z))}{1-2z(1+M(z))}$ takes into consideration when the root block is a single edge. The multiplication $\frac{1}{1-2z(1+M(z))}$ in the first term accounts for the extra submap attachment due to split of the root edge (recall the construction preceding Proposition 1.6).

Lemma 3.6. *We have the following substitution relation between $C(z)$ and $B(z)$:*

$$C(z) = B(U(z)) \cdot (1 + C(z)) . \quad (3.7)$$

To see that this identity gives a composition schema, note that it may equivalently be written as $C(z) = \hat{B}(U(z))$ with $\hat{B} = B/(1 - B)$. The multiplication $(1 + C(z))$ accounts for the extra submap attachment due to the split of the root edge (see the construction preceding Proposition 1.7).

4. AIRY DISTRIBUTION FOR QUADRANGULATIONS

In this section, we describe composition schemata for both rooted connected and 2-connected quadrangulations, and then establish corresponding distributional results.

Proposition 4.1. *The triple (M, C, H) is a map schema with*

$$\alpha_0 = \frac{7}{15}, \quad \alpha = \frac{40}{27}, \quad \beta = \frac{5^{2/3} \cdot 15}{28}, \quad \gamma = \frac{9}{5^{1/3} \cdot 4} . \quad (4.1)$$

Proposition 4.2. *The triple (C, B, U) is a map schema with*

$$\alpha_0 = \frac{5}{7}, \quad \alpha = \frac{21^{1/2} \cdot 9}{5^{1/2} \cdot 50}, \quad \beta = \frac{3^{2/3} \cdot 45}{14^{1/3} \cdot 8}, \quad \gamma = \frac{3^{2/3} \cdot 225}{14^{1/3} \cdot 56} . \quad (4.2)$$

These two map schemata lead to the Airy distributional results stated in Propositions 1.8 and 4.3.

Proof of Proposition 1.8. This now follows immediately from Corollary 3.4 and Proposition 4.2. \square

We will also need the following analogue of Proposition 1.8 for the largest 2-connected block of a general quadrangulation.

Proposition 4.3. *Let $\mathbf{Q}_q \in_u \mathcal{Q}_q$, then for any $A > 0$, uniformly over $x \in [-A, A]$,*

$$\mathbb{P} \left(\mathfrak{b}(\mathbf{Q}_q) = \left\lfloor 7q/15 + xq^{2/3} \right\rfloor \right) = \frac{\beta \mathcal{A}(\beta x)}{q^{2/3}} (1 + o(1)) ,$$

where β is given in Proposition 4.1.

Proof. This follows from Corollary 3.4 and Proposition 4.1. \square

Before proving Propositions 4.1 and 4.2, we first establish a system of parameterization for $M(z)$, which is the key to showing that $M(z)$ is singular with exponent $3/2$ and to extracting the coefficients of $M(z)$, $C(z)$ and $H(z)$.

Lemma 4.4. *Let $\psi_M(t) = \frac{t(2-9t)}{(1-3t)^2}$, let $\phi_M(t) = \frac{1}{1-3t}$, and let $L_M(z)$ be defined by the implicit relation $L_M(z) = z\phi_M(L_M(z))$, then*

$$M(z) = \psi_M(L_M(z)) .$$

Algebraic functions with such parameterization are called *Lagrangean*. The proof is a textbook application of the so-called *quadratic method*. A Lagrangean parametrization for M is derived in [3, Proposition 1]. The parameterization in Lemma 4.4 is slightly different, but the derivation is quite the same, and refer readers to that work the idea of the proof. (Also, in [3] the parameterization is stated for the generating function of general maps but this is equivalent, using Tutte's angular bijection, to quadrangulations. See also Goulden & Jackson [10, Section 2.9] for a detailed explanation of the quadratic method for map enumeration.) One may inspect the Taylor expansion of $\psi_M(L_M(z))$ at $z = 0$ to conclude that this parametrization gives $M(z) = 2z + 9z^2 + 54z^3 + 378z^4 + O(z^5)$.

Corollary 4.5 (Tutte [16]).

$$L_M(z) = \frac{1}{6} - \frac{1}{6}(1 - 12z)^{1/2} , \quad (4.3)$$

$$M(z) = \frac{1}{3} - \frac{4}{3}(1 - 12z) + \frac{8}{3}(1 - 12z)^{3/2} + O((1 - 12z)^2) . \quad (4.4)$$

In particular, $M(z)$ is singular with exponent $3/2$.

Proof. Using Lemma 4.4, Lagrange inversion yields the explicit formulas

$$L_M(z) = \frac{1}{6} - \frac{1}{6}(1 - 12z)^{1/2} , \quad M(z) = -1 + \frac{1}{54z^2}(-1 - 18z) + (1 - 12z)^{3/2} .$$

Writing $y = 1 - 12z$, the asymptotic expansion for M follows easily by rearrangement. \square

Implicit functional equations can be used to derive asymptotic expansions in great generality, even when no closed form is available, and we exploit this machinery in the current paper. We now sketch how the method is applied in our setting in slightly more detail, referring the reader to [3] and [8, VII.8] for a full exposition.

Suppose we are given y defined by an implicit formula $y(z) = z\phi(y(z))$, where ϕ is analytic, non-zero at 0, has non-negative Taylor coefficients, and has $\lim_{x \rightarrow r_y} x\phi'(x)/\phi(x) > 1$, where $r_y \in (0, \infty]$ is the radius of convergence of ϕ . (In our case, ϕ will always in fact be a *rational* function satisfying the preceding conditions.) Then, using Lagrange inversion, one obtains an asymptotic expansion of y around its dominant singularity (see [8, Section VI.7]). Given another function m expressible as $m(z) = \psi(y(z))$ where ψ is a rational function whose radius of convergence is at least as large as that of y , this yields an asymptotic expansion for m as follows.

First, we locate the radius of convergence for y . By [8, Theorem VI.6], we can expand $y(z)$ as

$$y(z) = \tau - l_{1/2}(1 - z/r_y)^{1/2} + l_1(1 - z/r_y) + O((1 - z/r_y)^{3/2}) , \quad (4.5)$$

where the coefficients $l_{i/2}$ are to be determined for $i \in \mathbb{N}$, and r_y and τ are determined by the equations

$$\tau\phi'(\tau) - \phi(\tau) = 0, \quad r_y = \frac{\tau}{\phi(\tau)} .$$

To determine $l_{1/2}$ and l_1 , let $h(t) = r_y - \frac{t}{\phi(t)}$. Then $h(\tau) = 0 = h'(\tau)$, so expanding $h(t)$ around τ yields

$$\begin{aligned} & 1 - z/r_y \\ &= \frac{h(t)}{r_y} \end{aligned}$$

$$\begin{aligned}
&= \frac{h''(\tau)}{2r_y} (t - \tau)^2 + \frac{h'''(\tau)}{6r_y} (t - \tau)^3 + O((t - \tau)^4) \\
&= \frac{h''(\tau)}{2r_y} \left[-l_{1/2} (1 - z/r_y)^{1/2} + l_1 (1 - z/r_y) + O\left((1 - z/r_y)^{3/2}\right) \right]^2 \\
&\quad + \frac{h'''(\tau)}{6r_y} \left[-l_{1/2} (1 - z/r_y)^{1/2} + l_1 (1 - z/r_y) + O\left((1 - z/r_y)^{3/2}\right) \right]^3 + O((t - \tau)^4) \\
&= \frac{h''(\tau)}{2r_y} l_{1/2}^2 (1 - z/r_y) - \left(2 \frac{h''(\tau)}{2r_y} l_{1/2} l_1 + \frac{h'''(\tau)}{6r_y} l_{1/2}^3 \right) (1 - z/r_y)^{3/2} + O\left((1 - z/r_y)^2\right).
\end{aligned}$$

By comparing the coefficients of the terms $(1 - z/r_y)$ we obtain

$$l_{1/2} = \left(\frac{2r_y}{h''(\tau)} \right)^{1/2} = \left(\frac{2\phi(\tau)}{\phi''(\tau)} \right)^{1/2}, \quad (4.6)$$

and by comparing the coefficients of the terms $(1 - z/r_y)^{3/2}$ we have

$$l_1 = -\frac{h'''(\tau) l_{1/2}^3}{6h''(\tau) l_{1/2}}. \quad (4.7)$$

Now we use the expansion (4.5) to derive an expansion for $m(z)$ around its dominant singularity r_m . First, the equation $m(z) = \psi(y(z))$ and the assumption that $r_\phi \geq r_y$ together imply that $r_m = r_y$. In the current work, we always have that $\psi'(\tau) = 0$ (indeed, this seems to generally be the case in compositional schemata involving maps); together with (4.5), a Taylor expansion of ψ around τ then yields

$$\begin{aligned}
m(z) &= \psi(y(z)) \\
&= \psi \left(\tau - l_{1/2} (1 - z/r_y)^{1/2} + l_1 (1 - z/r_y) + O\left((1 - z/r_y)^{3/2}\right) \right) \\
&= \psi(\tau) + \frac{1}{2} \psi''(\tau) \left[-l_{1/2} (1 - z/r_y)^{1/2} + l_1 (1 - z/r_y) + O\left((1 - z/r_y)^{3/2}\right) \right]^2 \\
&\quad + \frac{1}{6} \psi'''(\tau) \left[-l_{1/2} (1 - z/r_y)^{1/2} + l_1 (1 - z/r_y) + O\left((1 - z/r_y)^{3/2}\right) \right]^3 \\
&\quad + O\left((1 - z/r_y)^2\right) \\
&= \psi(\tau) + \frac{1}{2} \psi''(\tau) l_{1/2}^2 (1 - z/r_y) - \left(\psi''(\tau) l_{1/2} l_1 + \frac{1}{6} \psi'''(\tau) l_{1/2}^3 \right) (1 - z/r_y)^{3/2} \\
&\quad + O\left((1 - z/r_y)^2\right).
\end{aligned}$$

We remark that the vanishing term $\psi'(\tau) = 0$ accounts for the shift of the singular exponent to $3/2$.

Using the compositional relation given in Lemma 3.5 together with the expansion of $L_M(z)$ given in Corollary 4.5, we show that $H(z)$ is also Lagrangean and expand $H(z)$ at its radius of convergence. We similarly deduce such kind of expansions for $C(z)$, $U(z)$ and $B(z)$.

Lemma 4.6. *Table 1 gives Lagrangean parameterizations for $M(z)$, $H(z)$, $C(z)$, $U(z)$, $B(z)$.*

Consequently, $H(z)$, $C(z)$, $U(z)$, and $B(z)$ each has radius of convergence and asymptotic expansion around r_H , r_C , r_U and r_B as given in Table 2.

Proof. Let $H(z)$ be defined as in Proposition 4.1, and let $\psi_M(t)$ be given in Lemma 4.4. Write $t = L_M(z)$, then

$$H(z) = z \left(\frac{1}{1 - 2z(1 + M(z))} \right)^2 = z \left(\frac{1}{1 - 2z(1 + \psi_M(L_M(z)))} \right)^2 = -\frac{t(-1 + 3t)^3}{(1 - 5t + 8t^2)^2};$$

this proves the first assertion. Then taking $\psi_H(t)$ as given by Table 1 yields $H(z) = \psi_H(L_M(z))$.

The remaining parameterizations of Table 1 are established similarly, using (3.5) for $C(z)$, and (3.7) for $U(z)$ and $B(z)$. The radius of convergence and expansions around the radius are derived using Lagrange inversion as in Corollary 4.5. \square

f	ϕ_f	ψ_f
H	$\frac{1}{1-3t}$	$-\frac{t(-1+3t)^3}{(1-5t+8t^2)^2}$
C	$-\frac{(1-5t+8t^2)^2}{(-1+3t)^3}$	$\frac{t^2(-1+5t)}{(-1+3t)^3}$
U	$-\frac{(1-5t+8t^2)^2}{(-1+3t)^3}$	$-\frac{t(-1+4t)^2}{(-1+3t)^3}$
B	$-\frac{(-1+3t)^3}{(-1+4t)^2}$	$\frac{t^2(-1+5t)}{(-1+4t)(1-5t+8t^2)}$

TABLE 1. (Always with $L_f(z) = z\phi_f(L_f(z))$, where f is one of the functions H, C, U , and B .)

f	r_f	expansion at r_f
H	1/12	$\frac{27}{196} - \frac{405}{1372}(1-12z) + \frac{54}{343}(1-12z)^{3/2} + O((1-12z)^2)$
C	27/196	$\frac{1}{27} - \frac{28}{135}(1-196z/27) + \frac{392}{675}\sqrt{\frac{7}{15}}(1-196z/27)^{3/2} + O((1-196z/27)^2)$
U	27/196	$\frac{4}{27} - \frac{28}{135}(1-196z/27) + \frac{112}{675}\sqrt{\frac{7}{15}}(1-196z/27)^{3/2} + O((1-196z/27)^2)$
B	4/27	$\frac{1}{28} - \frac{27}{196}(1-27z/4) + \frac{9\sqrt{3}}{98}(1-27z/4)^{3/2} + O((1-27z/4)^2)$

TABLE 2

Remark. One of the fundamental facts of singularity analysis is that the radius of convergence of a generating function determines the exponential growth rate of the associated combinatorial family. Under Tutte’s angular bijection (see [16]), 2-connected and simple quadrangulations respectively correspond to 2-edge-connected and 2-connected maps. In view of this, the values $r_C = 27/196$ and $r_B = 4/27$ agree with the known exponential growth rates for loopless bridgeless maps Walsh & Lehman [17, (7)] and for 2-connected maps [3, Table 2] (noting that the coefficients of the expansion for $B(z)$ are slightly different than in [3], because in that work a single loop is counted as a 2-connected map).

Proofs of Propositions 4.1 and 4.2. We have verified that $M(z)$ and $H(z)$ are singular with exponent 3/2 in Lemmas 4.5 and 4.6 respectively. The facts that $H(r_H) = r_C$ and that $U(r_U) = r_B$ are immediate from the values and expansions given in Table 2. Thus, (M, C, H) and (C, B, U) are map schemata. The values claimed in (4.1) and (4.2) are then derived by routine arithmetic. \square

5. SIZES AND DIAMETERS OF PENDANT SUBMAPS

In this section, we first obtain a size bound for the decorations of the largest simple block in a uniform rooted 2-connected quadrangulation. Using this we then derive a corresponding diameter bound which leads to a “GH convergence version” of Proposition 1.5, shown in Section 5.1. Analogous tail bounds for uniform rooted quadrangulations are stated in Section 5.2.

Proof of Proposition 1.9. Let $\lambda : \mathbb{N} \rightarrow \mathbb{N}$ be a function tending to infinity with $\lambda(r) \leq \frac{r^{1/3}}{(\log r)^2}$. For $k \leq r - r^{2/3}\lambda(r)$, the bound follows straightforwardly from Proposition 1.8, Corollary 3.3, and the fact that the largest block is larger than or equal in size to the root block. We hereafter assume that $r - r^{2/3}\lambda(r) < k \leq r$. Note that for r large enough, $r - r^{2/3}\lambda(r) > r/2$, so there must be less than one largest simple block of size k .

Let $\mathbf{R}_r \in_u \mathcal{R}_r$, and note that $\mathbb{P}(\text{sb}(\mathbf{R}_r) = k) = \frac{|\mathcal{R}_{r,k}|}{|\mathcal{R}_r|}$. By Proposition 3.1 and Lemma 4.6, $|\mathcal{R}_r| = [z^{r-2}]C(z) = \Theta(1) \cdot r^{-5/2} r_C^{-r}$ as $r \rightarrow \infty$ with $r_C = \frac{27}{196}$.

Each element of $\mathcal{R}_{r,k}$ may be constructed by first choosing $S \in \mathcal{S}_k$ and a collection $(M_e, e \in e(S))$ of rooted 2-connected quadrangulations, each with at most k vertices, and with $\sum_{e \in e(S)} |e(M_e)| = 2(r - k)$; then attaching each M_e to $e \in e(S)$ to obtain a map R with r edges; and, finally, fixing a root edge \bar{e} in R . This builds a map $(R, \bar{e}) \in \mathcal{R}_{r,k}$, and any element of $\mathcal{R}_{r,k}$ may be so built. It follows that

$$|\mathcal{R}_{r,k}| \leq (4r - 8) \cdot (2k - 4)^{r-k} \cdot |\mathcal{S}_k| \cdot |\mathcal{R}_{r-k}|,$$

where $(4r - 8)$ accounts for the re-assignment of the oriented root edge, and $|\mathcal{S}_k| = [z^{k-2}]B(z)$, $|\mathcal{R}_{r-k}| = [z^{r-k-2}]C(z)$. So

$$|\mathcal{R}_{r,k}| \leq (4r - 8) \cdot (2k - 4)^{r-k} \cdot k^{-5/2} \cdot r_B^{-k} \cdot (r - k)^{-5/2} \cdot r_C^{-r+k},$$

where r_B and r_C appear in Table 2.

Altogether, for $r - r^{2/3}\lambda(r) < k \leq r$,

$$\mathbb{P}(\text{sb}(\mathbf{R}_r) = k) = \frac{|\mathcal{R}_{r,k}|}{|\mathcal{R}_r|} \leq \Theta(r) \cdot (2k - 4)^{r-k} \cdot k^{-5/2} \cdot (r - k)^{-5/2} \cdot r^{5/2} \cdot \left(\frac{r_C}{r_B}\right)^k.$$

For large enough r we have $r - k < \lambda(r)r^{2/3} \leq \frac{r}{(\log r)^2}$, so for such r ,

$$(2k - 4)^{r-k} = \exp((r - k) \cdot \log(2k - 4)) \leq \exp\left(\frac{r}{(\log r)^2} \cdot \log(2k - 4)\right) \leq \exp\left(\frac{r}{\log r}\right).$$

We have $\frac{r_C}{r_B} < 1$, so there exists $b > 0$ such that $\frac{r_C}{r_B} \leq \exp(-b)$. It follows that for some positive constants c_1 and c_2 ,

$$\begin{aligned} \mathbb{P}(\text{sb}(\mathbf{R}_r) = k) &\leq (4r - 8) \cdot k^{-5/2} \cdot (r - k)^{-5/2} \cdot r^{5/2} \cdot \exp\left(-b \cdot k + \frac{r}{\log r}\right) \\ &\leq c_1 \exp(-c_2 r (k/r - 5/7)^3). \end{aligned} \quad \square$$

Proof of Proposition 1.10. For all positive integers r and x with $x \leq r - s(r) + 2$ write

$$\mathcal{L}_{r,x} = \{\mathbf{R} \in \mathcal{R}_{r,s(r)} : L(\mathbf{R}) = x\}.$$

Fix $\varepsilon \in (0, 1/3)$ for the remainder of the proof. Letting $\mathbf{R}_r \in_u \mathcal{R}_{r,s(r)}$,

$$\mathbb{P}\left(L(\mathbf{R}_r) \geq r^{2/3+\varepsilon}\right) = |\mathcal{R}_{r,s(r)}|^{-1} \sum_{x=\lceil r^{2/3+\varepsilon} \rceil}^{r-s(r)+2} |\mathcal{L}_{r,x}| \quad (5.1)$$

Since $s(r) = 5r/7 + O(r^{2/3})$ as $r \rightarrow \infty$, by Proposition 1.8,

$$|\mathcal{R}_{r,s(r)}| = \Theta\left(r^{-2/3}\right) \cdot |\mathcal{R}_r| = \Theta\left(r^{-2/3}\right) \cdot r^{-5/2} \cdot r_C^{-r} = \Theta\left(r^{-19/6}\right) \cdot r_C^{-r}. \quad (5.2)$$

Thus, it remains to bound $|\mathcal{L}_{r,x}|$.

Each element of $\mathcal{L}_{r,x}$ can be obtained by attaching some $R_x \in \mathcal{R}_x$ to the largest simple block of some $\mathbf{R} \in \mathcal{R}_{r-x+2,s(r)}$ with $\text{sb}(\mathbf{R}) \leq x$, then possibly re-assigning the root edge. We therefore have

$$|\mathcal{L}_{r,x}| \leq \Theta(r \cdot s(r)) \cdot |\mathcal{R}_{r-x+2,s(r)}| \cdot |\mathcal{R}_x| \quad (5.3)$$

as $r \rightarrow \infty$. Then to bound $|\mathcal{L}_{r,x}|$, it suffices to bound $|\mathcal{R}_{r-x+2,s(r)}|$ and $|\mathcal{R}_x|$. For large enough r and for all $x \in [r^{2/3+\varepsilon}, r - s(r) + 2]$, we have $5(r-x+2)/7 + (r-x+2)^{2/3} \leq s(r) \leq r-x+2$. For x in this range, we may apply Proposition 1.9: we obtain that for some $C' > 0$,

$$\frac{|\mathcal{R}_{r-x+2,s(r)}|}{|\mathcal{R}_{r-x+2}|} = O(1) \cdot \exp\left(-C'(r-x) \left(\frac{s(r)}{r-x} - 5/7\right)^3\right).$$

For all possible x , by Proposition 3.2 and Lemma 4.6 we have $|\mathcal{R}_{r-x+2}| = \Theta(1) \cdot (r-x)^{-5/2} r_C^{-r+x}$, so

$$|\mathcal{R}_{r-x+2,s(r)}| = O(1) \cdot (r-x)^{-5/2} \cdot r_C^{-r+x} \cdot \exp\left(-C'(r-x) \left(\frac{s(r)}{r-x} - 5/7\right)^3\right).$$

Then (5.3) gives

$$|\mathcal{L}_{r,x}| = O(r \cdot s(r)) \cdot x^{-5/2} \cdot (r-x)^{-5/2} \cdot r_C^{-r} \cdot \exp\left(-C'(r-x) \left(\frac{s(r)}{r-x} - 5/7\right)^3\right).$$

Since $\frac{s(r)}{r} \geq 5/7 - Cr^{-1/3}$, we have for large r ,

$$\exp\left(-C'(r-x) \left(\frac{s(r)}{r-x} - 5/7\right)^3\right) \leq \exp\left(-C'(r-x)^{-2} \left(5x/7 - Cr^{2/3}\right)^3\right).$$

For $r^{2/3+\varepsilon} \leq x \leq r - s(r)$, and for large enough r , we thus have

$$\begin{aligned} & |\mathcal{L}_{r,x}| \\ &= O(r \cdot s(r)) \cdot x^{-5/2} \cdot (r-x)^{-5/2} \cdot r_C^{-r} \cdot \exp\left(-C'(r-x)^{-2} \left(5x/7 - Cr^{2/3}\right)^3\right) \\ &= O(r \cdot s(r)) \cdot x^{-5/2} \cdot (r-x)^{-5/2} \cdot r_C^{-r} \cdot \exp\left(-C' \left(r - r^{2/3+\varepsilon}\right)^{-2} \left(5r^{2/3+\varepsilon}/7 - Cr^{2/3}\right)^3\right) \\ &= r_C^{-r} \cdot \exp(-C'' \cdot r^{3\varepsilon}), \end{aligned} \tag{5.4}$$

for some $C'' > 0$.

Finally, combining (5.1), (5.2), (5.4) and the fact that $s(r) = 5r/7 + O(r^{2/3})$, there exist positive constants c_1, c_2 and $c_3 = c_3(\varepsilon)$ such that

$$\mathbb{P}\left(L(\mathbf{R}_r) \geq r^{2/3+\varepsilon}\right) = |\mathcal{R}_{r,s(r)}|^{-1} \sum_{x=[r^{2/3+\varepsilon}]^{r-s(r)+2}} |\mathcal{L}_{r,x}| \leq c_1 \exp(-c_2 r^{c_3}). \quad \square$$

5.1. Diameters of submaps pendant to the largest simple block. We want to apply [6] to obtain a diameter bound, but first we need to transfer the diameter tail bound from [6] to the setting of 2-connected quadrangulations.

Lemma 5.1. *Let $\mathbf{R}_r \in_u \mathcal{R}_r$, then there exist positive constants x_0, c_1 and c_2 such that for all $x > x_0$,*

$$\mathbb{P}\left(\text{diam}(\mathbf{R}_r) > xr^{1/4}\right) \leq c_1 r^{2/3} \exp(-c_2 x).$$

Proof. For $q \in \mathbb{N}$, let $\mathbf{Q}_q \in_u \mathcal{Q}_q$. Given that $b(\mathbf{Q}_q) = r$, $\mathbf{R}(\mathbf{Q}_q)$ has the same distribution as \mathbf{R}_r . So for all $q \geq r$ and $x > 0$, we have

$$\begin{aligned} \mathbb{P}\left(\text{diam}(\mathbf{R}_r) > xr^{1/4}\right) &= \mathbb{P}\left(\text{diam}(\mathbf{R}(\mathbf{Q}_q)) > xr^{1/4} \mid b(\mathbf{Q}_q) = r\right) \\ &\leq \mathbb{P}\left(\text{diam}(\mathbf{Q}_q) > xr^{1/4} \mid b(\mathbf{Q}_q) = r\right) \end{aligned}$$

$$\leq \frac{\mathbb{P}(\text{diam}(\mathbf{Q}_q) > xr^{1/4})}{\mathbb{P}(\text{b}(\mathbf{Q}_q) = r)}.$$

Now let $q = \lfloor 15r/7 \rfloor$, then $xr^{1/4} \geq x(7/15)^{1/4}q^{1/4}$, so by Proposition 1.11, there exist positive constants x_0, C_1, C_2 such that for all $x > x_0$,

$$\mathbb{P}(\text{diam}(\mathbf{Q}_q) > xr^{1/4}) \leq C_1 \exp(-C_2x).$$

On the other hand, by Proposition 4.3, there exists $C_3 > 0$ such that for all $r \in \mathbb{N}$,

$$\mathbb{P}(\text{b}(\mathbf{Q}_q) = r) \geq C_3 r^{-2/3}.$$

Altogether, we have

$$\mathbb{P}(\text{diam}(\mathbf{R}_r) > xr^{1/4}) \leq \frac{C_1 \exp(-C_2x)}{C_3 r^{-2/3}}$$

Then setting $c_1 = C_1/C_3$ and $c_2 = C_2$ concludes the proof. \square

Proof of Proposition 1.12. Fix a positive integer r and let $k \in \mathbb{N}$ with $k \leq \min\{s(r), r - s(r)\}$. Let $\mathbf{R}_r = (R_r, e_r) \in_u \mathcal{R}_{r,s(r)}$, write $\mathbf{S}_r = \mathbf{S}(\mathbf{R}_r)$, let e' be the $\prec_{\mathbf{R}_r}$ -minimal oriented edge of \mathbf{S}_r , and write $\mathbf{R}'_r = (R_r, e')$. It follows from Proposition 1.6 that \mathbf{R}'_r uniquely decomposes as $(\mathbf{S}_r, \Theta) \in \mathcal{S} \times \mathcal{R}^{|\mathbf{S}(\mathbf{R}_r)|+1}$. Write $\Theta = (\Theta_0, \Theta_1, \dots, \Theta_{|\mathbf{S}(\mathbf{R}_r)|})$; recall that Θ has two entries corresponding to the root edge.

For any $0 \leq i \leq |\mathbf{S}(\mathbf{R}_r)|$, given that $|v(\Theta_i)| = k$, Θ_i is uniformly distributed over \mathcal{R}_k . By Lemma 5.1 and since $k \leq r$, there exist positive constants x_0, c_1 and c_2 such that for all $x \geq x_0$, and for all $0 \leq i \leq |\mathbf{S}(\mathbf{R}_r)|$,

$$\mathbb{P}(\text{diam}(\Theta_i) \geq xk^{1/4} \mid |v(\Theta_i)| = k) \leq c_1 r^{2/3} \exp(-c_2x). \quad (5.5)$$

Note that $|\mathbf{S}(\mathbf{R}_r)| = 2s(r) - 4$ and recall that $D(\mathbf{R}_r) = \max(\text{diam}(\Theta_i) : 0 \leq i \leq 2s(r) - 4)$. Fix $\varepsilon \in (0, 1/7)$. Using a union bound,

$$\begin{aligned} & \mathbb{P}(D(\mathbf{R}_r) \geq r^{5/24}) \\ & \leq \sum_{i=0}^{2s(r)-4} \left[\sum_{k=1}^{\lfloor r^{2/3+\varepsilon} \rfloor} \mathbb{P}(\text{diam}(\Theta_i) \geq r^{5/24}, |v(\Theta_i)| = k) + \mathbb{P}(|v(\Theta_i)| \geq r^{2/3+\varepsilon}) \right] \\ & \leq \sum_{i=0}^{2s(r)-4} \left[\sum_{k=1}^{\lfloor r^{2/3+\varepsilon} \rfloor} \mathbb{P}(\text{diam}(\Theta_i) \geq r^{5/24} \mid |v(\Theta_i)| = k) + \mathbb{P}(|v(\Theta_i)| \geq r^{2/3+\varepsilon}) \right]. \end{aligned}$$

By (5.5), for $k \leq r^{2/3+\varepsilon}$ and for each $0 \leq i \leq 2s(r) - 4$,

$$\begin{aligned} \mathbb{P}(\text{diam}(\Theta_i) \geq r^{5/24} \mid |v(\Theta_i)| = k) & \leq c_1 r^{2/3} \exp(-c_2 r^{5/24} k^{-1/4}) \\ & \leq c_1 r^{2/3} \exp(-c_2 r^{1/24 - \varepsilon/4}) \end{aligned}$$

Finally, by Proposition 1.10, there exist positive constants k_1, k_2 and $k_3 = k_3(\varepsilon)$ such that for each $0 \leq i \leq 2s(r) - 4$,

$$\mathbb{P}(|v(\Theta_i)| \geq r^{2/3+\varepsilon}) \leq \mathbb{P}(L(\mathbf{R}_r) \geq r^{2/3+\varepsilon}) \leq k_1 \exp(-k_2 r^{k_3});$$

combining the preceding 3 inequalities and using that $s(r) \leq r$ and that $1/24 - \varepsilon/4 > 1/168$ yields

$$\begin{aligned} \mathbb{P}(D(\mathbf{R}_r) \geq r^{5/24}) & \leq (2s(r) - 3) \left[r^{2/3+\varepsilon} \cdot c_1 r^{2/3} \exp(-c_2 r^{1/24 - \varepsilon/4}) + k_1 \exp(-k_2 r^{k_3}) \right] \\ & = O\left(r^{7/3+\varepsilon} \exp(-c_2 r^{1/168})\right) + O\left(r \cdot \exp(-k_2 r^{k_3})\right). \end{aligned}$$

By choosing the constants C_1, C_2 and C_3 carefully, we may conclude the proof. \square

Given the diameter bound, we immediately have the ‘‘GH convergence version’’ of Proposition 1.5:

Proposition 5.2. *Let $\mathbf{R}_r \in_u \mathcal{R}_{r,s(r)}$ and write $\mathbf{S}_r = S(\mathbf{R}_r)$, then as $r \rightarrow \infty$,*

$$\left(\left(v(\mathbf{R}_r), \left(\frac{21}{40r} \right)^{1/4} \cdot d_{\mathbf{R}_r} \right), \left(v(\mathbf{S}_r), \left(\frac{21}{40r} \right)^{1/4} \cdot d_{\mathbf{S}_r} \right) \right) \xrightarrow{d} ((\mathcal{M}, d), (\mathcal{M}, d)) \quad (5.6)$$

in distribution for the Gromov-Hausdorff topology.

Proof. For any compact metric space $X = (X, d)$ and any subspace $Y = (Y, d|_{Y \times Y})$ we have $d_{\text{GH}}(X, Y) \leq \frac{1}{2} \sup_{x \in X} d(x, Y)$. By Proposition 1.12, $\sup_{v \in v(\mathbf{R}_r)} r^{-1/4} d_{\mathbf{R}_r}(v, \mathbf{S}_r) \xrightarrow{P} 0$, and the result follows. \square

5.2. Analogous results for the largest 2-connected block. By analogy to Propositions 1.10 and 1.12, we have the following bounds for the submaps pendant to the largest 2-connected block in a uniform quadrangulation.

Given $\mathbf{Q}_q = (Q_q, e_q) \in \mathcal{Q}_q$, write $\mathbf{R}_q = R(Q_q)$, let e' be the $<_{Q_q}$ -minimal oriented edge of \mathbf{R}_q , and write $\mathbf{Q}'_q = (Q_q, e')$. By Proposition 1.7, \mathbf{Q}'_q uniquely decomposes as

$$(\mathbf{R}_q, ((L_i, b_i) : 0 \leq i \leq 2|e(\mathbf{R}_q)| - 4)) ,$$

where $L_i = (M_{i,j} : 1 \leq j \leq l_i) \in \mathcal{Q}^{l_i}$ and $b_i = (b_{i,j} : 1 \leq j \leq l_i) \in \{0, 1\}^{l_i}$, and $(l_i : 0 \leq i \leq 2|e(\mathbf{R}_q)| - 4)$ are suitable non-negative integers. Write

$$L'(Q_q) = \max(|v(M_{i,j})| : 0 \leq i \leq 2|e(\mathbf{R}_q)| - 4, 1 \leq j \leq l_i) , \quad (5.7)$$

$$D'(Q_q) = \max(\text{diam}(M_{i,j}) : 0 \leq i \leq 2|e(\mathbf{R}_q)| - 4, 1 \leq j \leq l_i) . \quad (5.8)$$

Proposition 5.3. *For all $\varepsilon \in (0, 1/3)$, there exist positive constants c_1, c_2 , and $c_3 = c_3(\varepsilon)$ such that, if $\mathbf{Q}_q \in_u \mathcal{Q}_{q,r(q)}$,*

$$\mathbb{P} \left(L'(\mathbf{Q}_q) \geq q^{2/3+\varepsilon} \right) \leq c_1 \exp(-c_2 q^{c_3}) .$$

Proposition 5.4. *Let $\mathbf{Q}_q \in_u \mathcal{Q}_{q,r(q)}$, then there exist positive constants C_1, C_2 , and C_3 such that*

$$\mathbb{P} \left(D'(\mathbf{Q}_q) \geq q^{5/24} \right) \leq C_1 \exp(-C_2 q^{C_3}) .$$

We omit proofs for the above propositions since the arguments are very similar as those for Propositions 1.10 and 1.12.

6. EXCHANGEABLE DECORATIONS

This section provides bounds on the Prokhorov distance between three sorts of measures on the vertices of a graph: the uniform measure, the degree-biased measure, and measures obtained by assigning vertices exchangeable random masses. In subsequent sections, these bounds help control the GHP distance between a map and its largest block.

Recall from the introduction that for a map G , and $c > 0$, cG denotes the measured metric space $(v(G), c \cdot d_G, \mu_G)$. Given a map G , write $\deg_G(v)$ for the degree of $v \in v(G)$ in G ; the *degree-biased measure* on G is the measure μ_G^B on $v(G)$ satisfying $\mu_G^B(S) = \sum_{v \in S} \deg_G(v) / (2|e(G)|)$.

Lemma 6.1. *For any quadrangulation Q and any $\varepsilon > 0$, with μ_Q and μ_Q^B viewed as measures on εQ , we have*

$$d_P(\mu_Q, \mu_Q^B) \leq \max(\varepsilon, 1/|v(Q)|).$$

Proof. Let n be the number of faces of Q , so that $|v(Q)| = n + 2$ and $|e(Q)| = 2n$. Fix $V \subsetneq v(Q)$ (the remaining case is trivial). Write $N(V) = \{v \in v(Q) : d_Q(v, V) \leq 1\}$. Let $V^+ = V \cup N(V)$; we claim that

$$\mu_Q^B(V^+) \geq \mu_Q(V) - \frac{1}{|v(Q)|}. \quad (6.1)$$

If this is so then in εQ we obtain $\mu_Q(V) \leq \mu_Q^B(V^\varepsilon) + 1/|v(Q)|$; since V was arbitrary, the lemma then follows easily. We now prove (6.1).

First suppose $Q[V]$ is connected, and write $p = |V|$. If $p = 1$ then the inequality is easily checked. If $p \geq 2$ then view $Q[V]$ as a quadrangulation with boundaries; let the boundaries have lengths ℓ_1, \dots, ℓ_k and write $\sum_{i=1}^k \ell_i = \ell$. We have $k \geq 1$ since $V \neq v(Q)$. A face f of Q is an *internal* face of $Q[V]$ if all vertices of f lie in V ; it is a *boundary face* of $Q[V]$ if some edge of $Q[V]$ is incident to f , but not all edges incident to f belong to $Q[V]$.

Writing i for the number of internal faces of $Q[V]$, Euler's formula straightforwardly yields $p = i + 2 + \ell/2 - k$. Furthermore, if f is a boundary face of $Q[V]$ then all edges of f lie within $Q[V^+]$. Now, a boundary face can be incident to at most two edges of $Q[V]$, so $Q[V]$ must have at least $\ell/2$ boundary faces. It follows that

$$\sum_{v \in V^+} \deg_Q(v) \geq \sum_{f \text{ internal to } Q[V^+]} 4 \geq 4(i + \ell/2) = 4(p + k - 2) \geq 4(p - 1).$$

Since $\sum_{v \in v(Q)} \deg_Q(v) = 2|e(Q)| = 4n$, it follows that

$$\mu_Q^B(V) \geq \frac{p-1}{n} = \mu_Q(V) - \frac{n+2-2p}{n(n+2)} \geq \mu_Q(V) - \frac{1}{|v(Q)|}.$$

Finally, if $Q[V]$ is not connected, the same argument applied component-wise yields the same bound. \square

Since both $1/s \rightarrow 0$ and $(\frac{3}{8s})^{1/4} \rightarrow 0$ as $s \rightarrow \infty$, the following is immediate.

Corollary 6.2. *For $S_s \in \mathcal{S}_s$, with μ_{S_s} and $\mu_{S_s}^B$ viewed as measures on $(\frac{3}{8s})^{1/4} S_s$, we have $d_P(\mu_{S_s}, \mu_{S_s}^B) \rightarrow 0$ as $s \rightarrow \infty$.*

In what follows, for a vector $\mathbf{x} = (x_1, \dots, x_k) \in \mathbb{R}^k$ write $|\mathbf{x}|_p = (\sum_{i=1}^k x_i^p)^{1/p}$. Suppose that $G = (G, e)$ is a rooted map. Enumerate the edges of G as e_1, \dots, e_m , where $m = |e(G)|$, and let e_0 be a second copy of the root edge e . (This makes sense even if G is random, as long as it is possible to specify a canonical way to order the edges of G ; for example, we may use the order \prec_G described in the introduction.)

For each $0 \leq i \leq m$, let w_i be a uniformly random endpoint of e_i . Let $\mathbf{n} = (n_0, \dots, n_m)$ be a vector of non-negative real numbers with $|\mathbf{n}|_1 > 0$. Define a (random) probability measure $\nu_G^{\mathbf{n}}$ on $v(G)$ as follows: for $V \subset v(G)$, let

$$\nu_G^{\mathbf{n}}(V) = \frac{1}{|\mathbf{n}|_1} \cdot \sum_{\{i: w_i \in V\}} n_i. \quad (6.2)$$

If one views $(w_i : 0 \leq i \leq 2s - 4)$ as attachment locations for pendant submaps, and \mathbf{n} as listing the masses of these submaps, then $\nu_G^{\mathbf{n}}$ is the probability measure assigning each vertex v a mass proportional to the total mass of submaps pendant to v .

Lemma 6.3. *Let $G = (G, e)$ have $|e(G)| = m$ and let $\mathbf{n} = (n_0, \dots, n_m)$ be an exchangeable random vector of non-negative real numbers with $|\mathbf{n}|_2$ deterministic and strictly positive. Then for any $V \subset v(G)$,*

$$\mathbb{P} \left(\left| \nu_G^{\mathbf{n}}(V) - \mu_G^B(V) \right| > \frac{2t}{|\mathbf{n}|_1} + \frac{1}{m+1} \right) \leq 4 \exp \left(-\frac{2t^2}{|\mathbf{n}|_2^2} \right).$$

In the proof, we will use the following result of Aldous [2, Theorem 20.6], which informally says that partial sums constructed by sampling without replacement may be obtained by first sampling with replacement and then taking a suitable projection.

Proposition 6.4 ([2], Proposition 20.6). *Fix $x_1, \dots, x_m \in \mathbb{R}$ and $k \in \{1, \dots, m\}$, let σ be a uniformly random permutation of $\{1, \dots, m\}$, and let I_1, \dots, I_k be independent and uniform on $\{1, \dots, m\}$. Then there exists a pair of random variables (X, Y) such that $\mathbb{E}[Y|X] = X$ and*

$$X \stackrel{d}{=} \sum_{j=1}^k x_{\sigma(j)}, \quad Y \stackrel{d}{=} \sum_{j=1}^k x_{I_j}.$$

Aldous [2] notes the following consequence of the preceding proposition, which is what we will in fact use.

Corollary 6.5 ([15], Theorem 2). *Under the conditions of Proposition 6.4, for all continuous convex functions $\phi : \mathbb{R} \rightarrow \mathbb{R}$,*

$$\mathbb{E}[\phi(X)] \leq \mathbb{E}[\phi(Y)].$$

Proof of Lemma 6.3. Given $V \subset v(G)$, write $\partial_e V = \{\{u, v\} \in e(G) : u \notin V, v \in V\}$ for the edge boundary of S . By definition, for $0 \leq j \leq m$, the vertex w_j is a uniformly random endpoint of e_j . We have

$$\nu_G^{\mathbf{n}}(V) = \frac{\sum_{\{j:e_j \in G[V]\}} n_j + \sum_{\{j:e_j \in \partial_e V\}} \mathbf{1}_{[w_j \in V]} n_j}{|\mathbf{n}|_1}. \quad (6.3)$$

We now show that $\nu_G^{\mathbf{n}}(V)$ is concentrated using Proposition 6.4. Independently for each $j \geq 1$ let $I_j \in_u \{0, \dots, m\}$. By the exchangeability of \mathbf{n} , it follows that for any continuous convex $\phi : \mathbb{R} \rightarrow \mathbb{R}$,

$$\mathbb{E} \left[\phi \left(\sum_{\{j:e_j \in G[V]\}} n_j \right) \right] \leq \mathbb{E} \left[\phi \left(\sum_{\{j:e_j \in G[V]\}} n_{I_j} \right) \right].$$

Also by exchangeability,

$$\mathbb{E} \left[\sum_{\{j:e_j \in G[V]\}} n_j \right] = |\mathbf{n}|_1 \cdot \frac{|e(G[V])|}{m+1}.$$

Taking $\phi(x) = e^{cx}$ for suitably chosen c and applying Markov's inequality as in [12, Theorem 2.5] yields Hoeffding's inequality-type bounds for $\sum_{\{j:e_j \in G[V]\}} n_j$:

$$\begin{aligned} & \mathbb{P} \left(\left| \sum_{\{j:e_j \in G[V]\}} \frac{n_j}{|\mathbf{n}|_1} - \frac{|e(G[V])|}{m+1} \right| > \frac{t}{|\mathbf{n}|_1} \right) \\ &= \mathbb{E} \left[\mathbb{P} \left(\left| \sum_{\{j:e_j \in G[V]\}} \frac{n_j}{|\mathbf{n}|_1} - \frac{|e(G[V])|}{m+1} \right| > \frac{t}{|\mathbf{n}|_1} \mid \mathbf{n} \right) \right] \\ &\leq \mathbb{E} \left[2 \exp \left(-\frac{2t^2}{|\mathbf{n}|_2^2} \right) \right] \\ &= 2 \exp \left(-\frac{2t^2}{|\mathbf{n}|_2^2} \right), \end{aligned}$$

recalling that $|\mathbf{n}|_2^2$ is deterministic. The random variables $\mathbf{1}_{[w_j \in V]}$ are iid Bernoulli(1/2), so a reprise of the argument yields

$$\mathbb{P} \left(\left| \sum_{\{j: e_j \in \partial_e V\}} \frac{\mathbf{1}_{[w_j \in V]} n_j}{|\mathbf{n}|_1} - \frac{|\partial_e V|}{2(m+1)} \right| > \frac{t}{|\mathbf{n}|_1} \right) \leq 2 \exp \left(-\frac{2t^2}{|\mathbf{n}|_2^2} \right).$$

We have

$$\mu_G^B(V) = \frac{1}{2|e(G)|} \sum_{v \in V} \deg(v) = \frac{2|e(G[V])| + |\partial_e V|}{2m},$$

so

$$\left| \mu_G^B(V) - \frac{|e(G[V])|}{m+1} - \frac{|\partial_e V|}{2(m+1)} \right| \leq \frac{1}{m+1}.$$

Considering (6.3), we then have

$$\begin{aligned} \mathbb{P} \left(|\nu_G^{\mathbf{n}}(V) - \mu_G^B(V)| > \frac{2t}{|\mathbf{n}|_1} + \frac{1}{m+1} \right) &\leq \mathbb{P} \left(\left| \sum_{\{j: e_j \in \partial_e V\}} \frac{\mathbf{1}_{[w_j \in V]} n_j}{|\mathbf{n}|_1} - \frac{|\partial_e V|}{2(m+1)} \right| > \frac{t}{|\mathbf{n}|_1} \right) \\ &+ \mathbb{P} \left(\left| \sum_{\{j: e_j \in G[V]\}} \frac{n_j}{|\mathbf{n}|_1} - \frac{|e(G[V])|}{m+1} \right| > \frac{t}{|\mathbf{n}|_1} \right). \end{aligned}$$

Now combine the three probability inequalities. \square

It is easily seen that the above lemma applies even for random sets V , so long as V is independent of the randomness used to select the endpoints w_i of the edges; we will use this in what follows.

7. PROJECTION OF MASSES IN RANDOM QUADRANGULATIONS

In this section we apply Lemma 6.3 to study projection of masses in large random quadrangulations, and in particular to prove Proposition 1.5. We begin by stating a straightforward corollary of Lemma 6.3. For a metric space $X = (X, d)$ and $x \in X$ write $B(x, r; X) = \{y : d(x, y) < r\}$.

Corollary 7.1. *For $s \in \mathbb{N}$ let $\mathbf{n}_s = (n_{s,0}, \dots, n_{s,2s-4})$ be an exchangeable random vector of non-negative real numbers. Let $\mathbf{S}_s \in_u \mathcal{S}_s$, and for $v \in v(\mathbf{S}_s)$ write $B(v, r) = B(v, r \cdot s^{1/4}; \mathbf{S}_s)$. Conditional on \mathbf{S}_s , let U and U' be independent, uniformly random elements of $v(\mathbf{S}_s)$. If $|\mathbf{n}_s|_1 \rightarrow \infty$ and $|\mathbf{n}_s|_2/|\mathbf{n}_s|_1 \rightarrow 0$ then for all $x \geq 0$,*

$$|\nu_{\mathbf{S}_s}^{\mathbf{n}_s}(B(U, x)) - \mu_{\mathbf{S}_s}^B(B(U, x))| \rightarrow 0, \quad (7.1)$$

$$|\nu_{\mathbf{S}_s}^{\mathbf{n}_s}(B(U, x) \cap B(U', x)) - \mu_{\mathbf{S}_s}^B(B(U, x) \cap B(U', x))| \rightarrow 0 \quad (7.2)$$

in probability as $s \rightarrow \infty$.

Proof. Fix $x \geq 0$. We assumed that $|\mathbf{n}_s|_1 \rightarrow \infty$ and $|\mathbf{n}_s|_2/|\mathbf{n}_s|_1 \rightarrow 0$; we may therefore choose a sequence $t(s)$ such that $t(s)/|\mathbf{n}_s|_1 \rightarrow 0$ and $t(s)/|\mathbf{n}_s|_2 \rightarrow \infty$. Now take $V = B(U, x)$. Recalling that $|e(\mathbf{S}_s)| + 1 = 2s - 3$, for any $\varepsilon > 0$, Lemma 6.3 gives

$$\begin{aligned} &\limsup_{s \rightarrow \infty} \mathbb{P} (|\nu_{\mathbf{S}_s}^{\mathbf{n}_s}(B(U, x)) - \mu_{\mathbf{S}_s}^B(B(U, x))| \geq \varepsilon) \\ &\leq \limsup_{s \rightarrow \infty} \mathbb{P} \left(|\nu_{\mathbf{S}_s}^{\mathbf{n}_s}(V) - \mu_{\mathbf{S}_s}^B(V)| \geq \frac{2t(s)}{|\mathbf{n}_s|_1} + \frac{1}{2s-3} \right) \\ &\leq \limsup_{s \rightarrow \infty} 4 \exp \left(-\frac{2t(s)^2}{|\mathbf{n}_s|_2^2} \right) \\ &= 0, \end{aligned}$$

which is (7.1). To prove (7.2) take $V = B(U, x) \cap B(U', x)$ and argue similarly. \square

Corollary 7.2. *Under the assumptions of Corollary 7.1, with $\nu_{\mathbf{S}_s}^{\mathbf{n}_s}$ and $\mu_{\mathbf{S}_s}^B$ viewed as measures on $(\frac{3}{8s})^{1/4} \mathbf{S}_s$, we have $d_P(\mu_{\mathbf{S}_s}, \nu_{\mathbf{S}_s}^{\mathbf{n}_s}) \rightarrow 0$ in probability as $s \rightarrow \infty$.*

Proof. By Corollary 6.2, it suffices to show that $d_P(\mu_{\mathbf{S}_s}^B, \nu_{\mathbf{S}_s}^{\mathbf{n}_s}) \xrightarrow{P} 0$. To achieve this, we use Corollary 7.1 and the compactness of the Brownian map $\mathbf{M} = (\mathcal{M}, d, \mu)$. For the remainder of the proof we abuse notation by writing $\mu_s = \mu_{\mathbf{S}_s}^B$ and $\nu_s = \nu_{\mathbf{S}_s}^{\mathbf{n}_s}$, for readability.

Fix $\varepsilon > 0$. By [1, Theorem 1], the triple $(v(\mathbf{S}_s), (\frac{3}{8s})^{1/4} d_{\mathbf{S}_s}, \mu_{\mathbf{S}_s})$ converges in distribution to \mathbf{M} as $r \rightarrow \infty$. Since \mathbf{M} is almost surely compact and μ a.s. has support \mathcal{M} , if $(U_i : i \in \mathbb{N})$ are independent with law μ then we almost surely have

$$K_\infty := \inf \left(k \in \mathbb{N} : \bigcup_{i=1}^k B(U_i, \varepsilon; \mathbf{M}) = \mathcal{M} \right) < \infty.$$

For $s \in \mathbb{N}$, let $(U_{s,i} : i \in \mathbb{N})$ be independent with law $\mu_{\mathbf{S}_s}$, and let

$$K_s = \inf \left(k \in \mathbb{N} : \bigcup_{i=1}^k B(U_{s,i}, \varepsilon; (3/8s)^{1/4} \mathbf{S}_s) = v(\mathbf{S}_s) \right).$$

The aforementioned distributional convergence and the a.s. finiteness of K_∞ together imply that there exists $K \in \mathbb{N}$ such that for all $s \in \mathbb{N}$, $\mathbb{P}(K_s > K) < \varepsilon$.

For $i \geq 1$ let

$$B_i = B \left(U_{s,i}, \varepsilon; (3/(8s))^{1/4} \mathbf{S}_s \right). \quad (7.3)$$

Let $A_1 = B_1$, and for $i > 1$ let $A_i = B_i \setminus \bigcup_{j=1}^{i-1} B_j$. Then A_1, \dots, A_{K_s} is a covering of $v(\mathbf{S}_s)$ by disjoint sets.

Suppose that $d_P(\mu_s, \nu_s) > \varepsilon$. Then there exists a set $S \subset v(\mathbf{S}_s)$ such that either $\mu_s(S^\varepsilon) < \nu_s(S) - \varepsilon$ or $\nu_s(S^\varepsilon) < \mu_s(S) - \varepsilon$. Since A_1, \dots, A_{K_s} partition $v(\mathbf{S}_s)$, there is j such that either

$$\begin{aligned} \mu_s(S^\varepsilon \cap A_j) &< \nu_s(S \cap A_j) - \varepsilon/K_s \leq \nu_s(A_j) - \varepsilon/K_s \text{ or} \\ \nu_s(S^\varepsilon \cap A_j) &< \mu_s(S \cap A_j) - \varepsilon/K_s \leq \mu_s(A_j) - \varepsilon/K_s. \end{aligned}$$

For one of these to occur we must have $S \cap A_j \neq \emptyset$. Since A_j has radius at most ε , it follows that $A_j \subset S^\varepsilon$. Thus, either

$$\mu_s(A_j) < \nu_s(A_j) - \varepsilon/K_s \quad \text{or} \quad \nu_s(A_j) < \mu_s(A_j) - \varepsilon/K_s.$$

This yields the bound

$$\begin{aligned} &\mathbb{P}(d_P(\mu_s, \nu_s) > \varepsilon) \\ &\leq \mathbb{P}(|\mu_s(A_j) - \nu_s(A_j)| > \varepsilon/K_s \text{ for some } 1 \leq j \leq K_s) \\ &\leq \sum_{j=1}^{K_s} \mathbb{P}(|\mu_s(A_j) - \nu_s(A_j)| > \varepsilon/K_s, K_s \leq K) + \mathbb{P}(K_s > K) \\ &\leq \sum_{j=1}^{K_s} \mathbb{P}(|\mu_s(A_j) - \nu_s(A_j)| > \varepsilon/K) + \varepsilon. \end{aligned} \quad (7.4)$$

By the triangle inequality, for all $1 \leq i \leq K$,

$$\begin{aligned} &|\mu_s(A_i) - \nu_s(A_i)| \\ &= \left| \mu_s \left(B_i \setminus \bigcup_{j=1}^{i-1} B_j \right) - \nu_s \left(B_i \setminus \bigcup_{j=1}^{i-1} B_j \right) \right| \\ &\leq |\mu_s(B_i) - \nu_s(B_i)| + \left| \mu_s \left(B_i \cap \bigcup_{j=1}^{i-1} B_j \right) - \nu_s \left(B_i \cap \bigcup_{j=1}^{i-1} B_j \right) \right| \end{aligned}$$

$$\leq |\mu_s(B_i) - \nu_s(B_i)| + \sum_{j=1}^{i-1} |\mu_s(B_i \cap B_j) - \nu_s(B_i \cap B_j)| ,$$

where the last sum equals 0 in the case $i = 1$. Recalling the definitions of the B_i from (7.3), the preceding inequality and Corollary 7.1 imply that for each fixed $i \geq 1$,

$$|\mu_s(A_i) - \nu_s(A_i)| \rightarrow 0$$

in probability as $s \rightarrow \infty$. Combining this with (7.4), we obtain

$$\limsup_{s \rightarrow \infty} \mathbb{P}(d_P(\mu_s, \nu_s) > \varepsilon) \leq \varepsilon .$$

Since $\varepsilon > 0$ was arbitrary, this completes the proof. \square

We are almost ready to prove Proposition 1.5; before doing so we state two easy facts, which each provide bounds on the GHP distance between a measured metric space and an induced (in some sense) subspace. The first fact is immediate from the definition of d_{GHP} .

Fact 7.3. *Fix a measured metric space $X = (X, d, \mu)$ and $Y \subset X$, and let μ_Y be a Borel measure on (Y, d_Y) , where $d_Y = d|_{Y \times Y}$ is the induced metric. Write $Y = (Y, d_Y, \mu_Y)$. Then $d_{\text{GHP}}(X, Y) \leq \max(d_{\text{H}}(X, Y), d_P(\mu, \mu_Y))$.*

The second fact informally says that in a compact measured metric space, projecting onto an ε -net does not change the space very much (in the GHP sense). The proof is left to the reader.

Fact 7.4. *Let $X = (X, d, \mu)$ be a compact measured metric space, and let $S \subset X$ be finite so that there exists $\varepsilon > 0$ with $X \subset S^\varepsilon$. Let $(X_s : s \in S)$ be such that $\bigcup_{s \in S} X_s = X$, that $\mu(X_s \cap X_{s'}) = 0$ for $s \neq s'$, and that $X_s \subset B(s, \varepsilon; X)$ for all $s \in S$. Define a measure ν on S by $\nu(s) = \mu(X_s)$ for any $s \in S$, and let $S = (S, d|_{S \times S}, \nu)$. Then*

$$d_{\text{GHP}}(X, S) \leq \varepsilon .$$

Proof of Proposition 1.5. For $r \in \mathbb{N}$, let $\mathbf{R}_r \in_u \mathcal{R}_{r, s(r)}$, and let $\mathbf{S}_r = \mathbf{S}(\mathbf{R}_r)$. Write $\mathbf{R}_r = (R_r, e_r)$, and let e' be the $\langle \mathbf{R}_r$ -minimal oriented edge of \mathbf{S}_r . Next, apply the bijection of Proposition 1.6 to the map $\mathbf{R}'_r = (R_r, e')$: this decomposes \mathbf{R}_r into \mathbf{S}_r together with a sequence $(\Theta_i : 0 \leq i \leq 2s(r) - 4)$ of submaps of \mathbf{R}_r . Let $n_{r,0} = |e(\Theta_0)|\mathbf{1}_{[|e(\Theta_i)| > 1]}$, and for $1 \leq i \leq 2s(r) - 4$ let $n_{r,i} = 1 + |e(\Theta_i)|\mathbf{1}_{[|e(\Theta_i)| > 1]}$. Then let $\mathbf{n}_r = (n_{r,0}, \dots, n_{r, 2s(r)-4})$ and construct the measure $\nu_{\mathbf{S}_r}^{\mathbf{n}_r}$ as in (6.2): for $0 \leq i \leq 2s(r) - 4$, w_i is a random endpoint of e_i , and

$$\nu_{\mathbf{S}_r}^{\mathbf{n}_r}(v) = \frac{1}{2r-4} \sum_{\{i: w_i=v\}} n_{r,i} .$$

(The difference of 1 in the definition of $n_{r,0}$ accounts for the fact that when reconstructing \mathbf{R}_r from \mathbf{S}_r and the Θ_i , we identify two copies of the root edge; the fact that $2r - 4$ is the correct normalization follows from (1.1).)

We have $|\mathbf{n}_r|_1 = 2r - 4 \rightarrow \infty$ as $r \rightarrow \infty$. Furthermore, if $L(\mathbf{R}_r) \leq r^{3/4}$ then $n_{r,i} \leq 2r^{3/4} - 3$ for all i , so $|\mathbf{n}_r|_2 / |\mathbf{n}_r|_1 = O(r^{-1/8}) \rightarrow 0$. By Proposition 1.10, we have $\mathbb{P}(L(\mathbf{R}_r) \leq r^{3/4}) \rightarrow 1$, so $|\mathbf{n}_r|_2 / |\mathbf{n}_r|_1 \xrightarrow{P} 0$.

Corollary 7.2 now implies that $d_P(\mu_{\mathbf{S}_r}, \nu_{\mathbf{S}_r}^{\mathbf{n}_r}) \rightarrow 0$ as $r \rightarrow \infty$, with the measures viewed as living on $(\frac{21}{40r})^{1/4} \mathbf{S}_r$. For Borel measures μ, ν on a compact metric space (X, d) , we have $d_{\text{GHP}}((X, d, \mu), (X, d, \nu)) = d_P(\mu, \nu)$, so

$$d_{\text{GHP}}\left(\left(\frac{21}{40r}\right)^{1/4} \mathbf{S}_r, \left(v(\mathbf{S}_r), \left(\frac{21}{40r}\right)^{1/4} \cdot d_{\mathbf{S}_r}, \nu_{\mathbf{S}_r}^{\mathbf{n}_r}\right)\right) \xrightarrow{P} 0 . \quad (7.5)$$

We now bound the distance from $(\frac{21}{40r})^{1/4} \mathbf{R}_r$ to the latter space. It is convenient to work with a graph with edge lengths rather than a finite measured metric space. More precisely, view each edge e of \mathbf{R}_r as an isometric copy I_e of the unit interval $[0, 1]$, endowed with the

rescaled Lebesgue measure $(2r-4)^{-1} \cdot \text{Leb}_{I_e}$, and write $\mathbf{R}' = (R'_r, d'_r, \mu'_r)$ for the resulting measured metric space. We then have $\mu'_r(R'_r) = (2r-4)^{-1} \cdot \sum_{e \in e(R_r)} \text{Leb}_{I_e}(I_e) = 1$.

We may naturally identify $v(R_r)$ with the set of endpoints of edges in \mathbf{R}' , and this is an isometric embedding in that with this identification we have $d_{R_r} = d'_r|_{v(R_r)}$. Furthermore, the degree-biased measure $\mu_{R_r}^B$ may be obtained by projection onto $v(R_r)$: for $v \in v(R_r)$ we have $\mu_{R_r}^B(v) = \mu'_r(B(v, 1/2; \mathbf{R}')) = \deg_{R_r}(v)/(2(2s-4))$. By Fact 7.4, it follows that for any $\varepsilon > 0$,

$$d_{\text{GHP}}(\varepsilon \mathbf{R}', (v(R_r), \varepsilon d_{R_r}, \mu_{R_r}^B)) \leq \varepsilon. \quad (7.6)$$

The space $\mathbf{S}_r = (v(S_r), d_{S_r})$ is likewise isometrically embedded within \mathbf{R}' , and we may also obtain the measure $\nu_{\mathbf{S}_r}^{\mathbf{n}_r}$ by projection. To do so, let $E_i = e(\Theta_i)$ for $1 \leq i \leq 2s(r)-4$, let $E_0 = E(\Theta_0) \setminus \{e'\}$, and for $v \in v(S_r)$ let

$$X_v = \bigcup_{\{i:w_i=v\}} \bigcup_{e \in E_i} I_e.$$

Then $\nu_{\mathbf{S}_r}^{\mathbf{n}_r}(v) = \mu'_r(X_v)$. Furthermore, $(X_v : v \in v(S_r))$ covers R'_r and $\mu'_r(X_u \cap X_v) = 0$ for $u \neq v$ since edges only intersect at their endpoints. Recalling the definition of $D(\mathbf{R}_r)$ from Section 1.2, for any $v \in V$ we have $X_v \subset B(v, D(\mathbf{R}_r); \mathbf{R}')$. It follows from Fact 7.4 that for all $\varepsilon > 0$,

$$d_{\text{GHP}}(\varepsilon \mathbf{R}', (v(S_r), \varepsilon d_{S_r}, \nu_{\mathbf{S}_r}^{\mathbf{n}_r})) \leq \varepsilon \cdot D(\mathbf{R}_r). \quad (7.7)$$

We always have $D(\mathbf{R}_r) \geq 1$, so combining (7.6), (7.7) gives

$$d_{\text{GHP}}((v(S_r), \varepsilon d_{S_r}, \nu_{\mathbf{S}_r}^{\mathbf{n}_r}), (v(R_r), \varepsilon d_{R_r}, \mu_{R_r}^B)) \leq 2\varepsilon \cdot D(\mathbf{R}_r)$$

Using Lemma 6.1 to bound $d_{\text{GHP}}((v(R_r), \varepsilon d_{R_r}, \mu_{R_r}^B), \varepsilon \mathbf{R}_r)$, the triangle inequality then gives

$$d_{\text{GHP}}((v(S_r), \varepsilon d_{S_r}, \nu_{\mathbf{S}_r}^{\mathbf{n}_r}), \varepsilon \mathbf{R}_r) \leq 2\varepsilon \cdot D(\mathbf{R}_r) + \max(\varepsilon, 1/r). \quad (7.8)$$

By Proposition 1.12, $r^{-1/4}D(\mathbf{R}_r) \xrightarrow{P} 0$, and (7.5) then implies that

$$d_{\text{GHP}}\left(\left(\frac{21}{40r}\right)^{1/4} \mathbf{S}_r, \left(\frac{21}{40r}\right)^{1/4} \mathbf{R}_r\right) \xrightarrow{P} 0.$$

Since $\frac{3}{8s(r)} \cdot \mathbf{S}_r \xrightarrow{d} \mathbf{M}$ as $r \rightarrow \infty$, and $\frac{3}{8s(r)} = (1 + o(1))\frac{21}{40r}$, the result follows. \square

8. THE REMAINING PROOFS

Recall that \mathbb{K} is the set of measured isometry classes of compact metric spaces, and that GHP convergence refers to convergence in the the Polish space $(\mathbb{K}, d_{\text{GHP}})$.

Proof of Theorem 1.1. Let $g : \mathbb{K} \rightarrow \mathbb{R}$ be a bounded continuous function, and write $\|g\| = \sup |g| < \infty$. Recall that $\mathbf{R}_r \in_u \mathcal{R}_r$, and let $\mathbf{M}_r = \left(\frac{21}{40r}\right)^{1/4} \mathbf{R}_r$. We show that

$$\mathbb{E}[g(\mathbf{M}_r)] \rightarrow \mathbb{E}[g(\mathbf{M})] \quad (8.1)$$

as $r \rightarrow \infty$; the result then follows by the Portmanteau theorem.

The proof of (8.1) is simply summarized: average over the size of $\text{sb}(\mathbf{R}_r)$. The details are also fairly straightforward. Fix $\varepsilon \in (0, 1/2)$ with $\varepsilon < 1/\|g\|$, let \mathcal{A} be the Airy density and let β given by Proposition 1.8. Then fix $C_\varepsilon > 0$ large enough that $\int_{-C_\varepsilon}^{C_\varepsilon} \beta \mathcal{A}(\beta x) dx > 1 - \varepsilon$. Recall from the introduction $s(r)$ satisfies $|s(r) - 5r/7| \leq Cr^{2/3}$ for large r . The constant C was fixed but arbitrary, so we may assume that $C > C_\varepsilon$.

Next, for $r, s \in \mathbb{N}$ with $s \leq r$, let $\mathbf{R}_{r,s} \in_u \mathcal{R}_{r,s}$ and write $\mathbf{M}_{r,s} = \left(\frac{21}{40r}\right)^{1/4} \mathbf{R}_{r,s}$. We claim that

$$\sup_{\{s \in \mathbb{N} : |s - 5r/7| \leq C_\varepsilon r^{2/3}\}} |g(\mathbf{M}_{r,s}) - g(\mathbf{M})| \rightarrow 0 \quad (8.2)$$

as $r \rightarrow \infty$. Indeed: otherwise we may find a sequence $(\hat{s}(r), r \geq 1)$ such that $|\hat{s}(r) - 5r/7| \leq C_\varepsilon r^{2/3} < Cr^{2/3}$ with $\limsup_{r \rightarrow \infty} |g(\mathbf{M}_{r, \hat{s}(r)}) - g(\mathbf{M})| \neq 0$. By the Portmanteau

theorem, this implies that $\mathbf{M}_{r,s}$ does not converge in distribution to \mathbf{M} , contradicting Proposition 1.5. This establishes (8.2).

Now for each $r \in \mathbb{N}$, let

$$E_r = \left\{ |\text{sb}(\mathbf{R}_r) - 5r/7| \leq C_\varepsilon r^{2/3} \right\} .$$

Recalling the definition of $\delta_s(\cdot)$ from Theorem 1.4, it follows from Proposition 1.8 and a Riemann approximation that for large enough r ,

$$\begin{aligned} \mathbb{P}(E_r) &= (1 + o(1)) \sum_{\{s \in \mathbb{N}: |s - 5r/7| \leq C_\varepsilon r^{2/3}\}} \frac{\beta \mathcal{A}(\beta \delta_s(q))}{r^{2/3}} \\ &= (1 + o(1)) \int_{-C_\varepsilon}^{C_\varepsilon} \beta \mathcal{A}(\beta s) ds \\ &> 1 - 2\varepsilon. \end{aligned}$$

Then for large enough r ,

$$\left| \mathbb{E}[g(\mathbf{M}_r)] - \mathbb{E}[g(\mathbf{M}_r \mathbf{1}_{[E_r]})] \right| \leq \mathbb{P}(E_r^c) \|g\| < 2\varepsilon \|g\|. \quad (8.3)$$

We now show that $\left| \mathbb{E}[g(\mathbf{M}_r \mathbf{1}_{[E_r]})] - \mathbb{E}[g(\mathbf{M})] \right|$ is also small. The conditional law of \mathbf{R}_r given that $\text{sb}(\mathbf{R}_r) = s$ is identical to that of $\mathbf{R}_{r,s}$, so

$$\mathbb{E}[g(\mathbf{M}_r \mathbf{1}_{[E_r]})] = \sum_{\{s \in \mathbb{N}: |s - 5r/7| \leq C_\varepsilon r^{2/3}\}} \mathbb{P}(\text{sb}(\mathbf{R}_r) = s) \mathbb{E}[g(\mathbf{M}_{r,s})] .$$

By the triangle inequality, we therefore have

$$\begin{aligned} &\left| \mathbb{E}[g(\mathbf{M}_r \mathbf{1}_{[E_r]})] - \mathbb{E}[g(\mathbf{M})] \right| \\ &\leq \sum_{\{s \in \mathbb{N}: |s - 5r/7| \leq C_\varepsilon r^{2/3}\}} \mathbb{P}(\text{sb}(\mathbf{R}_r) = s) \cdot \left| \mathbb{E}[g(\mathbf{M}_{r,s})] - \mathbb{E}[g(\mathbf{M})] \right| \\ &\leq \sup_{\{s \in \mathbb{N}: |s - 5r/7| \leq C_\varepsilon r^{2/3}\}} |g(\mathbf{M}_{r,s}) - g(\mathbf{M})|. \end{aligned} \quad (8.4)$$

This tends to 0 by (8.2), which with (8.3) gives $\limsup_{r \rightarrow \infty} \left| \mathbb{E}[g(\mathbf{M}_r)] - \mathbb{E}[g(\mathbf{M})] \right| \leq 2\varepsilon \|g\|$. Since $\varepsilon > 0$ was arbitrary, this establishes (8.1) and completes the proof. \square

In the remaining proofs, we use the following simple fact. Recall the definition of L' from (5.7).

Fact 8.1. *Let $\mathbf{Q}_q \in \mathcal{Q}_q$, write $\mathbf{R}_q = \mathbf{R}(\mathbf{Q}_q)$ and $\mathbf{S}_q = \mathbf{S}(\mathbf{Q}_q)$. Note that if $\mathbf{S}_q \neq \mathbf{S}(\mathbf{R}_q)$, then $\text{sb}(\mathbf{Q}_q) \geq \text{sb}(\mathbf{R}_q)$, and it follows that \mathbf{Q}_q contains at least two 2-connected blocks of size at least $\text{sb}(\mathbf{R}_q)$, implying that $L'(\mathbf{Q}_q) \geq \text{sb}(\mathbf{R}_q)$.*

Proof of Theorem 1.4. Fix $q \in \mathbb{N}$ and write $\mathbf{R}_q = \mathbf{R}(\mathbf{Q}_q)$. Let $\mathbf{R} \in_u \mathcal{R}_{r(q)}$. Given that $\mathbf{b}(\mathbf{Q}_q) = r(q)$, $\mathbf{R}_q \stackrel{d}{=} \mathbf{R}$, so

$$\begin{aligned} &\mathbb{P}(\mathbf{b}(\mathbf{Q}_q) = r(q), \text{sb}(\mathbf{R}_q) = s(r(q))) \\ &= \mathbb{P}(\text{sb}(\mathbf{R}_q) = s(r(q)) \mid \mathbf{b}(\mathbf{Q}_q) = r(q)) \cdot \mathbb{P}(\mathbf{b}(\mathbf{Q}_q) = r(q)) \\ &= \mathbb{P}(\text{sb}(\mathbf{R}) = s(r(q))) \cdot \mathbb{P}(\mathbf{b}(\mathbf{Q}_q) = r(q)) . \end{aligned}$$

Writing $\beta = \frac{5^{2/3} \cdot 15}{28}$ and $\beta' = \frac{3^{2/3} \cdot 45}{14^{1/3} \cdot 8}$, by Propositions 4.3 and 1.8, we thus have

$$\mathbb{P}(\mathbf{b}(\mathbf{Q}_q) = r(q), \text{sb}(\mathbf{R}_q) = s(r(q))) = \frac{\beta \mathcal{A}(\beta \delta_r(q))}{q^{2/3}} \frac{\beta' \mathcal{A}(\beta' \delta_s(q))}{r(q)^{2/3}} (1 + o(1)) . \quad (8.5)$$

Next,

$$\left| \mathbb{P}(\mathbf{b}(\mathbf{Q}_q) = r(q), \text{sb}(\mathbf{R}_q) = s(r(q))) - \mathbb{P}(\mathbf{b}(\mathbf{Q}_q) = r(q), \text{sb}(\mathbf{Q}_q) = s(r(q))) \right|$$

$$\begin{aligned} &\leq \mathbb{P}(\mathbf{b}(\mathbf{Q}_q) = r(q), \text{sb}(\mathbf{Q}_q) \neq s(r(q)), \text{sb}(\mathbf{R}_q) = s(r(q))) \\ &\quad + \mathbb{P}(\mathbf{b}(\mathbf{Q}_q) = r(q), \text{sb}(\mathbf{Q}_q) = s(r(q)), \text{sb}(\mathbf{R}_q) \neq s(r(q))) , \end{aligned} \quad (8.6)$$

If $\{\mathbf{b}(\mathbf{Q}_q) = r(q), \text{sb}(\mathbf{Q}_q) \neq s(r(q)), \text{sb}(\mathbf{R}_q) = s(r(q))\}$ occurs then $L'(\mathbf{Q}_q) \geq s(r(q))$, as explained in Fact 8.1. Similarly, if $\{\mathbf{b}(\mathbf{Q}_q) = r(q), \text{sb}(\mathbf{Q}_q) = s(r(q)), \text{sb}(\mathbf{R}_q) \neq s(r(q))\}$ occurs then \mathbf{Q}_q must contain a simple block of size $s(r(q))$ that does not lie within \mathbf{R}_q ; since $\mathbf{b}(\mathbf{Q}_q) = r(q) > s(r(q))$, in this case we also obtain $L'(\mathbf{Q}_q) \geq s(r(q))$. It follows from Proposition 5.3 that there exist positive constants c_1, c_2, c_3 such that

$$\begin{aligned} &|\mathbb{P}(\mathbf{b}(\mathbf{Q}_q) = r(q), \text{sb}(\mathbf{R}_q) = s(r(q))) - \mathbb{P}(\mathbf{b}(\mathbf{Q}_q) = r(q), \text{sb}(\mathbf{Q}_q) = s(r(q)))| \\ &\leq 2\mathbb{P}(L'(\mathbf{Q}_q) \geq s(r(q))) \\ &\leq c_1 \exp(-c_2 q^{c_3}) \\ &= o(q^{-2}) , \end{aligned}$$

which combined with (8.5) proves the theorem. \square

For the proof of Theorem 1.3, we require a lemma bounding the maximum degree in a quadrangulation uniformly drawn from $\mathcal{Q}_{q,r(q),s(q)}$; the lemma follows easily from the fact that degrees in uniform quadrangulations have exponential tails.

Lemma 8.2. *Let $\mathbf{Q}_q \in_u \mathcal{Q}_{q,r(q),s(q)}$. Then for all q sufficiently large,*

$$\mathbb{P}\left(\max(\deg_{\mathbf{Q}_q}(w) : w \in v(\mathbf{Q}_q)) \geq (\ln q)^2\right) < q^{-10} .$$

Proof. By [4, Theorem 2.1 (a)] (and Tutte's angular bijection between maps and quadrangulations), for all $\varepsilon > 0$ there exists $B > 0$ such that for all $q \geq 3$, if $\mathbf{Q} \in_u \mathcal{Q}_q$ and $u \in_u v(\mathbf{Q})$ then

$$\mathbb{P}(\deg_{\mathbf{Q}}(v) > d) < B \left(\frac{1}{2} + \varepsilon\right)^d . \quad (8.7)$$

Given that $\mathbf{b}(\mathbf{Q}) = r(q)$ and $\text{sb}(\mathbf{R}(\mathbf{Q})) = s(q)$, the conditional law of \mathbf{Q} is uniform on $\mathcal{Q}_{q,r(q),s(q)}$; so

$$\begin{aligned} &\mathbb{P}\left(\max(\deg_{\mathbf{Q}_q}(w) : w \in v(\mathbf{Q}_q)) > d\right) \\ &= \mathbb{P}\left(\max(\deg_{\mathbf{Q}}(w) : w \in v(\mathbf{Q})) > d \mid \mathbf{b}(\mathbf{Q}) = r(q), \text{sb}(\mathbf{R}(\mathbf{Q})) = s(q)\right) \\ &\leq q \cdot \mathbb{P}(\deg_{\mathbf{Q}}(u) > d \mid \mathbf{b}(\mathbf{Q}) = r(q), \text{sb}(\mathbf{R}(\mathbf{Q})) = s(q)) \\ &\leq q \cdot \frac{\mathbb{P}(\deg_{\mathbf{Q}}(u) > d)}{\mathbb{P}(\mathbf{b}(\mathbf{Q}) = r(q), \text{sb}(\mathbf{R}(\mathbf{Q})) = s(q))} \\ &= O(q^{7/3})\mathbb{P}(\deg_{\mathbf{Q}}(u) > d) , \end{aligned}$$

the final inequality by Theorem 1.4 and the definition of $r(q)$ and $s(q)$. Taking $d = \ln^2 q$ and $\varepsilon < 1/2$, the result then follows from (8.7). \square

Proof of Theorem 1.3. Recall that $\mathbf{Q}_q \in_u \mathcal{Q}_{q,r(q),s(q)}$, and $\mathbf{R}_q = R(\mathbf{Q}_q)$ and $\mathbf{S}_q = S(\mathbf{Q}_q)$. Let $\mathbf{Q} \in_u \mathcal{Q}_q$. Given that $\mathbf{b}(\mathbf{Q}) = r(q)$ and $\text{sb}(\mathbf{Q}) = s(q)$, we have $\mathbf{Q}_q \stackrel{d}{=} \mathbf{Q}$. By Fact 8.1, we then have

$$\begin{aligned} \mathbb{P}(\mathbf{S}_q \neq S(\mathbf{R}_q)) &= \mathbb{P}(S(\mathbf{Q}) \neq S(\mathbf{R}(\mathbf{Q})) \mid \mathbf{b}(\mathbf{Q}) = r(q), \text{sb}(\mathbf{Q}) = s(q)) \\ &\leq \mathbb{P}(L'(\mathbf{Q}) \geq s(q) \mid \mathbf{b}(\mathbf{Q}) = r(q), \text{sb}(\mathbf{Q}) = s(q)) \\ &\leq \frac{\mathbb{P}(L'(\mathbf{Q}) \geq s(q) \mid \mathbf{b}(\mathbf{Q}) = r(q))}{\mathbb{P}(\mathbf{b}(\mathbf{Q}) = r(q), \text{sb}(\mathbf{Q}) = s(q))} . \end{aligned}$$

Combined with Theorem 1.4, this gives

$$\mathbb{P}(\mathbf{S}_q \neq \mathbf{S}(\mathbf{R}_q)) = O\left(q^{4/3}\right) \cdot \mathbb{P}(L'(\mathbf{Q}) \geq s(q) | \mathbf{b}(\mathbf{Q}) = r(q)) .$$

Since $s(q) = q/3 + O(q^{2/3})$, by Proposition 5.3 there exist $c_2, c_3 > 0$ such that

$$\mathbb{P}(L'(\mathbf{Q}) \geq s(q) | \mathbf{b}(\mathbf{Q}) = r(q)) = O(\exp(-c_2 q^{c_3})) .$$

Hence,

$$\mathbb{P}(\mathbf{S}_q \neq \mathbf{S}(\mathbf{R}_q)) = O\left(q^{4/3} \cdot \exp(-c_2 q^{c_3})\right) . \quad (8.8)$$

Now let $\mathbf{R} \in_u \mathcal{R}_{r(q), s(q)}$. Given that $\mathbf{S}_q = \mathbf{S}(\mathbf{R}_q)$, we have $\mathbf{R}_q \in_u \mathcal{R}_{r(q), s(q)}$, so (8.8) implies easily that for any bounded continuous function $g : \mathbb{K}^2 \rightarrow \mathbb{R}$

$$\left| \mathbb{E} \left[g \left(\left(\frac{21}{40r(q)} \right)^{1/4} \mathbf{R}_q, \left(\frac{21}{40r(q)} \right)^{1/4} \mathbf{S}_q \right) \right] - \mathbb{E} \left[g \left(\left(\frac{21}{40r(q)} \right)^{1/4} \mathbf{R}, \left(\frac{21}{40r(q)} \right)^{1/4} \mathbf{S}(\mathbf{R}) \right) \right] \right| \rightarrow 0 ,$$

as $q \rightarrow \infty$. By Proposition 1.5 and the Portmanteau theorem, it follows that as $r(q) \rightarrow \infty$,

$$\left(\left(\frac{21}{40r(q)} \right)^{1/4} \mathbf{R}_q, \left(\frac{21}{40r(q)} \right)^{1/4} \mathbf{S}_q \right) \xrightarrow{d} (\mathbf{M}, \mathbf{M}) . \quad (8.9)$$

Moreover, by the definition of $r(q)$, there exist $C_1, C_2 > 0$ such that for all $q > 0$,

$$\frac{9}{8q + C_1 q^{2/3}} \leq \frac{21}{40r(q)} \leq \frac{9}{8q - C_2 q^{2/3}} .$$

From this and (8.9) we obtain

$$\left(\left(\frac{9}{8q} \right)^{1/4} \mathbf{R}_q, \left(\frac{9}{8q} \right)^{1/4} \mathbf{S}_q \right) \xrightarrow{d} (\mathbf{M}, \mathbf{M}) \quad (8.10)$$

as $q \rightarrow \infty$. To finish the proof, we show that also

$$\left(\left(\frac{9}{8q} \right)^{1/4} \mathbf{Q}_q, \left(\frac{9}{8q} \right)^{1/4} \mathbf{R}_q \right) \xrightarrow{d} (\mathbf{M}, \mathbf{M}) . \quad (8.11)$$

Joint convergence of the triple to the limit $(\mathbf{M}, \mathbf{M}, \mathbf{M})$ is immediate from (8.10) and (8.11), so it remains to prove (8.11). (Note that we may not simply invoke the result of Le Gall [11] and of Miermont [14] to conclude that the $\left(\frac{9}{8q}\right)^{1/4} \mathbf{Q}_q \xrightarrow{d} \mathbf{M}$ since \mathbf{Q}_q is not uniformly distributed over \mathcal{Q}_q , but over $\mathcal{Q}_{q, r(q), s(q)}$.) The argument is similar to that in Proposition 1.5, and we focus on explaining the points where it differs.

Let e' be the $\prec_{\mathbf{Q}_q}$ -minimal oriented edge of \mathbf{R}_q ; by definition, this is the root edge of \mathbf{R}_q . Write $\mathbf{Q}_q = (Q_q, e_q)$ and $\mathbf{R}_q = (R_q, e')$. Also, let $\mathbf{Q}'_q = (Q_q, e')$. The bijection ψ from Proposition 1.7 gives a decomposition of \mathbf{Q}'_q as

$$(R_q, ((L_i, b_i) : 0 \leq i \leq 2r(q) - 4)) ,$$

where the $L_i = (M_{i,j} : 1 \leq j \leq \ell_i) \in \mathcal{Q}^{\ell_i}$ satisfy (recalling (1.2))

$$|e(Q_q)| = |e(R_q)| + \sum_{i=0}^{|e(R_q)|} \sum_{j=1}^{\ell_i} (|e(M_{i,j})| + 1 + \mathbf{1}_{[|e(M_{i,j})|=1]}) , \quad (8.12)$$

and $b_i = (b_{i,j} : 1 \leq j \leq \ell_i) \in \{0, 1\}^{\ell_i}$.

List the elements of $e(R_q)$ as $(e_i : 1 \leq i \leq |e(R_q)|)$ according to the order $\prec_{\mathbf{R}_q}$; like in Section 1.1, we view e_i as oriented (we oriented so that the tail e_i^- precedes the head e_i^+ according to the breadth-first order described in the introduction, but this is unimportant; all that matters is to have a fixed rule for choosing the orientation). Also, let e_0 be a copy

of e' . Under the bijection ψ , for each $0 \leq i \leq |e(\mathbf{R}_q)|$ and $1 \leq j \leq \ell_i$, the value $b_{i,j}$ indicates the endpoint e_i at which $M_{i,j}$ is attached.

Recall that $\mu^B = \mu_{\mathbf{Q}_q}^B$ is the degree-biased measure on $v(\mathbf{Q}_q)$. We now compare μ^B with a random projection of μ^B onto \mathbf{R}_q . First define a vector \mathbf{n}_q as follows. Let $n_0 = 0$ if $\ell_0 = 0$ and otherwise let $n_0 = \sum_{j=1}^{\ell_0} (|e(M_{0,j})| + 1 + \mathbf{1}_{[|e(M_{0,j})| \neq 1]})$, and for $1 \leq i \leq 2r(q) - 4$ let

$$n_i = 1 + \sum_{j=1}^{\ell_i} (|e(M_{i,j})| + 1 + \mathbf{1}_{[|e(M_{i,j})| \neq 1]}) \quad (8.13)$$

Set $\mathbf{n}_q = (n_i : 0 \leq i \leq 2r(q) - 4)$; it is immediate from Proposition 1.7 that \mathbf{n}_q is exchangeable. Now define the measure $\nu^{\mathbf{n}_q} = \nu_{\mathbf{R}_q}^{\mathbf{n}_q}$ as in (6.2): more precisely, for each edge $e_i \in e(\mathbf{R}_q)$ choose a uniformly random endpoint w_i of e_i . Then $\nu^{\mathbf{n}_q}$ is specified by letting

$$\nu^{\mathbf{n}_q}(V) = \frac{1}{2q-4} \sum_{\{i:w_i \in V\}} n_i$$

for $V \subset v(\mathbf{R}_q)$. (The fact that $2q - 4$ is the correct normalizing constant follows from (8.12).)

If $\max(n_i : 0 \leq i \leq 2r(q) - 4)/(2q - 4) \xrightarrow{\mathbb{P}} 0$ then $|\mathbf{n}_q|_2/|\mathbf{n}_q|_1 \xrightarrow{\mathbb{P}} 0$ and the same argument which led to Corollary 7.2 gives $d_{\mathbb{P}}(\mu_{\mathbf{R}_q}, \nu^{\mathbf{n}_q}) \xrightarrow{\mathbb{P}} 0$. Assuming this holds then just as in (7.5) we obtain

$$d_{\text{GHP}}\left(\left(\frac{9}{8q}\right)^{1/4} \mathbf{R}_q, \left(v(\mathbf{R}_q), \left(\frac{9}{8q}\right)^{1/4} \cdot d_{\mathbf{R}_q}, \nu^{\mathbf{n}_q}\right)\right) \xrightarrow{\mathbb{P}} 0. \quad (8.14)$$

Recall the definition of D' from (5.8). Reprising the argument for (7.8) now gives that for $\varepsilon > 0$,

$$d_{\text{GHP}}((v(\mathbf{R}_q), \varepsilon d_{\mathbf{R}_q}, \nu^{\mathbf{n}_q}), \varepsilon \mathbf{Q}_q) \leq 2\varepsilon \cdot (D'(\mathbf{Q}_q) + 1) + \max(\varepsilon, 1/q). \quad (8.15)$$

This has a very slightly different form from (7.8), where the bound was $2\varepsilon D(\mathbf{R}_r) + \max(\varepsilon, 1/r)$. The reason for the difference is that in the current setting, the submaps of \mathbf{Q}_q pendant to \mathbf{R}_q only attach to one end of an edge of \mathbf{R}_q . When we project the mass to form $\nu^{\mathbf{n}_q}$ we may choose the “wrong end”. This source of error did not appear when projecting mass onto the largest simple block because the 2-connected “decorations” of the largest simple block are attached at *both* endpoints of their respective edges.

At any rate, by Proposition 5.4, $D'(\mathbf{Q}_q)/q^{1/4} \xrightarrow{\mathbb{P}} 0$, so (8.15) and (8.14) together give $d_{\text{GHP}}((\frac{9}{8q})^{1/4} \mathbf{Q}_q, (\frac{9}{8q})^{1/4} \mathbf{R}_q) \xrightarrow{\mathbb{P}} 0$. But by (8.10) we know that the second argument converges to \mathbf{M} , and (8.11) follows.

It thus remains to prove that $\max(n_i : 0 \leq i \leq 2r(q) - 4)/(2q - 4) \xrightarrow{\mathbb{P}} 0$. But this is easy: ℓ_i is the number of copies of a particular edge in \mathbf{Q}_q , so $\max_{0 \leq i \leq 2q-4} \ell_i$ is at most $\max(\deg_{\mathbf{Q}_q}(w) : w \in v(\mathbf{Q}_q))$. By (8.13) we then have

$$\max(n_i : 0 \leq i \leq 2r(q) - 4) \leq 1 + \max(\deg_{\mathbf{Q}_q}(w) : w \in v(\mathbf{Q}_q)) \cdot (2 + \max_{i,j} |e(M_{i,j})|).$$

By Lemma 8.2, the largest degree is at most $\ln^2 q$ with high probability, and Proposition 5.3 gives that $q^{-3/4} \cdot \max_{i,j} |e(M_{i,j})| \leq q^{-3/4} \cdot (2L'(\mathbf{Q}_q) - 4) \xrightarrow{\mathbb{P}} 0$. The result follows. \square

Proof of Theorem 1.2. The theorem follows from Theorem 1.3 in exactly the same way as Theorem 1.1 followed from Proposition 1.5, using Theorem 1.4 in place of Proposition 1.8 for the averaging argument. \square

ACKNOWLEDGEMENTS

LAB was supported by an NSERC Discovery grant and an FQRNT New Researcher grant throughout this research, and thanks both institutions for their support. YW was supported for part of this research by an FQRNT International Internship sponsored by Montréal's Centre de Recherches Mathématiques, and thanks both institutions for their support. Both authors thank the University of Oxford and the Newton Institute, where the final work on this manuscript was completed.

REFERENCES

- [1] Addario-Berry, L., & Albenque, M. (2013). The scaling limit of random simple triangulations and random simple quadrangulations. *arXiv preprint arXiv:1306.5227*. [1](#), [3](#), [4](#), [6](#), [23](#)
- [2] Aldous, D. J. (1985). Exchangeability and related topics (pp. 1-198). Springer Berlin Heidelberg. [21](#)
- [3] Banderier, C., Flajolet, P., Schaeffer, G., & Soria, M. (2001). Random maps, coalescing saddles, singularity analysis, and Airy phenomena. *Random Structures & Algorithms*, 19(3-4), 194-246. [1](#), [3](#), [7](#), [10](#), [11](#), [13](#), [15](#)
- [4] Bender, E. A., & Canfield, E. R. (1989). Face sizes of 3-polytopes. *Journal of Combinatorial Theory, Series B*, 46(1), 58-65. [27](#)
- [5] Burago, D., Burago, Y., & Ivanov, S. (2001). A course in metric geometry (Vol. 33, pp. 371-374). Providence: American Mathematical Society. [9](#)
- [6] Chassaing, P., & Schaeffer, G. (2004). Random planar lattices and integrated superBrownian excursion. *Probability Theory and Related Fields*, 128(2), 161-212. [7](#), [17](#)
- [7] Even, S. (2011). *Graph algorithms*. Cambridge University Press. [2](#)
- [8] Flajolet, P., & Sedgewick, R. (2009). *Analytic combinatorics*. Cambridge University Press. [13](#)
- [9] Gao, Z., & Wormald, N. C. (1999). The size of the largest components in random maps. *SIAM Journal on Discrete Mathematics*, 12(2), 217-228. [3](#), [9](#)
- [10] Gouldn, I.P. & Jackson, D.M. (1983). *Combinatorial Enumeration*. John Wiley & Sons. [13](#)
- [11] Le Gall, J. F. (2013). Uniqueness and universality of the Brownian map. *The Annals of Probability*, 41(4), 2880-2960. [1](#), [3](#), [4](#), [8](#), [9](#), [28](#)
- [12] McDiarmid, C. (1998). Concentration. In *Probabilistic methods for algorithmic discrete mathematics* (pp. 195-248). Springer Berlin Heidelberg. [21](#)
- [13] Miermont, G. (2009). Tessellations of random maps of arbitrary genus. In *Annales Scientifiques de l'cole Normale Suprieure* (Vol. 42, No. 5, pp. 725-781). [9](#)
- [14] Miermont, G. (2013). The Brownian map is the scaling limit of uniform random plane quadrangulations. *Acta mathematica*, 210(2), 319-401. [1](#), [3](#), [4](#), [8](#), [9](#), [28](#)
- [15] Strassen, V. (1965). The existence of probability measures with given marginals. *Ann. Math. Statist.* 36, 423-439. [21](#)
- [16] Tutte, W. T. (1963). A census of planar maps. *Canad. J. Math*, 15(2), 249-271. [13](#), [15](#)
- [17] Walsh, T. R., & Lehman, A. B. (1975). Counting rooted maps by genus III: Nonseparable maps. *Journal of Combinatorial Theory, Series B*, 18(3), 222-259. [15](#)

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCGILL UNIVERSITY, 805 SHERBROOKE STREET WEST, MONTRÉAL, QUÉBEC, H3A 0B9, CANADA

E-mail address: louigi.addario@mcgill.ca, yuting.wen@mail.mcgill.ca

URL: <http://problab.ca/louigi>, <http://www.math.mcgill.ca/ywen>