

LOCALIZATION PRINCIPLE FOR COMPACT HANKEL OPERATORS

ALEXANDER PUSHNITSKI AND DMITRI YAFAEV

ABSTRACT. In the power scale, the asymptotic behavior of the singular values of a compact Hankel operator is determined by the behaviour of the symbol in a neighbourhood of its singular support. In this paper, we discuss the localization principle which says that the contributions of disjoint parts of the singular support to this asymptotic behaviour are independent of each other. We apply this principle to Hankel integral operators and to infinite Hankel matrices. In both cases, we describe a wide class of Hankel operators with power-like asymptotics of singular values. The leading term of this asymptotics is found explicitly.

1. INTRODUCTION AND MAIN RESULTS

1.1. Hankel operators on the unit circle. Hankel operators admit various unitarily equivalent descriptions. We start by recalling the definition of Hankel operators on the Hardy class $H^2(\mathbb{T})$. Here \mathbb{T} is the unit circle in the complex plane, equipped with the normalized Lebesgue measure $dm(\mu) = (2\pi i\mu)^{-1}d\mu$, $\mu \in \mathbb{T}$; the Hardy class $H^2(\mathbb{T}) \subset L^2(\mathbb{T})$ is defined in the standard way as the subspace of $L^2(\mathbb{T})$ spanned by the functions $1, \mu, \mu^2, \dots$. Let $P_+ : L^2(\mathbb{T}) \rightarrow H^2(\mathbb{T})$ be the orthogonal projection onto $H^2(\mathbb{T})$, and let W be the involution in $L^2(\mathbb{T})$ defined by $(Wf)(\mu) = f(\bar{\mu})$. For a function $\omega \in L^\infty(\mathbb{T})$, which is called a *symbol* in this context, the *Hankel operator* $H(\omega)$ is defined by the relation

$$H(\omega)f = P_+(\omega Wf). \quad (1.1)$$

Background information on the theory of Hankel operators can be found e.g. in the books [6, 7].

Recall that the singular values of a compact operator H are defined by the relation $s_n(H) = \lambda_n(|H|)$, where $\{\lambda_n(|H|)\}_{n=1}^\infty$ is the non-increasing sequence of eigenvalues of the compact positive operator $|H| = \sqrt{H^*H}$ (enumerated with multiplicities taken into account). The study of singular values of compact Hankel operators has a long history and is linked to rational approximation, control theory and other subjects, see, e.g. [7]. In fact, this paper is in part motivated by its applications in [12] to the rational approximation of functions with logarithmic singularities.

2010 *Mathematics Subject Classification.* 47B35, 47B06.

Key words and phrases. Hankel operators, singular values, spectral asymptotics.

Singular values $s_n(H(\omega))$ of a Hankel operator with a symbol $\omega \in C^\infty(\mathbb{T})$ decay faster than any power of n^{-1} as $n \rightarrow \infty$. On the other hand, singularities of ω generate a slower decay of singular values. Here we will be interested in the case when the singular values behave as some power of n^{-1} . Optimal upper estimates on singular values of Hankel operators are due to V. Peller, see [7, Section 6.4]. He found necessary and sufficient conditions on ω for the estimate

$$s_n(H(\omega)) \leq Cn^{-\alpha}$$

for some $\alpha > 0$. These conditions are stated in terms of the Besov-Lorentz classes.

It is natural to expect that the asymptotic behavior of singular values is determined by the behaviour of the symbol ω in a neighbourhood of its singular support. We justify this thesis and show that the contributions of the disjoint components of the singular support of ω to the asymptotics of the singular values of $H(\omega)$ are independent of each other. We use the term ‘‘localization principle’’ for this fact. This principle is well understood in the context of the study of the essential spectrum [8] and of the absolutely continuous spectrum [4] of non-compact Hankel operators. Our aim here is to bring this principle to the fore in the question of the asymptotics of singular values of compact Hankel operators.

We combine the localization principle with the results of our previous publication [11] (where the history of the problem is described) to determine the asymptotics of singular values of Hankel operators of various natural classes. In particular, for Hankel matrices with oscillating matrix elements we show that the contributions of different oscillating terms to the asymptotics of singular values are independent of each other. We also establish similar results for Hankel integral operators whose integral kernels have a singularity at some finite point $t_0 \geq 0$ and several oscillating terms at infinity.

1.2. Localization principle. Recall that the singular support $\text{sing supp } \omega$ of a function $\omega \in L^\infty(\mathbb{T})$ is defined as the smallest closed set $X \subset \mathbb{T}$ such that $\omega \in C^\infty(\mathbb{T} \setminus X)$. Localization principle is stated as follows.

Theorem 1.1. *Let $\omega_1, \omega_2, \dots, \omega_L$ be bounded functions on \mathbb{T} such that*

$$\text{sing supp } \omega_\ell \cap \text{sing supp } \omega_j = \emptyset, \quad \ell \neq j. \quad (1.2)$$

Set $\omega = \omega_1 + \dots + \omega_L$. Then for all $p > 0$ we have the relations

$$\limsup_{n \rightarrow \infty} ns_n(H(\omega))^p = \sum_{\ell=1}^L \limsup_{n \rightarrow \infty} ns_n(H(\omega_\ell))^p, \quad (1.3)$$

$$\liminf_{n \rightarrow \infty} ns_n(H(\omega))^p = \sum_{\ell=1}^L \liminf_{n \rightarrow \infty} ns_n(H(\omega_\ell))^p. \quad (1.4)$$

In applications, the upper and lower limits in this theorem usually coincide. However, we prefer to work with these limits separately both because it is more

general and because it is technically more convenient. The limits in the right-hand sides of (1.3), (1.4) may be infinite; in such cases the left-hand sides of (1.3) or (1.4) are also infinite. It is not excluded in Theorem 1.1 that the singular support of each ω_ℓ consists of one point only. In fact, this is exactly this case that we will see in our applications.

1.3. Discussion. Theorem 1.1 can be equivalently stated in terms of the counting functions. For a compact operator H , the singular value counting function is defined by

$$n(\varepsilon; H) = \#\{n : s_n(H) > \varepsilon\}, \quad \varepsilon > 0. \quad (1.5)$$

We have

$$\limsup_{n \rightarrow \infty} n s_n(H)^p = \limsup_{\varepsilon \rightarrow 0} \varepsilon^p n(\varepsilon; H)$$

and similarly for the lower limits. It follows that (1.3), (1.4) may be restated as

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} \varepsilon^p n(\varepsilon; H(\omega)) &= \sum_{\ell=1}^L \limsup_{\varepsilon \rightarrow 0} \varepsilon^p n(\varepsilon; H(\omega_\ell)), \\ \liminf_{\varepsilon \rightarrow 0} \varepsilon^p n(\varepsilon; H(\omega)) &= \sum_{\ell=1}^L \liminf_{\varepsilon \rightarrow 0} \varepsilon^p n(\varepsilon; H(\omega_\ell)). \end{aligned}$$

In particular, in the case when all upper limits in the right-hand sides coincide with the lower limits and are finite, we have

$$n(\varepsilon; H(\omega)) = \sum_{\ell=1}^L n(\varepsilon; H(\omega_\ell)) + o(\varepsilon^{-p}), \quad \varepsilon \rightarrow 0.$$

Our proof of Theorem 1.1 consists of two steps. The first one is to check that under the assumption (1.2) the operators $H(\omega_\ell)$ are *asymptotically orthogonal* in the sense that for all $j \neq \ell$ and all $\alpha > 0$ we have

$$s_n(H(\omega_\ell)^* H(\omega_j)) = O(n^{-\alpha}), \quad s_n(H(\omega_\ell) H(\omega_j)^*) = O(n^{-\alpha}), \quad n \rightarrow \infty. \quad (1.6)$$

This result follows from the reduction of the products of Hankel operators in (1.6) to integral operators in $L^2(\mathbb{T})$ with smooth kernels.

The second step is to show that (1.6) implies relations (1.3) and (1.4). This fact is not specific for Hankel operators. In order to get some intuition into its proof, let us suppose for a moment that the operators $H(\omega_\ell)$ are pairwise *orthogonal* in the sense that

$$H(\omega_j)^* H(\omega_\ell) = 0 \quad \text{and} \quad H(\omega_j) H(\omega_\ell)^* = 0, \quad \forall j \neq \ell. \quad (1.7)$$

Then

$$\text{Ran } H(\omega_j) \perp \text{Ran } H(\omega_\ell) \quad \text{and} \quad \text{Ran } H(\omega_j)^* \perp \text{Ran } H(\omega_\ell)^*, \quad \forall j \neq \ell.$$

Thus, representing the sum $H(\omega) = H(\omega_1) + \dots + H(\omega_L)$ as a “block-diagonal” operator acting from $\oplus_{\ell=1}^L \overline{\text{Ran } H(\omega_\ell)^*}$ to $\oplus_{\ell=1}^L \overline{\text{Ran } H(\omega_\ell)}$, we conclude that

$$n(\varepsilon; H(\omega)) = \sum_{\ell=1}^L n(\varepsilon; H(\omega_\ell)), \quad \forall \varepsilon > 0.$$

Of course, the orthogonality condition (1.7) is too strong. In fact, an operator theoretic result, Theorem 2.2, shows that the asymptotic orthogonality (1.6) ensures the relations (1.3), (1.4) for $p = 1/\alpha$.

Representing Hankel operators in the basis $\{\mu^j\}_{j=0}^\infty$ in $H^2(\mathbb{T})$, one obtains the class of infinite Hankel matrices of the form $\{h(j+k)\}_{j,k=0}^\infty$ in the space $\ell^2(\mathbb{Z}_+)$. We give an application of the localization principle to such Hankel matrices in Theorem 3.1. Although the localization principle in the form stated above (Theorem 1.1) is quite natural, this application looks far less obvious.

Theorem 1.1 can be equivalently stated (see Theorem 2.7) in terms of Hankel operators $\mathbf{H}(\omega)$ acting in the Hardy space $H_+^2(\mathbb{R})$ of functions analytic in the upper half-plane. In this case the symbol $\omega(x)$ is a function of $x \in \mathbb{R}$. This leads to new results for Hankel operators defined as integral operators in the space $L^2(\mathbb{R}_+)$.

We will refer to the Hankel operators in $H^2(\mathbb{T})$ and in $\ell^2(\mathbb{Z}_+)$ as to the discrete case, and to the Hankel operators in $H_+^2(\mathbb{R})$ and in $L^2(\mathbb{R}_+)$ as to the continuous case. We will use boldface font for objects associated with the continuous case. We have tried to make exposition in the discrete and continuous cases parallel as much as possible.

1.4. Related work. Recall that for a bounded operator H , the non-zero parts of the operators

$$(-|H|) \oplus |H| \quad \text{and} \quad \begin{pmatrix} 0 & H \\ H^* & 0 \end{pmatrix}$$

are unitarily equivalent. Therefore various spectral results for $|H(\omega)|$ are equivalent to those for the self-adjoint Hankel operator with the matrix valued symbol

$$\begin{pmatrix} 0 & \omega(\mu) \\ \omega(\bar{\mu}) & 0 \end{pmatrix}.$$

Some forms of localization principle are known in the study of the continuous spectrum of $|H(\omega)|$. As far as we are aware, the idea of separation of singularities of the symbol goes back to the work [8] of S. R. Power on the essential spectrum spec_{ess} of Hankel operators with piecewise continuous symbols ω . Let $a_j \in \mathbb{T}$ be the points where ω has the jumps

$$\varkappa(a_j) = \lim_{\varepsilon \rightarrow +0} \omega(a_j e^{i\varepsilon}) - \lim_{\varepsilon \rightarrow +0} \omega(a_j e^{-i\varepsilon}) \neq 0.$$

Although Power was interested in the essential spectrum of $H(\omega)$ (which we do not discuss here), it follows from the matrix version of his results that

$$\text{spec}_{\text{ess}}(|H(\omega)|) = [0, M], \quad M = \frac{1}{2} \sup_{a_j \in \mathbb{T}} |\mathcal{Z}(a_j)|, \quad (1.8)$$

where the supremum is taken over all points a_j where ω has a jump.

A description of the absolutely continuous spectrum of $|H(\omega)|$ with piecewise continuous symbol ω follows from the matrix version of the results of Howland [4], where the trace class method of scattering theory was used. This question was also studied in our previous paper [9] by using the so-called smooth method of scattering theory. In both cases, under some mild additional assumptions, including the condition that ω has finitely many jumps, it can be shown that

$$\text{spec}_{\text{ac}}(|H(\omega)|) = \bigcup_{a_j \in \mathbb{T}} [0, \frac{1}{2} |\mathcal{Z}(a_j)|]. \quad (1.9)$$

Every term in the right-hand side of (1.9) gives its own band of the absolutely continuous spectrum of multiplicity one. Thus, formula (1.9) can be regarded as the continuous spectrum analogue of the localisation principle discussed in this paper: the contributions of different jumps of ω to $\text{spec}_{\text{ac}}(|H(\omega)|)$ are independent of each other. Of course, formulas (1.8) and (1.9) are consistent with each other.

1.5. The structure of the paper. In Section 2 we prove the localization principle in the discrete case (Theorem 1.1) and also state and prove its analogue in the continuous case (Theorem 2.7). In Section 3, we describe the applications of localization principle to the Hankel operators acting in $\ell^2(\mathbb{Z}_+)$; the proofs are given in Section 4. In Section 5 we give applications to integral Hankel operators in $L^2(\mathbb{R}_+)$; the proofs are given in Section 6. In Section 7 we consider integral Hankel operators with kernels with local singularities in \mathbb{R}_+ .

1.6. Some notation. For $\omega \in L^2(\mathbb{T})$, the Fourier coefficients of ω are denoted as usual by

$$\widehat{\omega}(j) = \int_{\mathbb{T}} \omega(\mu) \mu^{-j} dm(\mu), \quad j \in \mathbb{Z}.$$

We will consistently make use of the following constant, which appears in our asymptotic formulas:

$$v(\alpha) = 2^{-\alpha} \pi^{1-2\alpha} \left(B\left(\frac{1}{2\alpha}, \frac{1}{2}\right) \right)^\alpha, \quad \alpha > 0; \quad (1.10)$$

here $B(\cdot, \cdot)$ is the Beta function. We make a standing assumption that the exponents $p > 0$ and $\alpha > 0$ are related by $\alpha = 1/p$.

2. PROOF OF LOCALIZATION PRINCIPLE

In this section, we prove Theorem 1.1. We also prove a similar statement, Theorem 2.7, for Hankel operators in the Hardy space $H_+^2(\mathbb{R})$ of functions analytic in the upper half-plane.

2.1. Preliminaries. Let \mathcal{B} be the algebra of bounded operators in a Hilbert space \mathcal{H} , and let \mathbf{S}_∞ be the ideal of compact operators in \mathcal{B} . For $p > 0$, the weak Schatten class $\mathbf{S}_{p,\infty}$ consists of all compact operators A such that

$$\sup_n ns_n(A)^p < \infty.$$

The subclass $\mathbf{S}_{p,\infty}^0 \subset \mathbf{S}_{p,\infty}$ is defined by the condition

$$\lim_{n \rightarrow \infty} ns_n(A)^p = 0.$$

It is well known that both $\mathbf{S}_{p,\infty}$ and $\mathbf{S}_{p,\infty}^0$ are ideals of \mathcal{B} ; in particular, they are linear spaces. Of course $A \in \mathbf{S}_{p,\infty}$ (or $A \in \mathbf{S}_{p,\infty}^0$) if and only if the same is true for its adjoint A^* . We set $\mathbf{S}_0 = \bigcap_{p>0} \mathbf{S}_{p,\infty}$, that is,

$$A \in \mathbf{S}_0 \iff s_n(A) = O(n^{-\alpha}), \quad n \rightarrow \infty, \quad \forall \alpha > 0. \quad (2.1)$$

First we recall a classical result in perturbation theory (see e.g. [1, Theorem 11.6.8]) on the spectral stability of singular values.

Lemma 2.1. *Let $A \in \mathbf{S}_\infty$ and $B \in \mathbf{S}_{p,\infty}^0$ for some $p > 0$. Then*

$$\limsup_{n \rightarrow \infty} ns_n(A+B)^p = \limsup_{n \rightarrow \infty} ns_n(A)^p, \quad (2.2)$$

$$\liminf_{n \rightarrow \infty} ns_n(A+B)^p = \liminf_{n \rightarrow \infty} ns_n(A)^p. \quad (2.3)$$

Lemma 2.1 is stated in a slightly more general form than usual (see, e.g., Theorem 11.6.8 in [1]) because we do not require that $A \in \mathbf{S}_{p,\infty}$ and hence the limits in (2.2) and (2.3) may be infinite; in this case Lemma 2.1 means that both sides in (2.2) and (2.3) are infinite simultaneously. Note that if $A \notin \mathbf{S}_{p,\infty}$, then the expression (2.2) is infinite, but the expression (2.3) may be finite. Lemma 2.1 can also be equivalently stated in terms of the singular value counting functions $n(\varepsilon, A)$ defined by (1.5).

2.2. Asymptotically orthogonal operators. Note the implication

$$A \in \mathbf{S}_{p,\infty}, \quad B \in \mathbf{S}_{p,\infty} \implies A^*B \in \mathbf{S}_{p/2,\infty}, \quad AB^* \in \mathbf{S}_{p/2,\infty} \quad (2.4)$$

(see, e.g. [1, Theorem 11.6.9]). We say that the operators A and B in $\mathbf{S}_{p,\infty}$ are asymptotically orthogonal if the class $\mathbf{S}_{p/2,\infty}$ in the right side of (2.4) can be replaced by its subclass $\mathbf{S}_{p/2,\infty}^0$. The following theorem allows us to study singular values of sums of asymptotically orthogonal operators. This result is the key operator theoretic ingredient of our construction.

Theorem 2.2. *Let $p > 0$. Assume that $A_1, \dots, A_L \in \mathbf{S}_\infty$ and*

$$A_\ell^*A_j \in \mathbf{S}_{p/2,\infty}^0, \quad A_\ell A_j^* \in \mathbf{S}_{p/2,\infty}^0 \quad \text{for all } \ell \neq j. \quad (2.5)$$

Then for $A = A_1 + \cdots + A_L$, we have

$$\limsup_{n \rightarrow \infty} n s_n(A)^p = \sum_{\ell=1}^L \limsup_{n \rightarrow \infty} n s_n(A_\ell)^p, \quad (2.6)$$

$$\liminf_{n \rightarrow \infty} n s_n(A)^p = \sum_{\ell=1}^L \liminf_{n \rightarrow \infty} n s_n(A_\ell)^p. \quad (2.7)$$

Proof. Let us prove the first relation (2.6); the second one is proven in the same way. We argue in terms of counting functions (1.5). For an operator $A \in \mathbf{S}_\infty$, let us denote

$$\Delta_p(A) = \limsup_{\varepsilon \rightarrow 0} \varepsilon^{1/p} n(\varepsilon; A)$$

(this limit may be infinite). Then our aim is to prove that

$$\Delta_p(A) = \sum_{\ell=1}^L \Delta_p(A_\ell), \quad (2.8)$$

which is (2.6) in different notation. Put

$$\mathcal{H}^L = \underbrace{\mathcal{H} \oplus \cdots \oplus \mathcal{H}}_{L \text{ terms}}$$

and let $A_0 = \text{diag}\{A_1, \dots, A_L\}$ in \mathcal{H}^L , i.e.,

$$A_0(f_1, \dots, f_L) = (A_1 f_1, \dots, A_L f_L).$$

Since

$$A_0^* A_0 = \text{diag}\{A_1^* A_1, \dots, A_L^* A_L\}, \quad (2.9)$$

we see that

$$n(\varepsilon; A_0) = \sum_{\ell=1}^L n(\varepsilon; A_\ell)$$

and therefore

$$\Delta_{p/2}(A_0^* A_0) = \sum_{\ell=1}^L \Delta_{p/2}(A_\ell^* A_\ell) = \sum_{\ell=1}^L \Delta_p(A_\ell). \quad (2.10)$$

Next, let $J : \mathcal{H}^L \rightarrow \mathcal{H}$ be the operator given by

$$J(f_1, \dots, f_L) = f_1 + \cdots + f_L \quad \text{so that} \quad J^* f = (f, \dots, f).$$

Then

$$J A_0(f_1, \dots, f_L) = A_1 f_1 + \cdots + A_L f_L$$

and

$$(J A_0)^* f = (A_1^* f, \dots, A_L^* f).$$

It follows that

$$(J A_0)(J A_0)^* f = (A_1 A_1^* + \cdots + A_L A_L^*) f \quad (2.11)$$

and the operator $(JA_0)^*(JA_0)$ is a “matrix” in \mathcal{H}^L given by

$$(JA_0)^*(JA_0) = \begin{pmatrix} A_1^*A_1 & A_1^*A_2 & \dots & A_1^*A_L \\ A_2^*A_1 & A_2^*A_2 & \dots & A_2^*A_L \\ \vdots & \vdots & \ddots & \vdots \\ A_L^*A_1 & A_L^*A_2 & \dots & A_L^*A_L \end{pmatrix}. \quad (2.12)$$

According to (2.9) and (2.12) we have

$$(JA_0)^*(JA_0) - A_0^*A_0 \in \mathbf{S}_{p/2, \infty}^0. \quad (2.13)$$

Indeed, the “matrix” of the operator in (2.13) has zeros on the diagonal, and its off-diagonal elements are given by $A_\ell^*A_j$, $\ell \neq j$. Thus (2.13) follows from the first assumption (2.5). Therefore Lemma 2.1 implies that

$$\Delta_{p/2}((JA_0)^*(JA_0)) = \Delta_{p/2}(A_0^*A_0)$$

or

$$\Delta_{p/2}((JA_0)(JA_0)^*) = \Delta_{p/2}(A_0^*A_0) \quad (2.14)$$

because for any compact operator T the non-zero singular values of T^*T and TT^* coincide.

Further, since $AA^* = \sum_{\ell, j=1}^L A_\ell A_j^*$, it follows from (2.11) and the second assumption (2.5) that

$$AA^* - (JA_0)(JA_0)^* = \sum_{j \neq \ell} A_\ell A_j^* \in \mathbf{S}_{p/2, \infty}^0.$$

Using again Lemma 2.1, from here we obtain

$$\Delta_p(A) = \Delta_{p/2}(AA^*) = \Delta_{p/2}((JA_0)(JA_0)^*).$$

Combining the last equality with (2.14), we see that $\Delta_p(A) = \Delta_{p/2}(A_0^*A_0)$. Thus (2.10) yields the relation (2.8). \square

Corollary 2.3. *Under the assumption (2.5) we have*

$$\lim_{n \rightarrow \infty} ns_n(A)^p = \sum_{\ell=1}^L \lim_{n \rightarrow \infty} ns_n(A_\ell)^p$$

provided the limits in the right-hand side exist.

Under slightly more restrictive assumptions Theorem 2.2 appeared first in [2, Theorem 3]. Our proof is quite different from that of [2].

Remark 2.4. Let us mention two known statements that are similar in spirit to Theorem 2.2. Below A_1, \dots, A_L are bounded operators and $A = A_1 + \dots + A_L$.

(i) If the products $A_\ell^*A_j$, $A_\ell A_j^*$ are compact for all $j \neq \ell$, then for the essential spectra of A one has the formula

$$\text{spec}_{\text{ess}}(A) \cup \{0\} = \bigcup_{\ell=1}^L \text{spec}_{\text{ess}}(A_\ell),$$

see, e.g. [7, Section 10.1].

(ii) If A_1, \dots, A_L are bounded self-adjoint operators such that $A_\ell A_j$ are trace class for all $j \neq \ell$, then the absolutely continuous part of A is unitarily equivalent to the orthogonal sum of the absolutely continuous parts of the operators A_ℓ . This is known as Ismagilov's theorem, see [5].

2.3. Proof of localization principle for Hankel operators in $H^2(\mathbb{T})$. First we state two well-known facts that will be needed below. We recall that the Hankel operators $H(\omega)$ are defined by (1.1); the class \mathbf{S}_0 is defined by (2.1).

Lemma 2.5. (i) *Let K be an integral operator in $L^2(\mathbb{T})$ with an integral kernel of the class $C^\infty(\mathbb{T} \times \mathbb{T})$. Then $K \in \mathbf{S}_0$.*

(ii) *Let $\omega \in C^\infty(\mathbb{T})$; then $H(\omega) \in \mathbf{S}_0$.*

Proof. Part (i) is a classical fact; it can be obtained, for example, by approximating the integral kernel of K by trigonometric polynomials. This yields a fast approximation of K by finite rank operators.

Part (ii) is also well-known; let us show that it follows from part (i). It will be convenient to consider the projection P_+ here as an operator acting from $L^2(\mathbb{T})$ to $L^2(\mathbb{T})$ (rather than from $L^2(\mathbb{T})$ to $H^2(\mathbb{T})$). Recall that P_+ acts according to the formula

$$(P_+f)(\mu) = \lim_{\epsilon \rightarrow +0} \int_{\mathbb{T}} \frac{f(\mu')}{\mu' - (1 - \epsilon)\mu} \mu' dm(\mu'), \quad (2.15)$$

and that W is the involution $(Wf)(\mu) = f(\bar{\mu})$. We have to prove that the operator $P_+\omega WP_+$ in $L^2(\mathbb{T})$ belongs to the class \mathbf{S}_0 . Since P_+WP_+ is a rank one operator (projection onto constants), it suffices to check that

$$P_+\omega WP_+ - \omega P_+WP_+ = [P_+, \omega]WP_+ \in \mathbf{S}_0. \quad (2.16)$$

It follows from (2.15) that the commutator $[P_+, \omega]$ is an integral operator in $L^2(\mathbb{T})$ with the kernel

$$\frac{\omega(\mu') - \omega(\mu)}{\mu' - \mu} \mu', \quad \mu, \mu' \in \mathbb{T}.$$

This is a C^∞ function, and so $[P_+, \omega] \in \mathbf{S}_0$ which implies (2.16). \square

The following assertion allows us to separate the contributions of different singularities of the symbol. Essentially, this is a very well known argument, see, e.g. [8].

Lemma 2.6. *Let $\omega_1, \omega_2 \in L^\infty(\mathbb{T})$ be such that $\text{sing supp } \omega_1 \cap \text{sing supp } \omega_2 = \emptyset$. Then*

$$H(\omega_1)^*H(\omega_2) \in \mathbf{S}_0, \quad H(\omega_1)H(\omega_2)^* \in \mathbf{S}_0.$$

Proof. Let \varkappa_1, \varkappa_2 be functions in $C^\infty(\mathbb{T})$ with disjoint supports such that

$$(1 - \varkappa_k)\omega_k \in C^\infty(\mathbb{T}), \quad k = 1, 2.$$

By Lemma 2.5(ii), we have

$$H((1 - \varkappa_k)\omega_k) \in \mathbf{S}_0,$$

and hence it suffices to show that

$$H(\varkappa_1\omega_1)^*H(\varkappa_2\omega_2) \in \mathbf{S}_0, \quad H(\varkappa_1\omega_1)H(\varkappa_2\omega_2)^* \in \mathbf{S}_0. \quad (2.17)$$

It follows from definition (1.1) that

$$H(\varkappa_1\omega_1)^*H(\varkappa_2\omega_2)f = P_+W\overline{\omega_1}(\overline{\varkappa_1}P_+\varkappa_2)\omega_2Wf, \quad f \in H^2(\mathbb{T}).$$

Since the supports of \varkappa_1 and \varkappa_2 are disjoint, the operator $\overline{\varkappa_1}P_+\varkappa_2$ has a C^∞ -smooth integral kernel

$$\frac{\overline{\varkappa_1(\mu)}\varkappa_2(\mu')}{\mu' - \mu}\mu', \quad \mu, \mu' \in \mathbb{T},$$

and so by Lemma 2.5(i) it belongs to the class \mathbf{S}_0 . This ensures the first inclusion in (2.17). In view of the obvious identity

$$H(\omega)^* = H(\omega_*) \quad \text{where} \quad \omega_*(\mu) = \overline{\omega(\bar{\mu})},$$

the second inclusion (2.17) follows from the first one. \square

Proof of Theorem 1.1. Let us apply the abstract Theorem 2.2 to the Hankel operators $A_\ell = H(\omega_\ell)$, $\ell = 1, \dots, L$. Lemma 2.6 implies that the asymptotic orthogonality condition (2.5) is satisfied. Therefore the asymptotic relations (1.3) and (1.4) follow directly from (2.6) and (2.7). \square

2.4. Hankel operators in $H_+^2(\mathbb{R})$. Hankel operators can also be defined in the Hardy space $H_+^2(\mathbb{R})$ of functions analytic in the upper half-plane. We denote by Φ the unitary Fourier transform on $L^2(\mathbb{R})$,

$$\widehat{u}(t) = (\Phi u)(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} u(x)e^{-ixt} dx.$$

Let $H_+^2(\mathbb{R}) \subset L^2(\mathbb{R})$ be the Hardy class,

$$H_+^2(\mathbb{R}) = \{u \in L^2(\mathbb{R}) : \widehat{u}(t) = 0 \text{ for } t < 0\},$$

and let $\mathbf{P}_+ : L^2(\mathbb{R}) \rightarrow H_+^2(\mathbb{R})$ be the corresponding orthogonal projection. Let \mathbf{W} be the involution in $L^2(\mathbb{R})$, $(\mathbf{W}f)(x) = f(-x)$. For $\omega \in L^\infty(\mathbb{R})$, the operator $\mathbf{H}(\omega)$ in $H_+^2(\mathbb{R})$ is defined by the formula

$$\mathbf{H}(\omega)f = \mathbf{P}_+(\omega\mathbf{W}f), \quad f \in H_+^2(\mathbb{R}). \quad (2.18)$$

There is a unitary equivalence between the Hankel operators $H(\omega)$ defined in $H^2(\mathbb{T})$ by formula (1.1) and the Hankel operators $\mathbf{H}(\omega)$ defined in $H_+^2(\mathbb{R})$ by formula (2.18). Indeed, let

$$w = \frac{z - i/2}{z + i/2}, \quad z = \frac{i}{2} \frac{1 + w}{1 - w} \quad (2.19)$$

be the standard conformal map sending the real line onto the unit circle, and let $\mathcal{U} : H^2(\mathbb{T}) \rightarrow H_+^2(\mathbb{R})$ be the corresponding unitary operator defined by

$$(\mathcal{U}f)(x) = \frac{1}{\sqrt{2\pi}} \frac{1}{x+i/2} f\left(\frac{x-i/2}{x+i/2}\right), \quad (\mathcal{U}^*\mathbf{f})(\mu) = i\sqrt{2\pi} \frac{1}{1-\mu} \mathbf{f}\left(\frac{i}{2} \frac{1+\mu}{1-\mu}\right).$$

Then

$$\mathcal{U}H(\omega)\mathcal{U}^* = \mathbf{H}(\omega), \quad \text{if} \quad \omega(x) = -\frac{x-i/2}{x+i/2} \omega\left(\frac{x-i/2}{x+i/2}\right). \quad (2.20)$$

So the localization principle stated for $H(\omega)$ can be automatically mapped to operators $\mathbf{H}(\omega)$. This is discussed below.

2.5. Localization principle in $H_+^2(\mathbb{R})$. Symbols $\omega(x)$ of Hankel operators $\mathbf{H}(\omega)$ have the exceptional points $x = +\infty$ and $x = -\infty$; it will be convenient to identify these two points. The real line with such identification will be denoted \mathbb{R}_* . We write $\omega \in C(\mathbb{R}_*)$ if $\omega \in C(\mathbb{R})$ and

$$\lim_{x \rightarrow \infty} \omega(x) = \lim_{x \rightarrow -\infty} \omega(x),$$

where both limits are supposed to exist. Similarly, we write $\omega \in C^\infty(\mathbb{R}_*)$ if $\omega \in C^\infty(\mathbb{R})$ and, for all $m = 0, 1, \dots$,

$$\lim_{x \rightarrow \infty} \omega^{(m)}(x) = \lim_{x \rightarrow -\infty} \omega^{(m)}(x). \quad (2.21)$$

In particular, the point $x = \infty$ belongs to the singular support of ω if for at least one $m \geq 0$ the relation (2.21) fails (i.e. if one of the limits does not exist or if the limits are not equal).

Let us state the localization principle for Hankel operators in $H_+^2(\mathbb{R})$.

Theorem 2.7. *Let $\omega_\ell \in L^\infty(\mathbb{R})$, $\ell = 1, \dots, L < \infty$, be such that*

$$\text{sing supp } \omega_\ell \cap \text{sing supp } \omega_j = \emptyset, \quad \ell \neq j.$$

Set $\omega = \omega_1 + \dots + \omega_L$. Then for all $p > 0$ we have the relations

$$\begin{aligned} \limsup_{n \rightarrow \infty} n s_n(\mathbf{H}(\omega))^p &= \sum_{\ell=1}^L \limsup_{n \rightarrow \infty} n s_n(\mathbf{H}(\omega_\ell))^p, \\ \liminf_{n \rightarrow \infty} n s_n(\mathbf{H}(\omega))^p &= \sum_{\ell=1}^L \liminf_{n \rightarrow \infty} n s_n(\mathbf{H}(\omega_\ell))^p. \end{aligned}$$

Observe that formulas (2.19) establish a one-to-one correspondence between the unit circle \mathbb{T} and the real axis \mathbb{R}_* with the points $x = +\infty$ and $x = -\infty$ identified. They yield also the one-to-one correspondence between the singular supports of the symbols $\omega(\mu)$ and $\omega(x)$ linked by equality (2.20). Thus, Theorem 2.7 is a direct consequence of Theorem 1.1.

3. APPLICATIONS OF LOCALIZATION PRINCIPLE: DISCRETE CASE

3.1. Discrete representation. For a sequence $\{h(j)\}_{j=0}^{\infty}$ of complex numbers, the Hankel operator $\Gamma(h)$ in the space $\ell^2(\mathbb{Z}_+)$ is formally defined by the “infinite matrix” $\{h(j+k)\}_{j,k=0}^{\infty}$:

$$(\Gamma(h)u)(j) = \sum_{k=0}^{\infty} h(j+k)u(k), \quad u = \{u(k)\}_{k=0}^{\infty}. \quad (3.1)$$

The Hankel operators $\Gamma(h)$ in $\ell^2(\mathbb{Z}_+)$ and $H(\omega)$ in $H^2(\mathbb{T})$ are related as follows. Let

$$\mathcal{F} : f \mapsto \{\widehat{f}(j)\}_{j=0}^{\infty}, \quad \mathcal{F} : H^2(\mathbb{T}) \rightarrow \ell^2(\mathbb{Z}_+),$$

be the discrete Fourier transform. Then the matrix elements of $H(\omega)$ in the orthonormal basis $\{\mu^j\}_{j=0}^{\infty}$ are

$$(H(\omega)\mu^j, \mu^k)_{L^2(\mathbb{T})} = \widehat{\omega}(j+k), \quad j, k \geq 0,$$

so that

$$\Gamma(h) = \mathcal{F}H(\omega)\mathcal{F}^* \quad \text{if} \quad \widehat{\omega}(j) = h(j), \quad j \geq 0. \quad (3.2)$$

Since (3.2) involves only the coefficients with $j \geq 0$, for a given sequence h the symbol ω is not uniquely defined.

3.2. Plan of the approach. In our previous publication [11] we considered compact *self-adjoint* Hankel operators, corresponding to sequences of real numbers of the type

$$q(j) = j^{-1}(\log j)^{-\alpha} + \text{error term}, \quad j \rightarrow \infty, \quad (3.3)$$

where $\alpha > 0$. Under the appropriate assumptions on the error term, we proved in [11] that the positive eigenvalues of the Hankel operator $\Gamma(q)$ have the asymptotics

$$\lambda_n^+(\Gamma(q)) = v(\alpha)n^{-\alpha} + o(n^{-\alpha}), \quad n \rightarrow \infty,$$

where the coefficient $v(\alpha)$ is defined in (1.10). For negative eigenvalues, we have $\lambda_n^-(\Gamma(q)) = o(n^{-\alpha})$ as $n \rightarrow \infty$.

In [11] our analysis was based on the asymptotic form (3.3) and did not involve symbols directly. In this paper, we check (this is an easy calculation, see Lemma 4.3 below) that if $q(j) = j^{-1}(\log j)^{-\alpha}$, then a symbol σ of $\Gamma(q)$ can be chosen such that $\text{sing supp } \sigma = \{1\}$.

Theorem 1.1 allows us to find the asymptotics of singular values for more general “oscillating” sequences of the type

$$h(j) = \sum_{\ell=1}^L b_{\ell} j^{-1} (\log j)^{-\alpha} \zeta_{\ell}^{-j} + \text{error term}, \quad j \rightarrow \infty, \quad (3.4)$$

where $\zeta_1, \dots, \zeta_L \in \mathbb{T}$ are distinct points and $b_1, \dots, b_L \in \mathbb{C}$ are arbitrary coefficients. It is easy to see that the symbol corresponding to the ℓ 'th term in (3.4) equals $b_{\ell}\sigma(\mu/\zeta_{\ell})$. Hence its singular support consists of one point $\{\zeta_{\ell}\}$, and so we

are in the situation described by the localization principle for $p = 1/\alpha$. The error term in (3.4) is treated by using the estimates from [10] on singular values of Hankel operators.

Notice that the operators $\Gamma(h)$ corresponding to sequences h of the class (3.4) are in general not self-adjoint. We have information about the asymptotics of their singular values, but not of their eigenvalues.

3.3. Main result in the discrete case. In order to state our requirements on the error term in (3.4), we need some notation. Let

$$M(\alpha) = \begin{cases} [\alpha] + 1, & \text{if } \alpha \geq 1/2, \\ 0, & \text{if } \alpha < 1/2, \end{cases} \quad (3.5)$$

where $[\alpha]$ is the integer part of α . For a sequence $h = \{h(j)\}_{j=0}^{\infty}$, we define iteratively the sequences $h^{(m)} = \{h^{(m)}(j)\}_{j=0}^{\infty}$, $m = 0, 1, 2, \dots$, by setting $h^{(0)}(j) = h(j)$ and

$$h^{(m+1)}(j) = h^{(m)}(j+1) - h^{(m)}(j), \quad j \geq 0. \quad (3.6)$$

Note that if $h(j) = j^{-1}(\log j)^{-\alpha}$ for sufficiently large j , then for all $m \geq 1$ the sequences $h^{(m)}$ satisfy

$$h^{(m)}(j) = O(j^{-1-m}(\log j)^{-\alpha}), \quad j \rightarrow \infty. \quad (3.7)$$

Now we are in a position to state precisely our result on Hankel operators with matrix elements (3.4).

Theorem 3.1. *Let $\alpha > 0$, let $\zeta_1, \dots, \zeta_L \in \mathbb{T}$ be distinct numbers, and let $b_1, \dots, b_L \in \mathbb{C}$. Let h be a sequence of complex numbers such that*

$$h(j) = \sum_{\ell=1}^L (b_{\ell} j^{-1} (\log j)^{-\alpha} + g_{\ell}(j)) \zeta_{\ell}^{-j}, \quad j \geq 2, \quad (3.8)$$

where the error terms g_{ℓ} , $\ell = 1, \dots, L$, satisfy the estimates

$$g_{\ell}^{(m)}(j) = o(j^{-1-m}(\log j)^{-\alpha}), \quad j \rightarrow \infty, \quad (3.9)$$

for all $m = 0, 1, \dots, M(\alpha)$ ($M(\alpha)$ is given by (3.5)). Then the singular values of the Hankel operator $\Gamma(h)$ in $\ell^2(\mathbb{Z}_+)$ defined by formula (3.1) satisfy the asymptotic relation

$$s_n(\Gamma(h)) = c n^{-\alpha} + o(n^{-\alpha}), \quad n \rightarrow \infty, \quad (3.10)$$

where

$$c = v(\alpha) \left(\sum_{\ell=1}^L |b_{\ell}|^{1/\alpha} \right)^{\alpha} \quad (3.11)$$

and the coefficient $v(\alpha)$ is given by formula (1.10).

This result means that asymptotically the singular value counting function of the operator $\Gamma(h)$ is the sum of such functions for every term in the right-hand side of (3.8).

4. PROOF OF THEOREM 3.1

4.1. Singular value estimates and asymptotics. We need two results obtained in our papers [10, 11]. Let $M(\alpha)$ be as in (3.5).

Theorem 4.1. [10, Theorem 2.3] *Suppose that a sequence g satisfies*

$$g^{(m)}(j) = o(j^{-1-m}(\log j)^{-\alpha}), \quad j \rightarrow \infty, \quad (4.1)$$

for some $\alpha > 0$ and for all $m = 0, 1, \dots, M(\alpha)$. Then

$$s_n(\Gamma(g)) = o(n^{-\alpha}), \quad n \rightarrow \infty. \quad (4.2)$$

In [10] we also have a result with O instead of o in both (4.1) and (4.2), but we do not use it in this paper. Observe that for $\alpha < 1/2$ we need only the estimate on g , whereas for $\alpha \geq 1/2$ we also need estimates on the iterated differences $g^{(m)}$.

Theorem 4.2. [11, Theorem 1.1] *Let $\alpha > 0$, and let the “model sequence” q be defined by*

$$q(j) = j^{-1}(\log j)^{-\alpha} \quad (4.3)$$

for all sufficiently large j (the values $q(j)$ for any finite number of j are unimportant). Then

$$s_n(\Gamma(q)) = v(\alpha)n^{-\alpha} + o(n^{-\alpha}), \quad n \rightarrow \infty,$$

where $v(\alpha)$ is given by (1.10).

Of course, this result corresponds to a particular case of Theorem 3.1 with $L = 1$, $\zeta_1 = 1$, $b_1 = 1$.

4.2. The model symbol. In order to combine the contributions of different terms in (3.8), we use the localization principle (i.e. Theorem 1.1). To that end, we have to identify the singular support of the symbol corresponding to the model sequence q in Theorem 4.2. To be definite, we put $q(0) = q(1) = 0$ and define $q(j)$ by formula (4.3) for all $j \geq 2$. We need to find a function σ such that its Fourier coefficients $\widehat{\sigma}(j) = q(j)$ for $j \geq 0$. Of course, the choice of σ is not unique. We will choose σ corresponding to the *odd* extension of the sequence $q(j)$ to the negative j .

Lemma 4.3. *Let $\alpha \geq 0$, and let q be given by (4.3); set*

$$\sigma(\mu) = \sum_{j=2}^{\infty} q(j)(\mu^j - \bar{\mu}^j), \quad \mu \in \mathbb{T}. \quad (4.4)$$

Then $\sigma \in L^\infty(\mathbb{T})$ and $\sigma \in C^\infty(\mathbb{T} \setminus \{1\})$.

Proof. Note that for all $\mu \in \mathbb{T}$, the series (4.4) converges absolutely if $\alpha > 1$ and conditionally if $\alpha \leq 1$.

First, we check that $\sigma \in L^\infty(\mathbb{T})$. We write $\mu = e^{i\theta}$, $\theta \in (-\pi, \pi]$. For $\theta \neq 0$, we set $N = \lceil (2|\theta|)^{-1} \rceil$ and write $\sigma = \sigma_1 + \sigma_2$, where

$$\sigma_1(\mu) = \sum_{j=2}^N q(j)(\mu^j - \bar{\mu}^j), \quad \sigma_2(\mu) = \sum_{j=N+1}^{\infty} q(j)(\mu^j - \bar{\mu}^j).$$

Using the bounds $q(j) \leq (\log 2)^{-1}j^{-1}$ and

$$|\mu^j - \bar{\mu}^j| = 2|\sin(j\theta)| \leq 2j|\theta|$$

for σ_1 , we obtain the estimate

$$|\sigma_1(\mu)| \leq 2|\theta| \sum_{j=2}^N jq(j) \leq 2(\log 2)^{-1}|\theta|N \leq (\log 2)^{-1}.$$

In order to estimate σ_2 , let us use summation by parts:

$$\begin{aligned} (\mu - 1) \sum_{j=N+1}^{\infty} q(j)\mu^j &= \sum_{j=N+1}^{\infty} q(j)(\mu^{j+1} - \mu^j) \\ &= - \sum_{j=N+1}^{\infty} q^{(1)}(j)\mu^{j+1} - q(N+1)\mu^{N+1} \end{aligned} \quad (4.5)$$

where $q^{(1)}(j)$ is defined by (3.6). By (3.7), we have $q^{(1)}(j) = O(j^{-2})$, $j \rightarrow \infty$, and hence

$$\left| (\mu - 1) \sum_{j=N+1}^{\infty} q(j)\mu^j \right| \leq C_1 \left(\sum_{j=N+1}^{\infty} j^{-2} + N^{-1} \right) \leq C_2 N^{-1}.$$

It follows that

$$|\sigma_2(\mu)| \leq 2 \left| \sum_{j=N+1}^{\infty} q(j)\mu^j \right| \leq \frac{2C_2}{N|\mu - 1|} = \frac{2C_2}{[(2|\theta|)^{-1}]|e^{i\theta} - 1|} \leq C.$$

Thus $\sigma_2 \in L^\infty(\mathbb{T})$.

It remains to prove that $\sigma \in C^M(\mathbb{T} \setminus \{1\})$ for any $M \in \mathbb{N}$. Choose $\mu \in \mathbb{T}$ and put $a(j) = \mu^j$; then, by definition (3.6), $a^{(M+1)}(j) = (\mu - 1)^{M+1}\mu^j$. Similarly to (4.5), by a repeated summation by parts procedure, we obtain the identity

$$\begin{aligned} (\mu - 1)^{M+1} \sum_{j=2}^{\infty} q(j)\mu^j &= \sum_{j=2}^{\infty} q(j)a^{(M+1)}(j) \\ &= (-1)^{M+1} \sum_{j=2}^{\infty} q^{(M+1)}(j)a(j) + p_M(\mu) \end{aligned} \quad (4.6)$$

with some polynomial p_M . Since, by (3.7), $q^{(M+1)}(j) = O(j^{-2-M})$ as $j \rightarrow \infty$ and $a(j) = \mu^j$, the function of μ in the right-hand side of (4.6) is in $C^M(\mathbb{T})$. It follows that $\sigma \in C^M(\mathbb{T} \setminus \{1\})$ and hence $\sigma \in C^\infty(\mathbb{T} \setminus \{1\})$. \square

Remark 4.4. (i) It can be verified that the singular support of σ is non-empty, that is, $1 \in \text{sing supp } \sigma$. In fact, a more detailed calculation shows that

$$\sigma(e^{i\theta}) = \pi i \text{sign } \theta |\log|\theta||^{-\alpha} (1 + o(1)), \quad \theta \rightarrow 0.$$

(ii) If $\alpha > 1$, then instead of the odd extension of $q(j)$ to the negative j , one can extend it by zero, i.e. one can choose

$$\tilde{\sigma}(\mu) = \sum_{j=2}^{\infty} q(j) \mu^j.$$

This doesn't work for $\alpha \leq 1$ since $\tilde{\sigma}(\mu)$ is unbounded as $\mu \rightarrow 1$ in this case.

According to definition (4.4), we have $\hat{\sigma}(j) = q(j)$ for all $j \geq 0$. Hence, it follows from relation (3.2) that the operators $H(\sigma)$ and $\Gamma(q)$ are unitarily equivalent. So the next assertion is a direct consequence of Theorem 4.2.

Theorem 4.5. *Let the function $\sigma(\mu)$ be defined by formula (4.4) where $q(j)$ are given by (4.3) and $\alpha > 0$. Then the asymptotic relation holds*

$$s_n(H(\sigma)) = v(\alpha)n^{-\alpha} + o(n^{-\alpha}), \quad n \rightarrow \infty,$$

where $v(\alpha)$ is given by (1.10).

4.3. Rotation of the symbol. For a parameter $\zeta \in \mathbb{T}$, let R_ζ be the ‘‘rotation by ζ ’’ operator:

$$(R_\zeta f)(\mu) = f(\mu/\zeta).$$

Obviously, R_ζ is a unitary operator in $L^2(\mathbb{T})$ and in $H^2(\mathbb{T})$. Similarly, let V_ζ be the multiplication by ζ^{-j} :

$$(V_\zeta u)(j) = \zeta^{-j} u(j).$$

Obviously, V_ζ is a unitary operator in $\ell^2(\mathbb{Z}_+)$.

Lemma 4.6. (i) *For arbitrary $\zeta \in \mathbb{T}$ and $\omega \in L^\infty(\mathbb{T})$, we have*

$$H(R_\zeta \omega) = R_\zeta H(\omega) R_\zeta.$$

In particular, if $H(\omega)$ is compact, then

$$s_n(H(R_\zeta \omega)) = s_n(H(\omega)), \quad \forall n \geq 1.$$

(ii) *For any sequence h such that $\Gamma(h)$ is bounded, we have*

$$\Gamma(V_\zeta h) = V_\zeta \Gamma(h) V_\zeta.$$

In particular, if $\Gamma(h)$ is compact, then

$$s_n(\Gamma(V_\zeta h)) = s_n(\Gamma(h)), \quad \forall n \geq 1.$$

Proof. Since

$$P_+ R_\zeta = R_\zeta P_+ \quad \text{and} \quad R_\zeta W R_\zeta = W,$$

the first assertion is a direct consequence of the definition (1.1) of the Hankel operator $H(\omega)$ in $H^2(\mathbb{T})$. The second assertion immediately follows from definition (3.1). \square

4.4. **Proof of Theorem 3.1.** Let the symbol $\sigma(\mu)$ be defined by relation (4.4) and let

$$\omega_{\natural}(\mu) = \sum_{\ell=1}^L \omega_{\ell}(\mu) \quad \text{where} \quad \omega_{\ell}(\mu) = b_{\ell} \sigma(\mu/\zeta_{\ell}). \quad (4.7)$$

According to Theorem 4.5 and Lemma 4.6(i) we have

$$s_n(H(\omega_{\ell})) = |b_{\ell}| v(\alpha) n^{-\alpha} + o(n^{-\alpha}), \quad n \rightarrow \infty.$$

It follows from Lemma 4.3 that $\omega_{\ell} \in L^{\infty}(\mathbb{T})$ and $\omega_{\ell} \in C^{\infty}(\mathbb{T} \setminus \zeta_{\ell})$. Since ζ_1, \dots, ζ_L are distinct points, the localisation principle (Theorem 1.1) is applicable to the sum (4.7). This yields

$$\lim_{n \rightarrow \infty} n s_n(H(\omega_{\natural}))^p = \sum_{\ell=1}^L \lim_{n \rightarrow \infty} n s_n(H(\omega_{\ell}))^p = v(\alpha)^p \sum_{\ell=1}^L |b_{\ell}|^p, \quad p = 1/\alpha. \quad (4.8)$$

Note that, by the definition (4.7),

$$\widehat{\omega}_{\ell}(j) = b_{\ell} \widehat{\sigma}(j) \zeta_{\ell}^{-j}$$

and hence according to formula (4.4)

$$\widehat{\omega}_{\natural}(j) = \sum_{\ell=1}^L b_{\ell} j^{-1} (\log j)^{-\alpha} \zeta_{\ell}^{-j} =: h_{\natural}(j), \quad j \geq 2.$$

Set $h_{\natural}(0) = h_{\natural}(1) = 0$. Since the operators $H(\omega_{\natural})$ and $\Gamma(h_{\natural})$ are unitarily equivalent, it follows from (4.8) that

$$\lim_{n \rightarrow \infty} n s_n(\Gamma(h_{\natural}))^p = v(\alpha)^p \sum_{\ell=1}^L |b_{\ell}|^p. \quad (4.9)$$

Next, we consider the error term

$$g(j) = \sum_{\ell=1}^L g_{\ell}(j) \zeta_{\ell}^{-j}$$

in (3.8). According to condition (3.9) it follows from Theorem 4.1 that $s_n(\Gamma(g_{\ell})) = o(n^{-\alpha})$ as $n \rightarrow \infty$. By Lemma 4.6(ii), we also have $s_n(\Gamma(V_{\zeta_{\ell}} g_{\ell})) = o(n^{-\alpha})$ and hence

$$s_n(\Gamma(g)) = o(n^{-\alpha}) \quad \text{as} \quad n \rightarrow \infty. \quad (4.10)$$

Since

$$\Gamma(h) = \Gamma(h_{\natural}) + \Gamma(g),$$

we can use Lemma 2.1 with $A = \Gamma(h_{\natural})$ and $B = \Gamma(g)$. The required relations (3.10), (3.11) follow from (4.9) and (4.10). \square

5. APPLICATIONS OF LOCALIZATION PRINCIPLE: CONTINUOUS CASE

5.1. **Hankel operators in $L^2(\mathbb{R}_+)$.** Integral Hankel operators $\mathbf{\Gamma}(\mathbf{h})$ in the space $L^2(\mathbb{R}_+)$ are defined by the relation

$$(\mathbf{\Gamma}(\mathbf{h})\mathbf{u})(t) = \int_0^\infty \mathbf{h}(t+s)\mathbf{u}(s)ds, \quad \mathbf{u} \in C_0^\infty(\mathbb{R}_+), \quad (5.1)$$

where at least $\mathbf{h} \in L_{\text{loc}}^1(\mathbb{R}_+)$; this function is called the *kernel* of the Hankel operator $\mathbf{\Gamma}(\mathbf{h})$. Under the assumptions on \mathbf{h} below the operators $\mathbf{\Gamma}(\mathbf{h})$ are compact.

Similarly to the discrete case, Hankel operators in the Hardy space $H_+^2(\mathbb{R})$ are unitarily equivalent to integral operators $\mathbf{\Gamma}(\mathbf{h})$ in the space $L^2(\mathbb{R}_+)$:

$$\Phi\mathbf{H}(\boldsymbol{\omega})\Phi^* = \mathbf{\Gamma}(\mathbf{h}) \quad \text{if} \quad \mathbf{h}(t) = \frac{1}{\sqrt{2\pi}}\widehat{\boldsymbol{\omega}}(t) \quad \text{for } t > 0. \quad (5.2)$$

The Fourier transform $\widehat{\boldsymbol{\omega}}$ of $\boldsymbol{\omega} \in L^\infty(\mathbb{R})$ should in general be understood in the sense of distributions (for example, on the Schwartz class $\mathcal{S}'(\mathbb{R})$) and the precise meaning of (5.2) is given by the equation

$$(\mathbf{H}(\boldsymbol{\omega})\Phi^*\mathbf{u}, \Phi^*\mathbf{u}) = (\mathbf{\Gamma}(\mathbf{h})\mathbf{u}, \mathbf{u}), \quad \mathbf{u} \in C_0^\infty(\mathbb{R}_+).$$

A function $\boldsymbol{\omega}(x)$ satisfying (5.2) is known as a symbol of the Hankel operator $\mathbf{\Gamma}(\mathbf{h})$.

5.2. **Main result in the continuous case.** In the discrete case, the spectral asymptotics of $\Gamma(h)$ is determined by the asymptotic behaviour of the sequence $h(j)$ as $j \rightarrow \infty$. In the continuous case, the behaviour of the kernel $\mathbf{h}(t)$ for $t \rightarrow \infty$ and for $t \rightarrow 0$ as well as the local singularities of $\mathbf{h}(t)$ at positive points t contribute to the spectral asymptotics of $\mathbf{\Gamma}(\mathbf{h})$. In the following result we exclude local singularities. We denote $\langle x \rangle = \sqrt{1 + |x|^2}$.

Theorem 5.1. *Let $\alpha > 0$, let $a_1, \dots, a_L \in \mathbb{R}$ be distinct numbers and let $\mathbf{b}_0, \mathbf{b}_1, \dots, \mathbf{b}_L \in \mathbb{C}$. Let the number $M = M(\alpha)$ be given by (3.5). Suppose that $\mathbf{h} \in L_{\text{loc}}^\infty(\mathbb{R}_+)$ if $\alpha < 1/2$ and $\mathbf{h} \in C^M(\mathbb{R}_+)$ if $\alpha \geq 1/2$. Assume that*

$$\mathbf{h}(t) = \sum_{\ell=1}^L (\mathbf{b}_\ell t^{-1}(\log t)^{-\alpha} + \mathbf{g}_\ell(t))e^{-ia_\ell t}, \quad t \geq 2, \quad (5.3)$$

$$\mathbf{h}(t) = \mathbf{b}_0 t^{-1}(\log(1/t))^{-\alpha} + \mathbf{g}_0(t), \quad t \leq 1/2, \quad (5.4)$$

where the error terms \mathbf{g}_ℓ satisfy the estimates

$$\mathbf{g}_\ell^{(m)}(t) = o(t^{-1-m}\langle \log t \rangle^{-\alpha}), \quad m = 0, \dots, M(\alpha), \quad (5.5)$$

as $t \rightarrow \infty$ for $\ell = 1, \dots, L$ and as $t \rightarrow 0$ for $\ell = 0$. Then the singular values of the integral Hankel operator $\mathbf{\Gamma}(\mathbf{h})$ in $L^2(\mathbb{R}_+)$ satisfy the asymptotic relation

$$s_n(\mathbf{\Gamma}(\mathbf{h})) = \mathbf{c} n^{-\alpha} + o(n^{-\alpha}), \quad n \rightarrow \infty, \quad (5.6)$$

where

$$\mathbf{c} = v(\alpha) \left(\sum_{\ell=0}^L |\mathbf{b}_\ell|^{1/\alpha} \right)^\alpha \quad (5.7)$$

and the coefficient $v(\alpha)$ is given by formula (1.10).

The proof in the continuous case follows the same general outline as in the discrete case with the only difference that the singularity of the kernel $\mathbf{h}(t)$ at $t = 0$ has to be treated separately. It corresponds to the singularity of the symbol $\omega(x)$ at infinity.

In Section 7 we consider kernels $\mathbf{h}(t)$ that have a singularity at some positive point and admit representation (5.3) for large t . It turns out that, similarly to Theorem 5.1, the contributions of the singularities of these two types to the asymptotics of singular values are independent of each other.

6. PROOF OF THEOREM 5.1

The proof of Theorem 5.1 follows the scheme of the proof of Theorem 3.1. The only new point is that now we have to additionally establish the correspondence between symbols singular at infinity and kernels singular at $t = 0$.

6.1. Singular value estimates and asymptotics. Let us state the analogues of Theorems 4.1 and 4.2.

Theorem 6.1. [10, Theorem 2.8] *Let $\alpha > 0$, and let the number $M = M(\alpha)$ be given by (3.5). Suppose that $\mathbf{g} \in L_{\text{loc}}^\infty(\mathbb{R}_+)$ if $\alpha < 1/2$ and $\mathbf{g} \in C^M(\mathbb{R}_+)$ if $\alpha \geq 1/2$. Assume that*

$$\mathbf{g}^{(m)}(t) = o(t^{-1-m} \langle \log t \rangle^{-\alpha}) \quad \text{as } t \rightarrow 0 \text{ and as } t \rightarrow \infty \quad (6.1)$$

for all $m = 0, 1, \dots, M$. Then

$$s_n(\mathbf{\Gamma}(\mathbf{g})) = o(n^{-\alpha}), \quad n \rightarrow \infty. \quad (6.2)$$

In [10] we also have a result with O instead of o in (6.1) and (6.2), although we will not need it in this paper. Observe that for $\alpha < 1/2$ we need only the estimate on \mathbf{g} , whereas for $\alpha \geq 1/2$ we also need estimates on the derivatives $\mathbf{g}^{(m)}$.

Next, we define model kernels $\mathbf{q}_0, \mathbf{q}_\infty$. Choose some non-negative functions $\chi_0, \chi_\infty \in C^\infty(\mathbb{R})$ such that

$$\chi_0(x) = \begin{cases} 1 & \text{for } |x| \leq c_1, \\ 0 & \text{for } |x| \geq c_2, \end{cases} \quad \chi_\infty(x) = \begin{cases} 0 & \text{for } |x| \leq C_1, \\ 1 & \text{for } |x| \geq C_2, \end{cases}$$

for some $0 < c_1 < c_2 < 1$ and $1 < C_1 < C_2$.

Theorem 6.2. [11, Theorem 1.2] *For $\alpha > 0$, set*

$$\mathbf{q}_0(t) = \chi_0(t)t^{-1}(\log(1/t))^{-\alpha}, \quad \mathbf{q}_\infty(t) = \chi_\infty(t)t^{-1}(\log t)^{-\alpha}, \quad t > 0. \quad (6.3)$$

Then

$$s_n(\mathbf{\Gamma}(\mathbf{q}_0)) = v(\alpha) n^{-\alpha} + o(n^{-\alpha}), \quad s_n(\mathbf{\Gamma}(\mathbf{q}_\infty)) = v(\alpha) n^{-\alpha} + o(n^{-\alpha}), \quad n \rightarrow \infty,$$

where $v(\alpha)$ given by (1.10).

Of course, this result corresponds to particular cases of Theorem 5.1 with $L = 1$, $\mathbf{b}_0 = 1$, $\mathbf{b}_1 = 0$ and $\mathbf{b}_0 = 0$, $\mathbf{b}_1 = 1$.

6.2. Model symbols. In order to put together the contributions of different terms in (5.3) and (5.4), we use the localization principle in the form of Theorem 2.7. To that end, we need to determine the singular supports of the symbols corresponding to the model kernels \mathbf{q}_0 , \mathbf{q}_∞ . Again, we will choose functions σ_0 , σ_1 whose Fourier transform coincides with the odd extension of \mathbf{q}_0 , \mathbf{q}_∞ to the real line. The proof below is very similar to that of Lemma 4.3.

Lemma 6.3. *Let σ_0 , σ_∞ be defined by*

$$\sigma_0(x) = 2i \int_0^\infty \mathbf{q}_0(t) \sin(xt) dt, \quad \sigma_\infty(x) = 2i \int_0^\infty \mathbf{q}_\infty(t) \sin(xt) dt, \quad x \in \mathbb{R}, \quad (6.4)$$

where $\mathbf{q}_0(t)$ and $\mathbf{q}_\infty(t)$ are given by (6.3) with $\alpha \geq 0$. Then $\sigma_0, \sigma_\infty \in L^\infty(\mathbb{R})$ and $\sigma_0 \in C^\infty(\mathbb{R})$, $\sigma_\infty \in C^\infty(\mathbb{R}_* \setminus \{0\})$.

Proof. Note that for all $x \in \mathbb{R}$, the first integral in (6.4) converges absolutely while the second one converges absolutely for $\alpha > 1$ and conditionally for $\alpha \leq 1$.

Since the integral in the definition (6.4) of σ_0 is taken over a finite interval, we can differentiate this integral with respect to x arbitrary many times. Hence $\sigma_0 \in C^\infty(\mathbb{R})$. To prove that $\sigma_\infty \in C^\infty(\mathbb{R}_* \setminus \{0\})$, we integrate by parts $2M + 2$ times in the definition (6.4):

$$\sigma_\infty(x) = 2i(-1)^{M+1} x^{-2M-2} \int_0^\infty \mathbf{q}_\infty^{(2M+2)}(t) \sin(xt) dt.$$

Since $\mathbf{q}_\infty^{(2M+2)}(t) = O(|t|^{-2M-3})$ as $|t| \rightarrow \infty$, we see that $\sigma_\infty \in C^m(\mathbb{R} \setminus \{0\})$ and $\sigma_\infty^{(m)}(x) \rightarrow 0$ for $m = 0, 1, \dots, 2M$ as $|x| \rightarrow \infty$. Finally, we use that M is arbitrary.

It remains to prove that the functions σ_0 and σ_∞ are bounded. Below $\kappa = 0$ or $\kappa = \infty$. We may suppose that $x > 0$. Write $\sigma_\kappa = \sigma_\kappa^{(1)} + \sigma_\kappa^{(2)}$, where

$$\sigma_\kappa^{(1)}(x) = 2i \int_0^{1/x} \mathbf{q}_\kappa(t) \sin(xt) dt, \quad \sigma_\kappa^{(2)}(x) = 2i \int_{1/x}^\infty \mathbf{q}_\kappa(t) \sin(xt) dt.$$

Since $|\sin(xt)| \leq xt$, for σ_1 we have the estimate

$$|\sigma_\kappa^{(1)}(x)| \leq 2x \int_0^{1/x} \mathbf{q}_\kappa(t) t dt \leq C$$

because $\mathbf{q}_\kappa(t)t$ are bounded functions. For $\sigma_\kappa^{(2)}$, integrating by parts once, we get

$$\sigma_\kappa^{(2)}(x) = -\frac{2i}{x} \int_{1/x}^\infty \mathbf{q}_\kappa(t) (\cos(xt))' dt = \frac{2i}{x} \mathbf{q}_\kappa(1/x) \cos 1 + \frac{2i}{x} \int_{1/x}^\infty \mathbf{q}'_\kappa(t) \cos(xt) dt.$$

The first term in the right-hand side is bounded because $\mathbf{q}_\kappa(t)t$ are bounded functions. The second term is also bounded because the functions $\mathbf{q}'_\kappa(t)t^2$ are bounded. \square

Remark 6.4. (i) It can be verified that the singular supports of σ_0 and σ_∞ are non-empty, that is, $\text{sing supp } \sigma_0 = \{\infty\}$ and $\text{sing supp } \sigma_\infty = \{0\}$. In fact, a more detailed calculation shows that the symbols σ_0, σ_∞ satisfy the asymptotics

$$\begin{aligned}\sigma_0(x) &= \pi i \text{sign } x |\log|x||^{-\alpha} (1 + o(1)), & x \rightarrow \infty, \\ \sigma_\infty(x) &= \pi i \text{sign } x |\log|x||^{-\alpha} (1 + o(1)), & x \rightarrow 0.\end{aligned}$$

(ii) For some values of α , instead of the odd extension of $\mathbf{q}_0(t)$ and $\mathbf{q}_\infty(t)$ to the negative t , one can extend them by zero, i.e. one can choose

$$\begin{aligned}\tilde{\sigma}_0(x) &= \int_0^\infty \mathbf{q}_0(t) e^{ixt} dt, & \text{if } \alpha < 1, \\ \tilde{\sigma}_\infty(x) &= \int_0^\infty \mathbf{q}_\infty(t) e^{ixt} dt, & \text{if } \alpha > 1,\end{aligned}$$

instead of $\sigma_0(x), \sigma_\infty(x)$, respectively.

It easily follows from Lemma 6.3 that

$$\Phi \mathbf{H}(\sigma_0) \Phi^* = \Gamma(\mathbf{q}_0) \quad \text{and} \quad \Phi \mathbf{H}(\sigma_\infty) \Phi^* = \Gamma(\mathbf{q}_\infty). \quad (6.5)$$

Indeed, in view of relation (5.2), we only have to check that

$$\frac{1}{\sqrt{2\pi}} \widehat{\sigma}_0(t) = \mathbf{q}_0(t), \quad \frac{1}{\sqrt{2\pi}} \widehat{\sigma}_\infty(t) = \mathbf{q}_\infty(t), \quad t > 0, \quad (6.6)$$

where the Fourier transform is understood, for example, in the class of distributions $\mathcal{S}(\mathbb{R})'$. According to the first formula in (6.4), the function $(2\pi)^{-1/2} \sigma_0$ is the Fourier transform of the ‘‘extended’’ distribution $\mathbf{q}_0^{(\text{ext})}$ defined by the equation

$$\langle \mathbf{q}_0^{(\text{ext})}, \varphi \rangle = \int_0^\infty \mathbf{q}_0(t) (\bar{\varphi}(t) - \bar{\varphi}(-t)) dt.$$

Thus $(2\pi)^{-1/2} \widehat{\sigma}_0(t) = \mathbf{q}_0^{(\text{ext})}(t)$ which coincides with $\mathbf{q}_0(t)$ for $t > 0$. The second equality (6.6) is obvious because $\mathbf{q}_\infty \in \mathcal{S}(\mathbb{R})'$.

The next assertion is a direct consequence of relations (6.5) and Theorem 6.2.

Theorem 6.5. *Let the functions σ_0 and σ_∞ be defined by formulas (6.4) where $\mathbf{q}_0(t)$ and $\mathbf{q}_\infty(t)$ are given by (6.3) and $\alpha > 0$. Then the asymptotic relations hold*

$$s_n(\mathbf{H}(\sigma_0)) = v(\alpha) n^{-\alpha} + o(n^{-\alpha}), \quad s_n(\mathbf{H}(\sigma_\infty)) = v(\alpha) n^{-\alpha} + o(n^{-\alpha}), \quad n \rightarrow \infty,$$

where $v(\alpha)$ is given by (1.10).

6.3. **Shifts of symbols.** For a parameter $a \in \mathbb{R}$, let \mathbf{R}_a be the shift

$$(\mathbf{R}_a \mathbf{f})(x) = \mathbf{f}(x - a).$$

Obviously, \mathbf{R}_a is a unitary operator in $L^2(\mathbb{R})$ and $H_+^2(\mathbb{R})$. Of course, now \mathbf{R}_a is not a rotation, but we keep the letter \mathbf{R} in order to maintain the analogy between the discrete and continuous cases.

Similarly, let \mathbf{V}_a be the multiplication operator

$$(\mathbf{V}_a \mathbf{u})(t) = e^{-iat} \mathbf{u}(t), \quad t > 0.$$

Obviously, \mathbf{V}_a is a unitary operator in $L^2(\mathbb{R}_+)$.

Recall that the Hankel operators $\mathbf{H}(\boldsymbol{\omega})$ in $H^2(\mathbb{R})$ were defined by formula (2.18).

Lemma 6.6. *For arbitrary $a \in \mathbb{R}$, we have the following statements:*

(i) *For any $\boldsymbol{\omega} \in L^\infty(\mathbb{R})$, we have*

$$\mathbf{H}(\mathbf{R}_a \boldsymbol{\omega}) = \mathbf{R}_a \mathbf{H}(\boldsymbol{\omega}) \mathbf{R}_a.$$

In particular, if $\mathbf{H}(\boldsymbol{\omega})$ is compact, then

$$s_n(\mathbf{H}(\mathbf{R}_a \boldsymbol{\omega})) = s_n(\mathbf{H}(\boldsymbol{\omega})), \quad \forall n \geq 1.$$

(ii) *Suppose that $\Gamma(\mathbf{h})$ is bounded; then*

$$\Gamma(\mathbf{V}_a \mathbf{h}) = \mathbf{V}_a \Gamma(\mathbf{h}) \mathbf{V}_a.$$

In particular, if $\Gamma(\mathbf{h})$ is compact, then

$$s_n(\Gamma(\mathbf{V}_a \mathbf{h})) = s_n(\Gamma(\mathbf{h})), \quad \forall n \geq 1.$$

Proof. Since

$$\mathbf{P}_+ \mathbf{R}_a = \mathbf{R}_a \mathbf{P}_+ \quad \text{and} \quad \mathbf{R}_a \mathbf{W} \mathbf{R}_a = \mathbf{W},$$

the first assertion is a direct consequence of the definition (2.18) of the Hankel operator $\mathbf{H}(\boldsymbol{\omega})$ in $H^2(\mathbb{R})$. The second assertion immediately follows from the definition (5.1). \square

6.4. **Proof of Theorem 5.1.** Let the symbols $\boldsymbol{\sigma}_0(x)$ and $\boldsymbol{\sigma}_\infty(x)$ be defined by relations (6.4) and let

$$\boldsymbol{\omega}_\natural(x) = \boldsymbol{\omega}_0(x) + \sum_{\ell=1}^L \boldsymbol{\omega}_\ell(x) \quad \text{where} \quad \boldsymbol{\omega}_0(x) = \mathbf{b}_0 \boldsymbol{\sigma}_0(x), \quad \boldsymbol{\omega}_\ell(x) = \mathbf{b}_\ell \boldsymbol{\sigma}_\infty(x - a_\ell). \quad (6.7)$$

According to Theorem 6.5 and Lemma 6.6(i) we have

$$s_n(\mathbf{H}(\boldsymbol{\omega}_\ell)) = |\mathbf{b}_\ell| v(\alpha) n^{-\alpha} + o(n^{-\alpha}), \quad n \rightarrow \infty,$$

for all $\ell = 0, 1, \dots, L$. It follows from Lemma 6.3 that $\boldsymbol{\omega}_\ell \in L^\infty(\mathbb{R})$ for all $\ell = 0, 1, \dots, L$, $\boldsymbol{\omega}_0 \in C^\infty(\mathbb{R})$ and $\boldsymbol{\omega}_\ell \in C^\infty(\mathbb{R}_* \setminus a_\ell)$ for $\ell = 1, \dots, L$. Since a_1, \dots, a_L

are distinct points, the localisation principle (Theorem 2.7) is applicable to the sum (6.7). This yields

$$\lim_{n \rightarrow \infty} ns_n(H(\boldsymbol{\omega}_\natural))^p = \sum_{\ell=0}^L \lim_{n \rightarrow \infty} ns_n(H(\boldsymbol{\omega}_\ell))^p = v(\alpha)^p \sum_{\ell=0}^L |\mathbf{b}_\ell|^p, \quad p = 1/\alpha. \quad (6.8)$$

Note that, by its definition (6.7), $\widehat{\boldsymbol{\omega}}_0(t) = \mathbf{b}_0 \widehat{\boldsymbol{\sigma}}_0(t)$,

$$\widehat{\boldsymbol{\omega}}_\ell(t) = \mathbf{b}_\ell \widehat{\boldsymbol{\sigma}}_\infty(t) e^{-ia_\ell t}, \quad \ell = 1, \dots, L,$$

and hence according to formula (6.6)

$$\widehat{\boldsymbol{\omega}}_\natural(t) = \mathbf{b}_0 \chi_0(t) t^{-1} |\log t|^{-\alpha} + \sum_{\ell=1}^L \mathbf{b}_\ell \chi_\infty(t) t^{-1} |\log t|^{-\alpha} e^{-ia_\ell t} =: \mathbf{h}_\natural(t), \quad t > 0.$$

In view of relation (5.2) it now follows from (6.8) that

$$\lim_{n \rightarrow \infty} ns_n(\boldsymbol{\Gamma}(\mathbf{h}_\natural))^p = v(\alpha)^p \sum_{\ell=0}^L |\mathbf{b}_\ell|^p.$$

Next, we consider the error term

$$\mathbf{g}(t) = \mathbf{h}(t) - \mathbf{h}_\natural(t) = \mathbf{g}_0(t) + \sum_{\ell=1}^L \mathbf{g}_\ell(t) e^{-ia_\ell t}$$

where all functions $\mathbf{g}_\ell(t)$, $\ell = 0, 1, \dots, L$, satisfy the condition (5.5) both for $t \rightarrow 0$ and $t \rightarrow \infty$. It follows from Theorem 6.1 and Lemma 6.6(ii) that $s_n(\mathbf{H}(\mathbf{g}_\ell)) = o(n^{-\alpha})$ and hence

$$s_n(\mathbf{H}(\mathbf{g})) = o(n^{-\alpha}) \quad \text{as } n \rightarrow \infty. \quad (6.9)$$

Since

$$\mathbf{H}(h) = \mathbf{H}(\mathbf{h}_\natural) + \mathbf{H}(\mathbf{g}),$$

we can use Lemma 2.1 with $A = \mathbf{H}(\mathbf{h}_\natural)$ and $B = \mathbf{H}(\mathbf{g})$. The required relations (5.6), (5.7) follow from (6.8) and (6.9). \square

7. LOCAL SINGULARITIES OF THE KERNEL

The localization principle shows that the results on the asymptotics of singular values of different Hankel operators can be combined provided that the singular supports of their symbols are disjoint. This idea has already been illustrated by Theorems 3.1 and 5.1. Here we apply the same arguments to kernels $\mathbf{h}(t)$ satisfying condition (5.3) as $t \rightarrow \infty$ and singular at some point $t_0 > 0$. Below $\mathbb{1}_+(t)$ is the characteristic function of \mathbb{R}_+ .

The effect of local singularities of $\mathbf{h}(t)$ on the asymptotics of singular values of the corresponding Hankel operator $\boldsymbol{\Gamma}(\mathbf{h})$ was studied in [3] and later in [13]. The methods of these papers are quite different. We need the following result obtained in [13].

Lemma 7.1. *Let $t_0 > 0$, $m \in \mathbb{Z}_+$ and*

$$\mathbf{a}_m(t) = (t_0 - t)^m \mathbb{1}_+(t_0 - t). \quad (7.1)$$

Then $\text{Ker } \Gamma(\mathbf{a}_m) = L^2(t_0, \infty)$ and

$$\Gamma(\mathbf{a}_m)|_{L^2(0, t_0)} = m! A_m^{-1}$$

where the self-adjoint operator A_m in $L^2(0, t_0)$ is defined by the differential expression

$$(A_m \mathbf{u})(t) = (-1)^{m+1} \mathbf{u}^{(m+1)}(t_0 - t)$$

and the boundary conditions

$$\mathbf{u}(t_0) = \dots = \mathbf{u}^{(m)}(t_0) = 0. \quad (7.2)$$

Note that the operator A_m^2 is given by the differential expression

$$(A_m^2 \mathbf{u})(t) = (-1)^{m+1} \mathbf{u}^{(2m+2)}(t)$$

and the boundary conditions (7.2) and

$$\mathbf{u}^{(m+1)}(0) = \dots = \mathbf{u}^{(2m+1)}(0) = 0.$$

Thus A_m^2 is a regular differential operator and the asymptotics of its eigenvalues is given by the Weyl formula. Therefore the following result is an immediate consequence of Lemma 7.1.

Corollary 7.2. *Let the function $\mathbf{a}_m(t)$ be given by formula (7.1). Then*

$$s_n(\Gamma(\mathbf{a}_m)) = m! t_0^{m+1} (\pi n)^{-m-1} (1 + O(n^{-1})), \quad n \rightarrow \infty. \quad (7.3)$$

Notice that formula (7.3) was obtained much earlier in [3] by a completely different method.

We also note the explicit formula for the symbol $\tau_m(x)$ of the operator $\Gamma(\mathbf{a}_m)$:

$$\tau_m(x) = m! (ix)^{-m-1} \left(e^{it_0 x} - \sum_{k=0}^m \frac{1}{k!} (it_0 x)^k \right), \quad x \in \mathbb{R}. \quad (7.4)$$

Obviously, $\tau_m \in C^\infty(\mathbb{R})$ and $\tau_m(x)$ is an oscillating function as $|x| \rightarrow \infty$.

We are now in a position to consider the general case.

Theorem 7.3. *Let $t_0 > 0$, $m \in \mathbb{Z}_+$ and $\beta \in \mathbb{C}$. Set*

$$\mathbf{h}_m(t) = \beta (t_0 - t)^m \mathbb{1}_+(t_0 - t) + \mathbf{h}(t)$$

where $\mathbf{h}(t)$ satisfies the assumptions of Theorem 5.1 with $\mathbf{b}_0 = 0$ and $\alpha = m + 1$. Then the singular values of the operator $\Gamma(\mathbf{h}_m)$ satisfy the asymptotic relation

$$s_n(\Gamma(\mathbf{h}_m)) = \mathbf{c}_m n^{-m-1} + o(n^{-m-1})$$

with

$$\mathbf{c}_m = \left(\pi^{-1} t_0 (m! |\beta|)^{1/\alpha} + v(\alpha)^{1/\alpha} \sum_{\ell=1}^L |b_\ell|^{1/\alpha} \right)^\alpha, \quad \alpha = m + 1,$$

and $v(\alpha)$ defined by (1.10).

Proof. It is almost the same as that of Theorem 5.1. Let us use notation (7.1). The asymptotics of the singular values of the operator $\mathbf{\Gamma}(\mathbf{a}_m)$ is given by formula (7.3). The operator $\mathbf{\Gamma}(\mathbf{h})$ satisfies the assumptions of Theorem 5.1 so that the asymptotics of its eigenvalues is given by formula (5.6). The symbol (7.4) of the operator $\mathbf{\Gamma}(\mathbf{a}_m)$ is singular only at infinity. Neglecting the terms satisfying the assumptions of Theorem 6.1 and using Lemma 6.3, we see that the singular support of the symbol of the operator $\mathbf{\Gamma}(\mathbf{h})$ consists of the points $a_1, \dots, a_L \in \mathbb{R}$. Therefore applying Theorem 2.7, we conclude the proof. \square

Observe that we have excluded the term (5.4) singular at $t = 0$ in Theorem 7.3 because the corresponding symbol is singular at the same point $x = \infty$ as the function (7.4). In this case one might expect that the contributions of singularities of $\mathbf{h}(t)$ at $t = 0$ and $t = t_0 > 0$ are not independent of each other. In any case, our technique does not allow us to treat this situation.

Let us discuss the operator $\mathbf{\Gamma}(\mathbf{a}_m)$ in the representation $\ell^2(\mathbb{Z}_+)$, that is, the operator

$$\mathcal{F}\mathcal{U}^*\mathbf{H}(\boldsymbol{\tau}_m)\mathcal{U}\mathcal{F}^* = \Gamma(g_m).$$

Here $g_m(j)$ are the Fourier coefficients of the function $\tau_m(\mu)$ linked to $\boldsymbol{\tau}_m(x)$ by formula (2.20). Making the change of variables (2.19) in (7.4), we see that $\tau_m(\mu)$ is an oscillating function as $\mu \rightarrow 1$. Therefore the asymptotics of its Fourier coefficients $g_m(j)$ is determined by the stationary phase method which yields:

$$g_m(j) \sim m!\pi^{-1/2}2^{-(2m+1)/4}j^{-(2m+5)/4} \cos(2\sqrt{2j} - \pi(2m+1)/4).$$

Note that these sequences decay faster as $j \rightarrow \infty$ than the coefficients (3.8) where $\alpha = m + 1$. Nevertheless due to the oscillating factor their contribution to the asymptotics of singular values is of the same order.

ACKNOWLEDGEMENTS

The authors are grateful to the Departments of Mathematics of King's College London and of the University of Rennes 1 (France) for the financial support. The second author (D.Y.) acknowledges also the support and hospitality of the Isaak Newton Institute for Mathematical Sciences (Cambridge University, UK) where a part of this work has been done during the program Periodic and Ergodic Spectral Problems.

REFERENCES

- [1] M. SH. BIRMAN, M. Z. SOLOMYAK, *Spectral theory of selfadjoint operators in Hilbert space*. D. Reidel, Dordrecht, 1987.
- [2] M. SH. BIRMAN, M. Z. SOLOMYAK, *Compact operators with power asymptotic behavior of the singular numbers*. J. Sov. Math. **27** (1984), 2442–2447.

- [3] K. GLOVER, J. LAM, J. R. PARTINGTON, *Rational approximation of a class of infinite-dimensional systems I: singular values of Hankel operators*, Math. Control Signals Systems (1990) **3**, 325–344.
- [4] J. S. HOWLAND, *Spectral theory of self-adjoint Hankel matrices*, Michigan Math. J. **33** (1986), 145–153.
- [5] R. S. ISMAGILOV, *On the spectrum of Toeplitz matrices*, Sov. Math. Dokl. **4** (1963), 462–465.
- [6] N. K. NIKOLSKI, *Operators, functions, and systems: an easy reading*, vol. I: Hardy, Hankel, and Toeplitz, Math. Surveys and Monographs vol. 92, Amer. Math. Soc., Providence, Rhode Island, 2002.
- [7] V. PELLER, *Hankel operators and their applications*, Springer, 2003.
- [8] S. R. POWER, *Hankel operators with discontinuous symbols*, Proc. Amer. Math. Soc. **65** 1977, 77–79.
- [9] A. PUSHNITSKI, D. YAFAEV, *Spectral theory of piecewise continuous functions of self-adjoint operators*, Proc. London Math. Soc. **108** (2014), 1079–1115.
- [10] A. PUSHNITSKI, D. YAFAEV, *Sharp estimates for singular values of Hankel operators*, to appear in Integral Equations and Operator Theory, doi: 10.1007/s00020-015-2239-0.
- [11] A. PUSHNITSKI, D. YAFAEV, *Asymptotic behavior of eigenvalues of Hankel operators*, to appear in Int. Math. Res. Notices, doi: 10.1093/imrn/rnv048.
- [12] A. PUSHNITSKI, D. YAFAEV, *Rational approximation of functions with logarithmic singularities*, in preparation.
- [13] D. R. YAFAEV, *Criteria for Hankel operators to be sign-definite*, Analysis & PDE **8** (2015), no. 1, 183–221.

DEPARTMENT OF MATHEMATICS, KING'S COLLEGE LONDON, STRAND, LONDON, WC2R 2LS, U.K.

E-mail address: alexander.pushnitski@kcl.ac.uk

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF RENNES-1, CAMPUS BEAULIEU, 35042, RENNES, FRANCE

E-mail address: yafaev@univ-rennes1.fr