

DIAGONALISING AN ULTRAFILTER AND PRESERVING A P -POINT

HEIKE MILDENBERGER

ABSTRACT. With Ramsey-theoretic methods we show: It is consistent that there is a forcing that diagonalises one ultrafilter over ω and preserves another ultrafilter.

1. INTRODUCTION

Suppose M is a family of infinite sets such that M is a family that is large in some sense, e.g. unbounded or non-meagre, and that \mathbb{P} is a notion of forcing. We can ask whether M or some closure of M or reinterpretation of M in any \mathbb{P} -generic extension $V^{\mathbb{P}}$ are still large in $V^{\mathbb{P}}$. If this is the case we say that \mathbb{P} preserves that M is large. Various degrees of preservation can be distinguished: the forcing \mathbb{P} preserves the largeness of one particular set M , e.g. M being the set of all ground model reals, or \mathbb{P} preserves some large sets and makes others non-large, or \mathbb{P} preserves any large set of a certain form.

We are concerned with partial preservation for the largeness notion of generating an ultrafilter over ω , the set of natural numbers. Suppose that M is a family of infinite subsets of ω , the set of natural numbers. We say M generates an ultrafilter if $\{Y : (\exists X \in M)(X \subseteq Y)\}$ is an ultrafilter. For the case of unboundedness and Mathias forcing with a filter, examples of preserving one unbounded family and examples preserving all unbounded families are given [5]. Examples for partial preservation of the non-meagerness and non-nullness are given in [8]. Forcings with Milliken–Taylor ultrafilters preserving a P -point while destroying all P -points that are superfilters of a certain filter are given in [6] and in [11]. It is not known whether these forcings diagonalise an ultrafilter from the ground model. Here we add a new kind of example: Preserving one P -point and diagonalising another.

In Section 3 we introduce suitable families \mathcal{H} (see Def. 3.3) in a combinatorial space $(\mathcal{P})^\omega$ (see Def. 2.2(3)) and the notion “ \mathcal{H} avoids \mathcal{V} ” for a P -point \mathcal{V} . We prove a more general preservation theorem for a given P -point and a suitable family (Theorem 3.17). In the special case of a suitable family \mathcal{H} that projects to an ultrafilter over ω the theorem has the following form:

Date: March 11, 2016.

2010 Mathematics Subject Classification. 03E05, 03E35, 05C55.

Key words and phrases. Iterated proper forcing, normed creatures, P -points, preservation theorems.

The author would like to thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for support and hospitality during the programme Mathematical, Foundational and Computational Aspects of the Higher Infinite where work on this paper was undertaken.

Theorem 1.1. *Suppose that \mathcal{V} is a P -point and \mathcal{U} is an ultrafilter over ω such that there is a suitable family \mathcal{H} with projection $\Phi_2(\mathcal{H}) = \mathcal{U}$ and that $\Phi_2(\mathcal{H}) \not\leq_{RB} \mathcal{V}$. Then there is a proper notion of forcing that diagonalises \mathcal{U} and preserves \mathcal{V} .*

The premises to the theorem are consistent with ZFC:

Theorem 1.2. *Under CH or Martin's Axiom, given any P -point \mathcal{V} there is a suitable maximal centred family \mathcal{C} with projection $\Phi_2(\mathcal{C}) = \mathcal{U}$ such that \mathcal{U} is an ultrafilter and $\mathcal{U} \not\leq_{RB} \mathcal{V}$.*

Remark 1.3. In the first theorem we just use that \mathcal{H} is suitable and $\Phi_2(\mathcal{H})$ is centred. This might increase the versatility of suitable families. However, we do not know how to construct a family \mathcal{H} as in the first theorem without centredness. Moreover maximality can be added along the construction.

If any two ultrafilters over ω are nearly coherent (see definitions below) then there are no examples \mathcal{C}, \mathcal{V} with the stated properties.

We will prove the existence theorem in Section 4.

We recall definitions and facts: For a set X , we denote its powerset by $\mathcal{P}(X)$. By a *filter over ω* we mean a non-empty subset of $\mathcal{P}(\omega)$ that is closed under supersets and under finite intersections and that does not contain the empty set. We call a filter *non-principal* if it contains all cofinite subsets of ω and we call it an ultrafilter if it is a maximal filter.

For $B \subseteq \omega$ and $f: \omega \rightarrow \omega$, we let $f[B] = \{f(b) : b \in B\}$ and $f^{-1}[B] = \{n : f(n) \in B\}$. For $\mathcal{B} \subseteq \mathcal{P}(\omega)$ we let $f(\mathcal{B}) = \{X : f^{-1}[X] \in \mathcal{B}\}$. This double lifting is an important function from $\mathcal{P}(\mathcal{P}(\omega))$ into itself. In analysis the special case of f being finite-on-one (that means that the preimage of each natural number is finite) is particularly useful, see e.g., [3].

Let \mathcal{F} be a non-principal filter over ω and let $f: \omega \rightarrow \omega$ be finite-to-one. Then also $f(\mathcal{F})$ is a non-principal filter. It is the filter generated by $\{f[X] : X \in \mathcal{F}\}$. From now on we consider only non-principal filters and ultrafilters. Two filters \mathcal{F} and \mathcal{G} are *nearly coherent*, if there is some finite-to-one $f: \omega \rightarrow \omega$ such that $f(\mathcal{F}) \cup f(\mathcal{G})$ generates a filter. On the set of non-principal ultrafilters near coherence is an equivalence relation whose equivalence classes are called near-coherence classes. The principle near coherence of filters (short NCF) says that any two non-principal ultrafilters over ω are nearly coherent. Blass and Shelah [4] showed that NCF is consistent relative to ZFC.

The set of infinite subsets of ω is denoted by $[\omega]^\omega$, the set of finite subsets of ω is denoted by $[\omega]^{<\omega}$. We say “ A is almost a subset of B ” and write $A \subseteq^* B$ iff $A \setminus B$ is finite. Similarly, the symbol $=^*$ denotes equality up to finitely many exceptions in $[\omega]^\omega$ or in ${}^\omega\omega$, the set of functions from ω to ω . For $X \subseteq \omega$, we write X^c for $\omega \setminus X$.

Definition 1.4. Let κ be a regular uncountable cardinal. An ultrafilter \mathcal{U} is called a P_κ -point if for every $\gamma < \kappa$, for every $A_i \in \mathcal{U}$, $i < \gamma$, there is some $A \in \mathcal{U}$ such that for all $i < \gamma$, $A \subseteq^* A_i$; such an A is called a *pseudo-intersection* of the A_i , $i < \gamma$. A P_{\aleph_1} -point is called a P -point.

Let \mathbb{P} be a notion of forcing. We say that \mathbb{P} *preserves* an ultrafilter \mathcal{U} over I if

$$\Vdash_{\mathbb{P}} “(\forall X \subseteq I)(\exists Y \in \mathcal{U})(Y \subseteq X \vee Y \subseteq I \setminus X)”$$

and in the contrary case we say “ \mathbb{P} *destroys* \mathcal{U} ”. A particular way to destroy a non-principal ultrafilter is to diagonalise it, that means adding an infinite set X such that for any $Y \in \mathcal{U}$, $X \subseteq^* Y$.

If \mathbb{P} preserves \mathcal{U} then \mathcal{U} generates an ultrafilter in $\mathbf{V}[G]$. If \mathbb{P} is proper and preserves \mathcal{U} as an ultrafilter and \mathcal{U} is a P -point, then \mathcal{U} generates a P -point in the extension, since any countable set of ground model sets in the extension has a countable superset in the ground model, see [4, Lemma 3.2].

By nowadays, techniques for preserving ultrafilters that do not involve preservation of P -points are much more difficult than the known proofs of P -point preservation, see e.g. [12]. This experience is, at least partially, based on mathematical reasons: Any forcing that adds a real destroys an ultrafilter [1, Theorem 3.5], whereas Miller forcing, Sacks forcing and a few other tree forcings preserve any P -point.

Let \mathcal{V} be an ultrafilter over ω . Diagonalising $f(\mathcal{V})$ means destroying \mathcal{V} : If $(\forall A \in f(\mathcal{V})(X \subseteq^* A)$, then $f^{-1}[X] \notin \mathcal{V}$ and $(f^{-1}[X])^c \notin \mathcal{V}$. The function f need not be finite-to-one for this. Hence under NCF there are no \mathcal{C} and \mathcal{V} as in the conditions of Theorem 1.1. This shows that the premises to Theorem 1.1 are independent of ZFC.

The paper is organised as follows: In Section 2 we explain normed subsets of powersets, introduce $(\mathcal{P})^\omega$ and introduce suitable sets. In Section 3 we explain Blass–Shelah forcing and Blass–Shelah with a suitable set $\mathcal{H} \subseteq (\mathcal{P})^\omega$. We recall the definition of the Rudin–Blass order \leq_{RB} and we recall Eisworth’s work on the preservation of P -points for Matet forcing and prove Theorem 1.1. In Section 4 we prove Theorem 1.2. We close with a short discussion and some open questions.

In the forcing, the stronger condition is the *smaller* one. This direction fits to the \leq -relation on the sequences of possibilities (see Def. 2.2(4)) that form the second components of conditions. In addition we follow the alphabetical rule: Later letters are used for stronger conditions.

2. SETS OF NORMED SUBSETS OF POWERSETS

In this section we introduce a relative of Blass–Shelah forcing ([4]). Conditions in either version of Blass–Shelah forcing are of the form $p = (s, \bar{a})$. The component s is a finite subset of ω , which is usually called the trunk, and the component \bar{a} is an ω -sequence of hereditarily finite sets and is usually called the pure part.

Now we define a space of \mathcal{P} from which the entries of the pure parts of our posets will be taken.

Definition 2.1. (1) A finite subset s of ω is called a block. A *set of possibilities* is a subset of the power set of a block. We denote by \mathcal{P} the set of all sets of possibilities. Typically we use variables s, t, \dots for blocks and $a, b, c \dots$ for sets of possibilities.

So sets of possibilities are one powerset operation higher than blocks.

- (2) Let a be a set of possibilities and $Y \subseteq \omega$. We let $a \upharpoonright Y = \{s : s \in a, s \subseteq Y\}$.
- (3) We define $\circ : \mathcal{P} \times \mathcal{P} \rightarrow \mathcal{P}$ by $a \circ b = \{s \cup t : s \in a, t \in b\}$.
- (4) We define $\text{norm} : \mathcal{P} \rightarrow \omega$ as follows:
- (a) $\text{norm}(a) \geq 0$, always,
 - (b) $\text{norm}(a) \geq 1$ iff $\bigcup a \neq \emptyset$,
 - (c) $\text{norm}(a) \geq k + 1$ iff whenever $\bigcup a = Y_1 \cup Y_2$ then $\max(\text{norm}(a \upharpoonright Y_1), \text{norm}(a \upharpoonright Y_2)) \geq k$,
 - (d) $\text{norm}(a) = k$ iff ($\text{norm}(a) \geq k$ and $\text{norm}(a) \not\geq k + 1$).

The structure (\mathcal{P}, \circ) is a semigroup.

If $\text{norm}(a) \geq 1$, then a contains a non-empty set.

- Definition 2.2.** (1) For $a, b \in \mathcal{P}$ we write $a < b$ iff $\bigcup a, \bigcup b \neq \emptyset$ and $(\forall n \in \bigcup a)(\forall m \in \bigcup b)(n < m)$.
- (2) A sequence $\bar{a} = \langle a_n : n \in \omega \rangle$ of members of \mathcal{P} is called unmeshed if for all n , $a_n < a_{n+1}$.
- (3) By $(\mathcal{P})^\omega$ we denote the set of unmeshed sequences \bar{a} such that $\lim_{n \rightarrow \omega} \text{norm}(a_n) = \omega$.
- (4) For sequences $\bar{a}, \bar{b} \in (\mathcal{P})^\omega$ we write $\bar{b} \leq \bar{a}$ or “ \bar{b} is stronger than \bar{a} ” iff there are a strictly increasing sequence $\langle i_n : n \in \omega \rangle$ and a strictly increasing function $g \in {}^\omega \omega$ such that for any n ,

$$b_n \subseteq a_{i_{g(n)}} \circ a_{i_{g(n)+1}} \circ \cdots \circ a_{i_{g(n+1)}-1}.$$

So this means first $\langle a_n : n \in \omega \rangle$ is thinned out to $\langle a_{i_n} : n \in \omega \rangle$ and then finite intervals of members of the subsequence are merged and then a subset is taken.

- (5) For sequences $\bar{a}, \bar{b} \in (\mathcal{P})^\omega$ we write $\bar{a} \leq^* \bar{b}$ iff there is an n such that $\langle a_k : k \geq n \rangle \leq \bar{b}$.
- (6) For sequences $\bar{a}, \bar{b} \in (\mathcal{P})^\omega$ we write $\bar{a} \perp^* \bar{b}$ if they are incompatible in \leq^* , i.e., if there is no $\bar{c} \in (\mathcal{P})^\omega$ such that $\bar{c} \leq^* \bar{a}, \bar{b}$.

So for any sequence $\bar{a} \in (\mathcal{P})^\omega$ we have for any n , $\text{norm}(a_n) \geq 1$ and $\max(\bigcup a_n) \geq n$.

Lemma 2.3. *The relations \leq and \leq^* are transitive.*

The next two notions connect elements of $(\mathcal{P})^\omega$ with subsets of ω .

Definition 2.4. (1) For $\bar{a} \in (\mathcal{P})^\omega$ we let $\text{set}(\bar{a}) = \bigcup \{\bigcup a_n : n \in \omega\}$.

(2) Let $\mathcal{H} \subseteq (\mathcal{P})^\omega$. The *projection of \mathcal{H} into $[\omega]^\omega$* is $\Phi_2(\mathcal{H}) = \{\text{set}(\bar{a}) : \bar{a} \in \mathcal{H}\}$.¹

Definition 2.5. Let $\bar{a} \in (\mathcal{P})^\omega$, $n \in \omega$. We write $(\bar{a} \text{ past } n)$ for $\langle a_i : i \in [k, \omega) \rangle$, where k is the minimal number such that $n \leq \min \bigcup a_k$.

Now we introduce a version \mathbb{Q}^{242} of Blass-Shelah forcing (see [4]).

¹We write Φ_2 to distinguish it from the projection that is used for Matet forcing in [6].

Definition 2.6. In the forcing order \mathbb{Q}^{242} , conditions are pairs (s, \bar{a}) such that $s \in [\omega]^{<\omega}$ and $\bar{a} \in (\mathcal{P})^\omega$ and $(\forall n \in s)(\forall m \in \bigcup a_0)(n < m)$. We let $(t, \bar{b}) \leq (s, \bar{a})$ (recall the stronger condition is the smaller one) iff $s \subseteq t$ and there are $k \in \omega$ and $i_0 < i_1 \cdots < i_{k-1}$ such that $t \setminus s \in a_{i_0} \circ \cdots \circ a_{i_{k-1}}$ and $\bar{b} \leq (\bar{a} \text{ past } \max(a_{i_{k-1}}) + 1)$, where \leq is from Def. 2.2(4).

For readers of [4] we give a brief description of the differences: Instead of sequences of relations $(r_k)_k$, $r_k \in K_{n_k, m_k} \subseteq \mathcal{P}(\mathcal{P}(n_k) \times \mathcal{P}(m_k))$, $n_k < m_k \leq n_{k+1}$, we work with $\bar{a} \in (\mathcal{P})^\omega$. Our version is forgetful, this means that $(s, t) \in r \in K_{n, m}$ from [4] is replaced by $s \in a_i \wedge (t \setminus s \in a_j)$ for some $i < j$ with $\max(\bigcup a_i) < n$ and $n \leq \min(\bigcup a_j) \leq \max(\bigcup a_j) < m$. Hence the set of pure parts of q 's that are stronger than p depends only on the maximum of the trunk and on the pure part of p and not on the whole p .

Note: The trunks contain no information about the block structure. Moreover, there are gaps: if $(t, \bar{b}) \leq (s, \bar{a})$, and $t \setminus s \cap \bigcup a_n \neq \emptyset$, then \bar{b} must begin after the maximum of $\bigcup a_n$ and not just after $\max(t)$ as would be the case in Mathias forcing.

Definition 2.7. (1) Let $(s, \bar{a}) \in \mathbb{Q}^{242}$. We define a tree $T(s, \bar{a})$ as follows: Elements of the tree are $\{s \cup t : (\exists n \in \omega)(\exists i_0 < i_1 \cdots < i_{n-1})(t \in a_{i_0} \circ \cdots \circ a_{i_{n-1}})\}$. The tree is ordered by end extension.

(2) We let

$$\text{Lev}_{<k}(s, \bar{a}) = \{s \cup t : (\exists n \leq k)(\exists i_0 < i_1 \cdots < i_{n-1} < k)(t \in a_{i_0} \circ \cdots \circ a_{i_{n-1}})\}.$$

In other words, $t \in T(s, \bar{a})$ iff $t = s$ or $(t, \bar{a} \text{ past } \max(t) + 1) \leq (s, \bar{a})$. Here and henceforth we write $(t, \bar{a} \text{ past } k)$ for $(t, (\bar{a} \text{ past } k))$.

Definition 2.8. Let $n \in \omega$, $(s, \bar{a}), (t, \bar{b}) \in \mathbb{Q}^{242}$ and assume $(t, \bar{b}) \leq (s, \bar{a})$. We say (t, \bar{b}) is a 0-extension of (s, \bar{a}) and write $(t, \bar{b}) \leq_0 (s, \bar{a})$ iff $t = s$.

3. BLASS-SHELAH FORCING WITH SUITABLE SETS

Now we thin out the reservoir $(\mathcal{P})^\omega$ to a suitable subfamily. We choose \mathcal{H} in allusion to a happy family.

Definition 3.1. We assume that $\langle \bar{a}_n : n \in \omega \rangle$ is a \leq -descending sequence of members $\bar{a}_n \in (\mathbb{P})^\omega$. We say \bar{b} is a *diagonal lower bound* of $\langle \bar{a}_n : n \in \omega \rangle$ iff for any $n \in \omega$, we have

$$(\bar{b} \text{ past } n) \leq \bar{a}_n.$$

Remark 3.2. If \bar{b} is a diagonal lower bound of $\langle \bar{a}_n : n \in \omega \rangle$ then $\bar{b} \leq \bar{a}_0$. If $\bar{c} \leq \bar{b}$ and \bar{b} is a diagonal lower bound then also \bar{c} is a diagonal lower bound.

Definition 3.3. Compare to Mathias [10]. A set $\mathcal{H} \subseteq (\mathcal{P})^\omega$ is called a *suitable set* if the following hold:

- (1) (Non-emptiness, freeness, upwards closure) $\mathcal{H} \subseteq (\mathcal{P})^\omega$, $\mathcal{H} \neq \emptyset$. If $\bar{a} \in \mathcal{H}$ and $\bar{b} \geq^* \bar{a}$ then $\bar{b} \in \mathcal{H}$.
- (2) (Existence of diagonal lower bounds) Any \leq -descending ω -sequence in \mathcal{H} has a diagonal lower bound in \mathcal{H} .

- (3) (Fullness) For any $Y \subseteq \omega$ and $\bar{a} \in \mathcal{H}$ there is $\bar{b} \leq \bar{a}$, $\bar{b} \in \bar{a}$ such that $\text{set}(\bar{b}) \subseteq Y$ or $\text{set}(\bar{b}) \subseteq Y^c$.
- (4) (Ramsey property) For any $C: [\omega]^{<\omega} \rightarrow 2$ and any (s, \bar{a}) with $\bar{a} \in \mathcal{H}$ and $\max(s) < \min(\bigcup a_0)$ there is $(t, \bar{b}) \leq (s, \bar{a})$, $\bar{b} \in \mathcal{H}$ such that either $C \upharpoonright T(t, \bar{b})$ is constantly 1 or $C \upharpoonright (T(s, \bar{b}) \setminus \{s\})$ is constantly 0.

Here is a family of subforcings of \mathbb{Q}^{242} :

Definition 3.4. Given a suitable set \mathcal{H} in $(\mathcal{P})^\omega$, the notion of forcing $\mathbb{Q}^{242}(\mathcal{H})$ consists of all pairs $(s, \bar{a}) \in \mathbb{Q}^{242}$ such that $\bar{a} \in \mathcal{H}$. The order relation is as in \mathbb{Q}^{242} (see Def. 2.6).

In Definition 2.2(3), we take the limit requirement following [4]. All the proofs in this paper could be carried out equally with the requirement $\limsup_n \text{norm}(a_n) = \infty$ instead. We do not know any difference in the effect of the two variants. For the full forcing with $(\mathcal{P})^\omega$, the two variants are equivalent, since any sequence with $\limsup_n (a_n) = \infty$ has a subsequence that satisfies the lim-requirement.

Now we repeat some propositions from [4] for $\mathbb{Q}^{242}(\mathcal{H})$. Our proofs are slightly different from the original proofs since the second coordinates of conditions need to be elements of the given family \mathcal{H} and \mathcal{H} is less closed than $(\mathcal{P})^\omega$. We let $\chi \geq (2^\omega)^+$ be a regular cardinal and we fix a well-ordering \triangleleft of $H(\chi)$.

Lemma 3.5. (See [4, Proposition 2.4]) *Let τ_i , $i \in \omega$, be $\mathbb{Q}^{242}(\mathcal{H})$ names for ordinals. Then every condition (s, \bar{a}) has a 0-extension (s, \bar{b}) with the following property: If $\ell \geq 1$ and $t \in \text{Lev}_{<\ell}(T(s, \bar{a})) \cap T(s, \bar{b})$ and $i \leq \max(\bigcup a_{\ell-1})$ and $(t, \bar{b} \text{ past } \max(\bigcup a_{\ell-1}) + 1)$ has a 0-extension forcing a particular value for τ_i , then $(t, \bar{b} \text{ past } \max(\bigcup a_{\ell-1}) + 1)$ forces a particular value for τ_i .*

Proof. By induction on ℓ we choose a \leq -decreasing sequence $\langle \bar{a}_\ell : \ell \in \omega \rangle$ of elements of \mathcal{H} . We start with $\bar{a}_0 = \bar{a}$. We let $n(\ell) = \max(\bigcup a_\ell) + 1$ and $n(-1) = 0$. Suppose $\bar{a}_{n(\ell-1)}$, $\ell \geq 0$, is chosen. Let $\{(t_j, i_j) : j \leq k\}$ be the \triangleleft -least enumeration of $\text{Lev}_{<\ell+1}(T(s, \bar{a})) \times n(\ell)$.

Now by a subinduction on $j \leq k$ we choose \bar{a}^j , $j = 0, \dots, k$. We start with $\bar{a}^0 = (\bar{a}_{n(\ell-1)} \text{ past } n(\ell))$. Given \bar{a}^j we do the following: If there is $(t_j, \bar{b}) \leq (t_j, \bar{a}^j_{\ell-1})$ forcing a value to τ_{i_j} then we let \bar{a}^{j+1} be the \triangleleft -least such \bar{b} . Otherwise we let $\bar{a}^{j+1} = \bar{a}^j$. In the end we let $\bar{a}_{n(\ell-1)+1} = \dots = \bar{a}_{n(\ell)} = \bar{a}^k$. This ends the subinduction.

Having defined $\langle \bar{a}_\ell : \ell < \omega \rangle$ we let $\bar{b} \in \mathcal{H}$ be a the \triangleleft -least diagonal lower bound in \mathcal{H} to the sequence $\langle \bar{a}_\ell : \ell \in \omega \rangle$. Then (s, \bar{b}) has the desired properties: If $\ell \geq 1$ and $t \in \text{Lev}_{<\ell}(T(s, \bar{a}))$, $t \in T(s, \bar{b})$ and $i < n(\ell-1)$ and there is a 0-extension of $(t, \bar{b} \text{ past } n(\ell-1))$ forcing a value to τ_i then, since $(\bar{b} \text{ past } n(\ell-1)) \leq \bar{a}_{n(\ell-1)}$ there is a 0-extension of $(t, \bar{a}_{n(\ell-1)})$ forcing a value to τ_i . Hence by construction $(t, \bar{a}_{n(\ell-1)})$ itself forces a value to τ_i . Since $(\bar{b} \text{ past } n(\ell-1)) \leq \bar{a}_{n(\ell-1)}$, the condition $(t, \bar{b} \text{ past } n(\ell-1))$ forces a value to τ_i . \square

Lemma 3.6. *Assume that τ_i , $i \in \omega$, and (s, \bar{a}) are as in the previous lemma. Any \bar{b} as in the conclusion of the previous lemma has the following property: For any $\ell \geq 1$, if $t \in \text{Lev}_{<\ell}(T(s, \bar{b}))$ and $i \leq \max(\bigcup b_{\ell-1})$ and $(t, \bar{b} \text{ past } \max(\bigcup b_{\ell-1}) + 1)$ has a 0-extension*

forcing a particular value for τ_i , then $(t, \bar{b} \text{ past } \max(\bigcup b_{\ell-1}) + 1)$ forces a particular value for τ_i .

Proof. For any $\ell \geq 1$ there is $\ell' \geq 1$ such that $\max(\bigcup b_\ell) \leq \max(\bigcup a_{\ell'})$ and $a_{\ell'} < b_{\ell+1}$. By the definitions of \leq and of $\bar{a} \mapsto (\bar{a} \text{ past } n)$, for any $t \in \text{Lev}_{<\ell}(t, \bar{b})$ we have $(t, \bar{b} \text{ past } \max(\bigcup b_\ell) + 1) \leq (t, \bar{b} \text{ past } \max(\bigcup a_{\ell'}) + 1)$. Hence we are done. \square

Lemma 3.7. $\mathbb{Q}^{242}(\mathcal{H})$ is proper.

Proof. This is derived from Lemma 3.5 as in [4, Proposition 2.5]. Indeed, given $M \prec (H(\chi), \in, \triangleleft)$ and an enumeration $\tau_i, i \in \omega$, of all names in M of ordinals and $(s, \bar{a}) \in M$, the condition \bar{b} constructed in Lemma 3.5 is $(M, \mathbb{Q}^{242}(\mathcal{H}))$ -generic. \square

Can we work with weaker properties than suitability? We do not know. At least the requirement of fullness seems to be natural. In order to explain this we name the generic reals:

Definition 3.8. Let G be $\mathbb{Q}^{242}(\mathcal{H})$ -generic over \mathbf{V} . We call

$$W_G = \bigcup \{s : \exists \bar{a}(s, \bar{a}) \in G\}$$

the $\mathbb{Q}^{242}(\mathcal{H})$ -generic real and let W be a name for it.

The generic real of the full Blass–Shelah forcing \mathbb{Q}^{242} is not split by any real in the ground model:

Lemma 3.9. If $\bar{a} \in (\mathcal{P})^\omega$ and $X \subseteq \omega$ then there is $\bar{b} \leq \bar{a}$ such that $\text{set}(\bar{b}) \subseteq X$ or $\text{set}(\bar{b}) \subseteq (\omega \setminus X)$.

Proof. By definition, for $a \in \mathcal{P}$, $\text{norm}(a \upharpoonright X) \geq \text{norm}(a) - 1$ or $\text{norm}(a \upharpoonright X^c) \geq \text{norm}(a) - 1$. Now we choose $Y = X$ or $Y = X^c$ such that $C = \{n : \text{norm}(a_n \upharpoonright Y) \geq \text{norm}(a_n) - 1\}$ is infinite. Let $(n_k)_k$ enumerate C and let $\bar{b} = \langle a_{n_k} \upharpoonright Y : k \in \omega \rangle$. \square

By the fullness requirement we have the same result for the subforcing $\mathbb{Q}^{242}(\mathcal{H})$:

Lemma 3.10. Let \mathcal{H} be a suitable set. In $\mathbf{V}^{\mathbb{Q}^{242}(\mathcal{H})}$, for any $X \subseteq \omega$, $X \in \mathbf{V}$, $W_G \subseteq^* X$ or $W_G \subseteq X^c$.

Proof. Given $p = (s, \bar{a}) \in \mathbb{Q}^{242}(\mathcal{H})$ and $X \subseteq \omega$, by fullness there is $\bar{b} \leq \bar{a}$, $\bar{b} \in \mathcal{H}$ with $\text{set}(\bar{b}) \subseteq X$ or $\text{set}(\bar{b}) \subseteq X^c$. Hence $D = \{p : p \Vdash W \subseteq^* X \text{ or } p \Vdash W \subseteq^* X^c\}$ is dense. \square

So in the generic extension we have the set $\mathcal{U}' := \{X \in \mathbf{V} \cap [\omega]^\omega : W_G \subseteq^* X\}$ that decides each old subset of ω . If $\mathcal{U}' \in \mathbf{V}$ then \mathcal{U}' can serve as the ultrafilter \mathcal{U} that is diagonalised, as required in Theorem 1.1. This ends the discussion of fullness. Further below we work with centred suitable families, for which $\mathcal{U}' \in \mathbf{V}$.

Now we work towards preserving a given P -point.

Definition 3.11. Let $\mathcal{S}, \mathcal{S}' \subseteq [\omega]^\omega$ be closed under almost supersets. We write $\mathcal{S} \leq_{\text{RB}} \mathcal{S}'$ and say \mathcal{S} is *Rudin-Blass-below* \mathcal{S}' iff there is a finite-to-one f such that $f(\mathcal{S}) \subseteq f(\mathcal{S}')$.

This definition with f on both sides follows [6]. The frequently used variant of the definition in which $f(\mathcal{S}) \subseteq f(\mathcal{S}')$ is replaced by $f(\mathcal{S}) \subseteq \mathcal{S}'$ has very similar preservation properties when used for ultrafilters \mathcal{S}' .

Definition 3.12. We assume that \mathcal{V} is a P -point and $\mathcal{H} \subseteq (\mathcal{P})^\omega$. We let for $\bar{a} \in \mathcal{H}$, $\mathcal{H} \upharpoonright \bar{a} = \{\bar{b} \in \mathcal{H} : \bar{b} \leq \bar{a}\}$. We say \mathcal{H} *avoids* \mathcal{V} iff $(\forall \bar{a} \in \mathcal{H})(\Phi_2(\mathcal{H} \upharpoonright \bar{a}) \not\leq_{\text{RB}} \mathcal{V})$.

Since \mathcal{V} is an ultrafilter, \mathcal{H} avoids \mathcal{V} holds iff for any finite-to-one function h and any $\bar{a} \in \mathcal{H}$ there is $V \in \mathcal{V}$ and $\bar{b} \in \mathcal{H} \upharpoonright \bar{a}$ such that $h[\text{set}(\bar{b})] \cap h[V] = \emptyset$.

Definition 3.13. (1) Let $\langle \bar{a}_n : n \in \omega \rangle$ be a \leq -descending sequence of elements on $(\mathcal{P})^\omega$. A sequence $\bar{b} \in (\mathcal{P})^\omega$ is called a lower bound iff $(\forall n \in \omega)(\bar{b} \leq^* \bar{a}_n)$.
 (2) $\mathcal{H} \subseteq (\mathcal{P})^\omega$ is called stable if any \leq -descending sequence of elements of \mathcal{H} has a lower bound in \mathcal{H} .

Since diagonal lower bounds are lower bounds the double projection $\Phi_2(\mathcal{H})$ of any suitable \mathcal{H} contains lower bounds for \subseteq^* -descending sequences. So if the double projection of a suitable family \mathcal{H} is an ultrafilter, it is a P -point.

The following deep theorem is crucial for the construction and the evaluation of the forcing.

Theorem 3.14. ([4, Theorem 2.6], see also [2, Theorem 7.4.20]) $(\mathcal{P})^\omega$ has the Ramsey property.

The full set $(\mathcal{P})^\omega$ is suitable.

In preparation for the next theorem we need a consequence of the Ramsey property:

Proposition 3.15. (See [4, Prop. 2.9]) Let \underline{A} be a $\mathbb{Q}^{242}(\mathcal{H})$ -name for a subset of ω . Then every condition (s, \bar{a}) has an extension (t, \bar{b}) with the following property: If $\ell \geq 1$ and if $t' \in \text{Lev}_{<\ell}(T(t, \bar{b}))$, and if $i \leq \max(b_{\ell-1})$, then $(t, \bar{b} \text{ past } \max(b_{\ell-1}) + 1)$ decides whether $i \in \underline{A}$.

Proof. We let $\tau_i = 0$ iff $i \notin \underline{A}$ and $\tau_i = 1$ else. We assume that (s, \bar{a}) already has the property stated of (s, \bar{b}) in Lemma 3.6 for the sequence $\tau_i, i \in \omega$. We define $C: T(s, \bar{a}) \rightarrow 2$ by

$$C(t) = 0 \text{ iff } (\forall \ell \geq 1)(t \in \text{Lev}_{<\ell}(T(s, \bar{a})) \rightarrow (\forall i \leq \max(\bigcup a_{\ell-1}))((t, \bar{a} \text{ past } \max(\bigcup a_{\ell-1}) + 1) \text{ decides } i \in \underline{A})).$$

By the Ramsey property there is $(t, \bar{b}) \leq (s, \bar{a})$ such that for each $t' \in T(t, \bar{b})$, $C(t') = 0$ or there for each $t' \in T(s, \bar{b}) \setminus \{s\}$, $C(t') = 1$. The second possibility is rule out: Let $\ell \geq 1$ and $(t', \bar{c}) \leq (t, \bar{b})$, $t' \in \text{Lev}_{<\ell}(T(s, \bar{a}))$ be such that it decides $i \in \underline{A}$ and $i \leq \max(\bigcup a_{\ell-1})$. Then $(t', \bar{c}) \leq_0 (t', \bar{b} \text{ past } \max(\bigcup a_{\ell-1}) + 1) \leq_0 (t', \bar{a} \text{ past } \max(\bigcup a_{\ell-1}) + 1)$ and the latter already decides $i \in \underline{A}$.

So we have the first possibility. We fix some $t' \in T(s, \bar{b}) \setminus \{s\}$, $t' \in \text{Lev}_{<\ell_0}(T(s, \bar{a}))$. Then $(t', \bar{b} \text{ past } \max(\bigcup a_{\ell_0-1}) + 1)$ has the following property: If $\ell \geq 1$ and if $t'' \in (\text{Lev}_{<\ell}(T(s, \bar{a})) \cap T(t', \bar{b}))$, and if $i \leq \max(\bigcup a_{\ell-1})$, then $(t'', \bar{b} \text{ past } \max(\bigcup a_{\ell-1}) + 1)$ decides whether $i \in A$.

Now as in the proof of Lemma 3.6 we go over from the cut-points $\langle \max(\bigcup a_\ell) : \ell \geq 1 \rangle$ to the cut-points $\langle \max(\bigcup b_\ell) : \ell \geq 1 \rangle$ and thus see that $(t', \bar{b} \text{ past } \max(\bigcup a_{\ell_0-1}) + 1)$ has the desired properties. \square

We recall a very useful theorem:

Theorem 3.16. (*Eisworth* [6, “ \rightarrow ” Theorem 4, “ \leftarrow ” Cor. 2.5, this direction works also with non- P ultrafilters]) *Let \mathcal{U} be a stable ordered-union ultrafilter over $[\omega]^{<\omega} \setminus \{\emptyset\}$ and let \mathcal{V} be a P -point. Then we have: $\Phi(\mathcal{U}) \not\leq_{RB} \mathcal{V}$ iff \mathcal{V} continues to generate an ultrafilter after we force with $\mathbb{M}(\mathcal{U})$.*

Here $\mathbb{M}(\mathcal{U})$ stands for the Matet forcing [9] with a stable ordered-union ultrafilter \mathcal{U} , see [6]. Stable ordered-union ultrafilters are also called Milliken–Taylor ultrafilters. They are ultrafilters over the space \mathbb{F} of non-empty finite subsets of ω . The projection to a filter over ω is $\Phi(\mathcal{U}) = \{X : \exists R \in \mathcal{U}, X \supseteq \bigcup R\}$. For more details we refer to [6]. Stable ordered-union ultrafilters and the projection function Φ will not be used in this work.

Here is an analogue theorem for $\mathbb{Q}^{242}(\mathcal{H})$:

Theorem 3.17. *Let \mathcal{V} be a P -point and let $\mathcal{H} \subseteq (\mathcal{P})^\omega$ be a suitable set that avoids \mathcal{V} . Then \mathcal{V} continues to generate an ultrafilter after we force with $\mathbb{Q}^{242}(\mathcal{H})$.*

Proof. We adapt the proof of [4, Theorem 3.3]. Let A be a name for a subset of ω . By genericity, it suffices to show that, if (s, \bar{a}) forces $A \subseteq \omega$, then some extension forces either $B \subseteq A$ or $B \subseteq A^c$, for some $B \in \mathcal{V}$. According to Proposition 3.15, we may assume that, for $\ell \geq 1$, $i \leq \max(\bigcup a_{\ell-1})$, $t \in \text{Lev}_{<\ell}(T(s, \bar{a}))$ the condition $(t, \bar{a} \text{ past } \max(\bigcup a_{\ell-1}) + 1)$ decides whether $i \in A$. Consider any $t \in T(s, \bar{a})$.

Then $t \in \text{Lev}_{<\ell}(T(s, \bar{a}))$ for all sufficiently large ℓ . Thus, for any fixed $i \in \omega$, $(t, \bar{a} \text{ past } z)$ will decide whether $i \in A$ once z is large enough; the decisions agree as z varies, since $(t, \bar{a} \text{ past } z')$ extends $(t, \bar{a} \text{ past } z)$ if $z' \geq z$. Let $A(t)$ be the set of those $i \in \omega$ for which the decision is positive.

Partition $T(s, \bar{a})$ by putting into one class all those $t \in T(s, \bar{a})$ for which $A(t) \in \mathcal{V}$. By Theorem 3.14, we can extend (s, \bar{a}) to some $(s', \bar{b}) \in \mathbb{Q}^{242}(\mathcal{H})$ such that all of $T(s', \bar{b})$ is in a single class. When we form this extension, we do not destroy the fact that, for $t \in T(s', \bar{b})$ for $i \in A(t)$ (resp. $i \notin A(t)$), $(t, \bar{b} \text{ past } z) \Vdash i \in A$ ($i \notin A$) for all sufficiently large z . We assume henceforth that $A(t) \in \mathcal{V}$ for all $t \in T(s', \bar{b})$; in the other case A is replaced by its complement. As \mathcal{V} is a P -point, let $B \in \mathcal{V}$ be almost included in each $A(t)$.

Let for $\ell \geq 0$, $n(\ell) = \max(\bigcup b_\ell) + 1 \geq \ell + 1$. Inductively we define a sequence $\langle \zeta(k) : k \in \omega \rangle$ of natural numbers, starting with $\zeta(0) = 1$, and increasing so rapidly that, if $t \in \text{Lev}_{<\zeta(k)}(T(s', \bar{b}))$, then

- (i) $B \setminus A(t) \subseteq \zeta(k+1)$, and
- (ii) $\zeta(k+1) \geq n(\zeta(k))$.

We think of $n \circ \zeta$ as partitioning ω into blocks $[n(\zeta(k)), n(\zeta(k+1))]$, $k \in \omega$, and consider the four sets X_i , $i = 0, 1, 2, 3$, obtainable by taking the union of every fourth block:

$$X_i = \bigcup \{[n(\zeta(j)), n(\zeta(j+1))] : j = i \pmod{4}\}.$$

As \mathcal{V} is an ultrafilter, it contains exactly one of these sets. By omitting a few terms (at most 3) from the sequence ζ , we may assume $i = 2$. Replacing B with $X_2 \cap B$, which is also in \mathcal{V} , we may assume $B \subseteq X_2$.

Let h_1 map $i \in [n(\zeta_{2k}), n(\zeta_{2k+2})]$ to $h_1(i) = k$, let h_2 map $i \in [n(\zeta_{2k+1}), n(\zeta_{2k+3})]$ to $h_2(i) = k$. Now by $h_1(\Phi_2(\mathcal{H} \upharpoonright \bar{b})) \not\subseteq h_1(\mathcal{V})$ there are $\bar{c}_1 \in \mathcal{H}$, $\bar{c}_1 \leq \bar{b}$ and $B' \in \mathcal{V}$, $B' \subseteq B \cap X_2$, such that

$$h_1[\text{set}(\bar{c}_1)] \cap h_1[B'] = \emptyset.$$

Now by $h_2(\Phi_2(\mathcal{H} \upharpoonright \bar{c}_1)) \not\subseteq h_2(\mathcal{V})$ there are $\bar{c} \in \mathcal{H}$, $\bar{c} \leq \bar{c}_1$ and $B'' \in \mathcal{V}$, $B'' \subseteq B' \cap X_2$, such that

$$(iii) \quad h_i[\text{set}(\bar{c})] \cap h_i[B''] = \emptyset \text{ for } i = 1, 2.$$

To complete the proof of the theorem, we show that (s', \bar{c}) forces $B'' \subseteq A$.

We fix an element $i \in B''$ and an extension (t, \bar{d}) of (s', \bar{c}) deciding whether $i \in A$. Since $B'' \subseteq B \subseteq X_2$, there is k such that $i \in [n(\zeta(4k+2)), n(\zeta(4k+3))]$. We fix k . By the choice of \bar{b} we know that we can assume that $\bar{d} = \bar{d} \text{ past } n(\zeta(4k+4))$ and that $t \in \text{Lev}_{<\zeta(4k+4)}(T(s', \bar{b}))$ and $t \in T(s', \bar{c})$. We show that the decision is positive. Now by (iii), the set $\text{set}(\bar{c})$ avoids the interval $[n(\zeta(4k+1)), n(\zeta(4k+4))]$, and hence $t \in \text{Lev}_{<\zeta(4k+1)}(T(s', \bar{b}))$.

Since $t \in \text{Lev}_{<\zeta(4k+1)}(T(s', \bar{b}))$ and $i \in B''$ and $i \geq n(\zeta(4k+2)) \geq \zeta(4k+2)$, clause (i) in the definition of ζ implies that $i \in A(t)$. Since also $t \in \text{Lev}_{<\zeta(4k+3)}(T(s', \bar{b}))$ and $i < n(\zeta(4k+3))$, clause (ii) in the definition of ζ implies $n(\zeta(4k+4)) > \zeta(4k+4) \geq n(\zeta(4k+3))$ and $(t, \bar{b} \text{ past } n(\zeta(4k+4))) \Vdash i \in A$ and hence $(t, \bar{d}) \Vdash i \in A$. \square

Definition 3.18. (1) A subset $\mathcal{C} \subseteq (\mathcal{P})^\omega$ is called centred if any finitely many members of \mathcal{C} have a common lower bound in \mathcal{C} .

(2) A centred subset $\mathcal{C} \subseteq (\mathcal{P})^\omega$ is called maximal if for any $\bar{a} \notin \mathcal{C}$ there is $\bar{b} \in \mathcal{C}$, $\bar{a} \perp^* \bar{b}$.

In the case of a suitable centred family \mathcal{C} , we have: \mathcal{C} avoids \mathcal{V} iff $\Phi_2(\mathcal{C}) \not\subseteq_{RB} \mathcal{V}$. Now we read the theorem for the special case of $\mathcal{H} = \mathcal{C}$ and thus finish the proof of Theorem 1.1.

4. EXISTENCE OF CENTRED SUITABLE \mathcal{H} UNDER CH OR MA

In this section we show that under CH, given a P -point there is a suitable maximal centred family \mathcal{C} such that \mathcal{C} avoids the given P -point. A natural \leq^* -descending construction will give just a stable family. We add diagonal lower bounds in the family by explicit construction steps.

Remark 4.1. If we replace the lim-requirement in Def.2.2(3) by the weaker lim sup-requirement then maximal centred sets (in the altered space) are full.

Proof. Let \mathcal{C} be a maximal centred suitable set. Let $X \subseteq \omega$ be given. By the definition of the norm and the modified definition of $(\mathcal{P})^\omega$, for any $\bar{c} \in \mathcal{C}$ we have $\bar{c} \upharpoonright X \in (\mathcal{P})^\omega$ or $\bar{c} \upharpoonright X^c \in (\mathcal{P})^\omega$. Since \mathcal{C} is centred, we have for $Y = X$ or for $Y = X^c$, for any $\bar{c} \in \mathcal{C}$, $\bar{c} \upharpoonright Y \in (\mathcal{P})^\omega$. So $\mathcal{C}' = \{\bar{b} : (\exists \bar{c} \in \mathcal{C})(\bar{b} \geq^* \bar{c} \upharpoonright Y)\}$ is a centred suitable set. Since \mathcal{C} was maximal and $\mathcal{C}' \supseteq \mathcal{C}$, we have $\mathcal{C}' = \mathcal{C}$. So there is $\bar{d} \in \mathcal{C}$ with $\bar{d} = \bar{c} \upharpoonright Y$ and hence $\text{set}(\bar{d}) \subseteq Y$. \square

However, in the original definition of the space $(\mathcal{P})^\omega$ we do not know whether maximal centred families are full. For $\mathcal{H} = (\mathcal{P})^\omega$, the lim and the lim sup requirement on the norms $\text{norm}(a_n)$, $n \in \omega$, give equivalent forcings. We do not know whether this is still true for arbitrary (suitable) families.

If \mathcal{C} is a suitable maximal centred set, then forcing with $\mathbb{Q}^{2^{2^2}}(\mathcal{C})$ diagonalises the ultrafilter $\Phi_2(\mathcal{C})$ by adding W_G . Hence the fullness and the maximality of \mathcal{C} are destroyed.

Lemma 4.2. *Under CH or under Martin's Axiom for $< 2^\omega$ dense sets, given an \mathbb{P} -point \mathcal{V} there is a suitable maximal centred set \mathcal{C} such that \mathcal{C} avoids \mathcal{V} .*

Proof. Let $\langle (C_\alpha, n_\alpha, D_\alpha, X_\alpha, \bar{a}_\alpha, h_\alpha) : \alpha < 2^\omega \rangle$ enumerate all tuples (C, n, D, X, \bar{a}, h) such that $C : [\omega]^{<\omega} \rightarrow 2$ is a colouring, $n \in \omega$, $D = \langle \bar{d}_n : n \in \omega \rangle$ is a \leq -descending sequence, $X \subseteq \omega$, $\bar{a} \in (\mathcal{P})^\omega$ and h is a finite-to-one function, and such that each tuple appears cofinally often in the enumeration. By induction on $\alpha < 2^\omega$ we choose $\bar{c}_\alpha \in (\mathcal{P})^\omega$ such that $(\forall \beta < \alpha) \bar{c}_\alpha \leq^* \bar{c}_\beta$. We let \bar{c}_0 be any element of $(\mathcal{P})^\omega$.

In the successor steps, given \bar{c}_α and C_α we first take care of the Ramsey property: By Theorem 3.15 we can choose $\bar{c}_{\alpha+0.3} \leq (\bar{c}_\alpha \text{ past } n_\alpha)$ such that for some s , we have $(s, \bar{c}_{\alpha+0.3}) \leq (\emptyset, \bar{c}_\alpha \text{ past } n_\alpha)$ and the colouring C_α is monochromatic with colour 0 on $T(\emptyset, \bar{c}_{\alpha+0.3}) \setminus \{\emptyset\}$ or C_α is monochromatic with colour 1 on $T(s, \bar{c}_{\alpha+0.3})$.

Now we take care of the diagonal lower bounds: Given $D_\alpha = \langle \bar{d}_n : n \in \omega \rangle$, we distinguish two cases: First case: Each $\bar{d}_n \geq^* \bar{c}_{\alpha+0.3}$. We let $\bar{d}_n \geq (\bar{c}_{\alpha+0.3} \text{ past } j(n))$ for an increasing function $j(n)$. We choose $\bar{c}_{\alpha+0.5} = \langle c_k : k \in \omega \rangle$ by induction on k . We let $c_0 = c_{\alpha+0.3, i}$ with an i such that $c_{\alpha+0.3, i}$ lies past $j(0)$ and has norm at least 1. For $k \geq 0$, given c_k , we take c_{k+1} such that $c_{k+1} = c_{\alpha+0.3, i}$ with i so large that c_{k+1} lies past $j(\max(\bigcup c_k) + 1)$ and has norm at least $k+1$. By construction $\bar{c}_{\alpha+0.5} = \bar{c}$ is a diagonal lower bound of D and $\bar{c}_{\alpha+0.5} \leq \bar{c}_{\alpha+0.3}$. Second case: There is n such that $\bar{d}_n \not\geq^* \bar{c}_{\alpha+0.3}$. Then we let $\bar{c}_{\alpha+0.5} = \bar{c}_{\alpha+0.3}$.

Next we ensure an instance of fullness: We choose $\bar{c}_{\alpha+0.7} \leq \bar{c}_{\alpha+0.5}$ that that $\text{set}(\bar{c}_{\alpha+0.7})$ is a subset of X_α or of X_α^c .

Next we take care of the task $h_\alpha(\Phi_2(\mathcal{C})) \not\subseteq h_\alpha(\mathcal{V})$. The set $\{h_\alpha^{-1}[\text{set}(b)] : \bar{b} \leq \bar{c}_{\alpha+0.9}\}$ is an analytic set and by the Luzin–Sierpinski theorem (see e.g. [7, Theorem 21.6]) the set $\{h_\alpha^{-1}[X] : X \in \mathcal{V}\}$ is not analytic. By the definition of the norm, the first set contains $h_\alpha^{-1}(\mathcal{U})$ for some free ultrafilter \mathcal{U} . Hence the $\{h_\alpha^{-1}[\text{set}(b)] : \bar{b} \leq \bar{c}_{\alpha+0.9}\}$ is not a subset of $\{h_\alpha^{-1}[X] : X \in \mathcal{V}\}$. Thus there is an $\bar{c}_{\alpha+1} \leq \bar{c}_{\alpha+0.7}$ such that $\omega \setminus h_\alpha^{-1}[\text{set}(\bar{c}_{\alpha+1})] \in \mathcal{V}$. This finishes the successor step.

In the limit steps $\alpha < 2^\omega$ of countable cofinality, we choose a cofinal sequence α_n , $n \in \omega$, converging to α , and then we take as \bar{c}_α a \leq^* -lower bound of \bar{c}_{α_n} , $n \in \omega$. If the continuum is larger and Martin's Axiom holds, in the limit steps α of uncountable cofinality in the

construction we force a lower bound with the following σ -centred approximation forcing \mathbb{Q} : Conditions are pairs (\bar{a}, F) such that $\bar{a} \in (\mathcal{P})^{<\omega}$ and F is a finite subset of $\{\bar{c}_\beta : \beta < \alpha\}$. We let $(\bar{b}, F') \leq_{\mathbb{Q}} (\bar{a}, F) = (\langle a_0, \dots, a_{n-1} \rangle, F)$ iff $F' \supseteq F$ and any element of $\bar{b} \setminus \bar{a}$ is for every $\bar{c} \in F$ an element of a condensation of \bar{c} and has norm at least $\text{norm}(a_{n-1}) + 1$ (and norm at least 1 if $n = 0$).

Having chosen \bar{c}_α , $\alpha < 2^\omega$, we let

$$\mathcal{C} = \{\bar{a} : (\exists \alpha < 2^\omega)(\bar{a} \geq^* \bar{c}_\alpha)\}.$$

It is clear that \mathcal{C} is centred, maximal and full and closed under diagonal lower bounds. We show the Ramsey property. Let $C: [\omega]^{<\omega} \rightarrow 2$ and (s, \bar{a}) be given such that $\bar{a} \in \mathcal{C}$. We take $\alpha < 2^\omega$ such that $\bar{a} \geq (\bar{c}_\alpha \text{ past } n_\alpha)$, $\max(s) < n_\alpha$, and $C_\alpha(t) = C(s \cup t)$ for $t \in T(\emptyset, \bar{c}_\alpha \text{ past } n_\alpha)$. By construction there is some t , $(t, \bar{c}_{\alpha+0.3}) \leq (\emptyset, \bar{c}_\alpha \text{ past } n_\alpha)$ such that the colouring C_α is monochromatic with colour 0 on $T(\emptyset, \bar{c}_{\alpha+0.3}) \setminus \{\emptyset\}$ or that C_α is monochromatic with colour 1 on $T(t, \bar{c}_{\alpha+0.3})$. Then C is monochromatic with colour 0 on $T(s, \bar{c}_{\alpha+0.3}) \setminus \{s\}$ or that C is monochromatic with colour 1 on $T(s \cup t, \bar{c}_{\alpha+0.3})$. \square

Thus we finish the proof of Theorem 1.2.

5. DISCUSSION AND QUESTIONS

Recall in Mathias forcing with an ultrafilter \mathcal{U} conditions are of the form (s, A) , $s \in [\omega]^{<\omega}$, $\max(s) < \min(A)$, $A \in \mathcal{U}$ and $(t, B) \leq (s, A)$ if $t \setminus s \subset A$ and $t \supseteq s$ and $B \subseteq A$.

Our result can be seen as an answer to the following question.

Question 5.1. *Is there a relative of Mathias forcing with an ultrafilter \mathcal{U} that preserves another ultrafilter?*

For an affirmative answer the ultrafilter \mathcal{U} cannot be rapid and hence cannot be a Ramsey ultrafilter, as Mathias forcing would add a dominating real and thus destroy any ultrafilter. On the other hand, for preserving a P -point Ramsey theoretic properties of the forcing are often very useful. Thus we provided for Ramsey-theoretic properties by superposing more structure and norms and letting $\Phi_2(\mathcal{H}) = \mathcal{U}$ be an ultrafilter. There is a projection mapping $(s, \bar{a}) \in \mathbb{Q}^{242}(\mathcal{H})$ to $(a, \text{set}(\bar{a}))$. However, we do not know whether this projection can be inverted to a complete embedding.

Another topic is the possibility of iterating forcings of the type $\mathbb{Q}^{242}(\mathcal{H})$. We are interested in a tower of suitable maximal centred \mathcal{C} in successive forcing extensions. For building such a tower it would be some ease of work if we knew that our list of requirements is redundant. Let $\mathcal{H} \subseteq (\mathcal{P})^\omega$.

Question 5.2. *Does “ \mathcal{H} is maximal centred” imply “ \mathcal{H} is full”?*

Question 5.3. *Does the Ramsey property together with the existence of (diagonal) lower bounds for \mathcal{H} imply fullness? If we add maximal centredness to the premises?*

Question 5.4. *Does the Ramsey property together with the existence of (diagonal) lower bounds and centredness for \mathcal{H} imply maximal centredness? If we add fullness to the premises?*

Question 5.5. *Let \mathcal{C} be maximal centred and suitable and avoiding \mathcal{V} . Is there a forcing \mathbb{Q} of size at most 2^ω that preserves \mathcal{V} such that in $\mathbf{V}^{\mathbb{Q}^{2^{42}}(\mathcal{C}) * \mathbb{Q}}$, is there $\mathcal{C}' \supseteq \mathcal{C}$ that is centred and suitable?*

*If so, in the iteration step of countable cofinality of iterands of the type $\mathbb{Q}^{2^{42}}(\mathcal{C}_\alpha) * \mathbb{Q}_\alpha$ is there a suitable maximal centred \mathcal{C} extending the families used in the preceding iterands?*

This question has an interesting sub-question.

Question 5.6. *Is in the extension by a forcing of type $\mathbb{Q}^{2^{42}}(\mathcal{C}) * \mathbb{Q}$ the Ramsey property for a given coloring witnessed by a sequence $\bar{b} \in (\mathcal{P})^\omega$ that is positive in the sense of the family \mathcal{C} ?*

REFERENCES

- [1] Tomek Bartoszyński, Martin Goldstern, Haim Judah, and Saharon Shelah. All meager filters may be null. *Proc. Amer. Math. Soc.*, 117(2):515–521, 1993.
- [2] Tomek Bartoszyński and Haim Judah. *Set Theory, On the Structure of the Real Line*. A K Peters, 1995.
- [3] Andreas Blass. Near coherence of filters, II: Applications to operator ideals, the Stone-Čech Remainder of a half-line, order ideals of sequences, and slenderness of groups. *Trans. Amer. Math. Soc.*, 300:557–581, 1987.
- [4] Andreas Blass and Saharon Shelah. There may be simple P_{\aleph_1} - and P_{\aleph_2} -points and the Rudin-Keisler ordering may be downward directed. *Annals of Pure and Applied Logic*, 33:213–243, 1987.
- [5] David Chodounský, Dušan Repovš, and Lyubomyr Zdomskyy. Mathias forcing and combinatorial covering properties of filters. *J. Symb. Log.*, 80(4):1398–1410, 2015.
- [6] Todd Eisworth. Forcing and stable ordered-union ultrafilters. *J. Symbolic Logic*, 67:449–464, 2002.
- [7] Alexander Kechris. *Classical Descriptive Set Theory*. Number 156 in Graduate text in Mathematics. Springer-Verlag, Heidelberg New York, 1995.
- [8] Jakob Kellner and Saharon Shelah. Preserving preservation. *J. Symbolic Logic*, 70(3):914–945, 2005.
- [9] Pierre Matet. Partitions and filters. *J. Symbolic Logic*, 51:12–21, 1986.
- [10] Adrian Mathias. Happy families. *Ann, Math. Logic*, 12:59–111, 1977.
- [11] Heike Mildenberger and Saharon Shelah. The principle of near coherence of filters does not imply the filter dichotomy principle. *Trans. Amer. Math. Soc.*, 361:2305–2317, 2009.
- [12] Saharon Shelah. Nice \aleph_1 generated non- P -points, I. *Mathematical Logic Quarterly*, submitted/under revision.

HEIKE MILDENBERGER, ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG, MATHEMATISCHES INSTITUT, ABTEILUNG FÜR MATHEMATISCHE LOGIK, ECKERSTR. 1, 79104 FREIBURG IM BREISGAU, GERMANY
E-mail address: heike.mildenberger@math.uni-freiburg.de