

# Yang-Baxter representations of the infinite symmetric group

Gandalf Lechner, Ulrich Pennig, Simon Wood

School of Mathematics, Cardiff University

July 1, 2017

## Abstract

Every unitary involutive solution of the quantum Yang-Baxter equation (“R-matrix”) defines an extremal character and a representation of the infinite symmetric group  $S_\infty$ . We give a complete classification of all such Yang-Baxter characters and determine which extremal characters of  $S_\infty$  are of Yang-Baxter form.

Calling two involutive R-matrices equivalent if they have the same character and the same dimension, we show that equivalence classes are classified by pairs of Young diagrams, and construct an explicit normal form R-matrix for each class. Using operator-algebraic techniques (subfactors), we prove that two R-matrices are equivalent if and only if they have similar partial traces.

Furthermore, we describe the algebraic structure of the equivalence classes of all involutive R-matrices, and discuss several classes of examples. These include Yang-Baxter representations of the Temperley-Lieb algebra at parameter  $q = 2$ , which can be completely classified in terms of their rank and dimension.

## 1 Introduction

The Yang-Baxter equation originated in quantum mechanics and statistical mechanics, in particular in the works of Yang [Yan67] and Baxter [Bax72]. By now, it is known to be of fundamental importance also in many other areas: Solutions of the Yang-Baxter equation (usually called “R-matrices”) appear in integrable quantum field theory as scattering operators [AAR01], and as quantum logical gates in quantum information theory [KL02]. Any R-matrix defines a representation of the infinite braid group and can give rise to link invariants in knot theory [Tur88, Jon87], as originally discovered in the context of subfactors and Jones’ fundamental construction [Jon83]. Also quasitriangular Hopf algebras define universal R-matrices satisfying the Yang-Baxter equation [CP94].

This list of topics and references is far from exhaustive and serves only as a sample of the vast literature on the subject. For a more complete introduction to the Yang-Baxter equation and further references, see [Jim89].

In its most basic form, the (quantum) Yang-Baxter equation is an equation for an endomorphism  $R \in \text{End}(V \otimes V)$  on the tensor square of a vector space  $V$ , namely

$$(R \otimes \text{id}_V)(\text{id}_V \otimes R)(R \otimes \text{id}_V) = (\text{id}_V \otimes R)(R \otimes \text{id}_V)(\text{id}_V \otimes R) \quad (1.1)$$

as an equation in  $\text{End}(V \otimes V \otimes V)$ . Several variations of this equation exist, such as the classical Yang-Baxter equation [CP94], the Yang-Baxter equation with spectral parameters [Jim89], or the set-theoretic Yang-Baxter equation [LYZ<sup>+</sup>00]. We shall only consider the form (1.1).

Although (1.1) makes perfect sense for infinite-dimensional Hilbert spaces  $V$ , and in fact has many interesting infinite-dimensional solutions<sup>1</sup>, we restrict ourselves to finite-dimensional  $V$ , and agree to write  $d = \dim V$  throughout. We fix a scalar product on  $V$  (and hence its tensor powers), and denote the set of all unitary solutions of (1.1) by  $\mathcal{R}(V)$ . As usual, the elements of  $\mathcal{R}(V)$  are referred to as “R-matrices”. Given some  $R \in \mathcal{R}(V)$ , we also adopt the convention of referring to  $d$  as the “dimension of  $R$ ” – although  $R$  is an endomorphism of a space of dimension  $d^2$  – and to  $V$  as the “base space” of  $R$ .

As is well known, any  $R \in \mathcal{R}(V)$  generates (unitary) representations  $\rho_R^{(n)}$ ,  $n \in \mathbb{N}$ , of the braid groups  $B_n$  on  $V^{\otimes n}$ , by representing the elementary braid<sup>2</sup>  $b_k$ ,  $k \in \{1, \dots, n-1\}$ , as

$$\rho_R^{(n)}(b_k) := \text{id}_V^{\otimes(k-1)} \otimes R \otimes \text{id}_V^{\otimes(n-k-1)} \in \text{End } V^{\otimes n}. \quad (1.2)$$

We set  $\rho_R^{(1)} = \text{id}_V$ . Proceeding to the inductive limit  $B_\infty = \bigcup_n B_n$  of the infinite braid group, every R-matrix defines a homomorphism of  $\mathbb{C}[B_\infty]$  into the infinite (algebraic) tensor product  $\mathcal{E}_0 := \bigcup_n \text{End}(V)^{\otimes n}$ .

In this paper, we focus on representations  $\rho_R^{(n)}$  that factor through the surjective group homomorphism  $B_n \rightarrow S_n$  onto the symmetric group  $S_n$  of  $n$  letters. This is the case if and only if  $R$  is involutive,  $R^2 = 1$ . The simplest involutive R-matrices are, up to a sign, the identity and the tensor flip on  $V \otimes V$ ,

$$R = \pm \text{id}_{V \otimes V}, \quad R = \pm F, \quad F(v \otimes w) = w \otimes v, \quad (1.3)$$

however, infinitely many more exist. Involutive solutions appear in particular in integrable quantum field theory, as symmetries of categories of vector spaces [Lyu87] and in a recent construction of non-commutative spaces [DVL17].

We write  $\mathcal{R}_0(V) \subset \mathcal{R}(V)$  for the subclass of involutive unitary R-matrices, and  $\mathcal{R}$ ,  $\mathcal{R}_0$  for the union of  $\mathcal{R}(V)$ ,  $\mathcal{R}_0(V)$  over all finite-dimensional vector spaces.

---

<sup>1</sup>Note that a solution of the Yang-Baxter equation with spectral parameter can be rewritten as one without parameter, but on an infinite-dimensional base space – see, for example, [HL17, Lemma 2.2].

<sup>2</sup>Recall that a presentation of the braid group  $B_n$  on  $n$  strands is given by  $B_n = \langle b_1, \dots, b_{n-1} : b_i b_{i+1} b_i = b_{i+1} b_i b_{i+1}, b_i b_j = b_j b_i \text{ for } |i-j| > 1 \rangle$ .

One of the aims of this work is to find and classify all elements of  $\mathcal{R}_0$  up to a natural equivalence which has its origins in previous work of the first author in integrable quantum field theory [AL17]. The same notion can be found in [Gur86], and a weaker version of this equivalence relation also played a role in the context of computing all solutions of the Yang-Baxter equation for  $\dim V = 2$  [Hie92].

**Definition 1.1.** *Two R-matrices  $R, S \in \mathcal{R}$  are defined to be equivalent, denoted  $R \sim S$ , if and only if for each  $n \in \mathbb{N}$ , the representations  $\rho_R^{(n)}$  and  $\rho_S^{(n)}$  are equivalent.*

Considering the case  $n = 2$  and taking into account that the only possible eigenvalues of  $R = R^{-1} = R^*$  are  $\pm 1$ , one sees that  $\rho_R^{(2)} \cong \rho_S^{(2)}$  for  $R, S \in \mathcal{R}_0$  if and only if  $R$  and  $S$  have the same dimension and trace. There is an old conjecture of Gurevich to the effect that the converse also holds, that is, that  $R \sim S$  if and only if  $R$  and  $S$  have the same dimension and trace [Gur86, p. 760]. Our findings in this article will in particular disprove this conjecture. The full equivalence  $R \sim S$  is a much stronger condition than having the same dimension and trace, which is reflected in the rich structure of  $\mathcal{R}_0/\sim$  that we find.

Simple examples of equivalent R-matrices can be produced as follows: For any  $R \in \mathcal{R}_0(V)$  and  $A \in \text{GL}(V)$ , one has

$$R \sim (A \otimes A)R(A^{-1} \otimes A^{-1}); \quad (1.4)$$

here  $\rho_R^{(n)}$  and  $\rho_{(A \otimes A)R(A^{-1} \otimes A^{-1})}^{(n)}$  are intertwined by  $A^{\otimes n}$  (and the latter representation is unitary if, for example,  $A$  is unitary). Another example of equivalent R-matrices is given by

$$R \sim FRF, \quad (1.5)$$

where  $F$  is the tensor flip (1.3). Here  $\rho_{FRF}^{(n)}(\iota_n)^{-1}\rho_F^{(n)}(\iota_n)$  is an intertwiner, where  $\iota_n$  is the total inversion. But in general, these two operations do not generate the full equivalence class of  $R \in \mathcal{R}_0$ .

Any  $R \in \mathcal{R}_0$  defines a representation of the infinite symmetric group  $S_\infty$ , the group of all bijections of  $\mathbb{N}$  that move only finitely many points. As any infinite discrete group having no normal abelian subgroup of finite index,  $S_\infty$  admits unitary representations which are not of type I [Tho64b], meaning that its irreducible representations are not classifiable in a reasonable manner. However, it is still possible to classify its extremal characters, corresponding to finite factor representations. Thoma found an explicit parameterization of the extremal characters of  $S_\infty$  [Tho64a], see also [EI16] and the literature cited therein for similar classifications for other “wild” (non-type I) groups. This parameterization depends on countably many continuous variables in a simplex  $\mathbb{T}$ , and we will rely on it to analyze the equivalence  $\sim$ .

Characters and representations of  $S_\infty$  are a topic of ongoing research. We mention here the works of Vershik and Kerov [VK81, VK<sup>+</sup>82], Okounkov and Olshanski [OO98], and in particular the recent monograph [BO17] for an introduction to the subject. Since Thoma’s original work, various new proofs of his results have been obtained [VK81, Oko95, GK10].

We have now introduced sufficient context to state the three main questions that we ask and answer in this article.

- Q1)** How can one classify unitary involutive solutions of the Yang-Baxter equation up to equivalence?
- Q2)** Given  $R, S \in \mathcal{R}_0$ , how to efficiently decide whether  $R \sim S$ ?
- Q3)** Which representations  $\rho$  of  $S_\infty$  are *Yang-Baxter representations*, that is, of the form  $\rho \cong \rho_R$  for some involutive R-matrix  $R \in \mathcal{R}_0$ ?

To answer these questions, we start in Sect. 2 by recalling Thoma's simplex  $\mathbb{T}$  and how it parameterizes the extremal characters of  $S_\infty$ . We show that any involutive R-matrix  $R$  defines an extremal character  $\chi_R$  (Sect. 2.1), but that not every extremal character is of Yang-Baxter form. This poses the question of how to characterize the subset  $\mathbb{T}_{\text{YB}} \subset \mathbb{T}$  parameterizing the Yang-Baxter characters.

Despite the finite-dimensional appearance of the Yang-Baxter equation, a full understanding of  $\mathbb{T}_{\text{YB}}$  requires infinite-dimensional analysis and tools from operator algebras. We consider subfactors arising from a subgroup  $S_\infty^> \subset S_\infty$  in Yang-Baxter representations  $\rho_R$  in Sect. 3. Via this approach, we derive further properties of  $\mathbb{T}_{\text{YB}}$ , and arrive at an answer to Q2) in Thm. 3.3, stating that  $R \sim S$  if and only if these R-matrices have similar partial traces. This section builds on the works of Gohm and Köstler on noncommutative probability and generalizations of Thoma's theorem [GK10, GK11], and of Yamashita on  $S_\infty$ -subfactors [Yam12].

To show that the properties of  $\mathbb{T}_{\text{YB}}$  found up to this point already characterize this set within  $\mathbb{T}$ , we develop in Sect. 4.1 a constructive procedure for generating R-matrices, independent of subfactor theory. The main idea is to find a good replacement for taking direct sums of representations which respects the Yang-Baxter equation. We define a binary operation on  $\mathcal{R}_0$  which enables us to build non-trivial R-matrices from the trivial ones  $\pm \text{id}_{V \otimes V}$ . This procedure results in a classification of Yang-Baxter characters (Thm. 4.7), and thus also essentially answers Q3). The full answer to Q1) is then given in Sect. 4.2, where we establish a parameterization of  $\mathcal{R}_0/\sim$  by pairs of Young diagrams (Thm. 4.8).

In Sect. 5, we use  $K$ -theory to characterize the equivalence  $R \sim S$  in terms of approximate unitary equivalence of the homomorphisms  $\rho_R, \rho_S$  (Thm. 5.3) and recover a result of Kerov and Vershik [VK83] in our Yang-Baxter setting.

As  $K_0(C^*S_\infty)$  is isomorphic to a quotient of the ring of symmetric functions [VK83], this also enables us to give an explicit formula for the decomposition of the Yang-Baxter representations  $\rho_R^{(n)}$  into irreducibles in terms of symmetric functions (Prop. 5.7).

Sect. 6 is devoted to a discussion of examples, including in particular Yang-Baxter representations of the Temperley-Lieb algebra at parameter  $q = 2$  [TL71]. Our results allow us to classify such representations completely in terms of their dimension and rank (Prop. 6.3).

## 2 R-matrices and characters of $S_\infty$

The infinite symmetric group  $S_\infty$  is the group of all bijections of  $\mathbb{N}$  that move only finitely many points, a countable discrete group with infinite (non-trivial) conjugacy classes. A *character* of  $S_\infty$  is defined as a positive definite class function  $\chi : S_\infty \rightarrow \mathbb{C}$  that is *normalized* at the identity,  $\chi(e) = 1$ . For example, the trivial representation has the constant character 1. The characters of  $S_\infty$  form a simplex, the extreme points of which are called *extremal characters* (or indecomposable characters). An example of an extremal character is the Plancherel trace  $\chi(\sigma) = \delta_{\sigma,e}$ .

Thoma found the following characterization of extremality of characters of  $S_\infty$ , often called ‘‘Thoma Multiplicativity’’. In its formulation, we define the support of  $\sigma \in S_\infty$  as the complement of the fixed points of  $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ .

**Theorem 2.1. [Tho64a]** *A character  $\chi$  of  $S_\infty$  is extremal if and only if for  $\sigma, \sigma' \in S_\infty$  with disjoint supports, it holds that  $\chi(\sigma\sigma') = \chi(\sigma)\chi(\sigma')$ .*

Some elements of  $S_\infty$  will appear repeatedly. We write  $\sigma_{i,j} = (i, j)$  for two-cycles, and specifically  $\sigma_k = \sigma_{k,k+1}$  for neighboring transpositions, the standard generators of  $S_\infty$ . General  $n$ -cycles will be denoted  $c_n \in S_\infty$ . In case a specific choice of  $n$ -cycle is necessary, we choose

$$c_n = \sigma_{n-1} \cdots \sigma_2 \sigma_1 = \sigma_{1,2} \sigma_{1,3} \cdots \sigma_{1,n}, \quad (2.1a)$$

$$\sigma_{1,n} = \sigma_{n-1} \cdots \sigma_2 \sigma_1 \sigma_2 \cdots \sigma_{n-1}. \quad (2.1b)$$

In view of the above theorem and the cycle decomposition of permutations, an extremal character  $\chi$  of  $S_\infty$  is uniquely determined by its values on  $n$ -cycles. For a general group element  $\sigma \in S_\infty$ , one then has

$$\chi(\sigma) = \prod_{n \geq 2} \chi(c_n)^{k_n},$$

where  $k_n$  is the number of  $n$ -cycles in the decomposition of  $\sigma$  into disjoint cycles.

### 2.1 Extremality and Thoma’s parameterization

We now connect  $S_\infty$  to R-matrices by showing that any R-matrix defines an extremal character and corresponding factor representation of  $S_\infty$ . We will be working with the infinite tensor product  $\mathcal{E}_0 := \bigotimes_{n \geq 1} \text{End } V$  (defined only algebraically at this point), with inclusions fixed by tensoring with  $\text{id}_V$  in the last factor. With the group inclusions  $S_n \subset S_{n+1} \subset S_\infty$  defined by letting  $\sigma \in S_n$  act on  $\mathbb{N}$  by keeping all  $j > n$  fixed, the system of representations  $\rho_R^{(n)}$ ,  $R \in \mathcal{R}_0(V)$  is coherent and defines a \*-homomorphism  $\rho_R : \mathbb{C}[S_\infty] \rightarrow \mathcal{E}_0$ . The generators  $\sigma_i$ ,  $i \in \mathbb{N}$ , are mapped to

$$R_i := \rho_R(\sigma_i) = 1^{\otimes(i-1)} \otimes R \otimes 1 \otimes \dots, \quad (2.2)$$

where here and hereafter, we write 1 instead of  $\text{id}_V$  when the base space is clear from the context. Note that  $R_i$  can be viewed as an element of  $\text{End } V^{\otimes n}$  for  $n \geq i + 1$ , or of  $\mathcal{E}_0$ .

We refrain from viewing  $\rho_R$  as a representation on  $\bigotimes_{n \geq 1} V$ , as the definition of this space depends on choices. Our  $S_\infty$ -representations will be defined by composing  $\rho_R$  with the GNS representation of  $\mathcal{E}_0$  with respect to its unique normalized trace<sup>3</sup>,

$$\tau = \bigotimes_{n \geq 1} \frac{\text{Tr}_V}{d} : \mathcal{E}_0 \rightarrow \mathbb{C}. \quad (2.3)$$

**Proposition 2.2.** *Let  $R \in \mathcal{R}_0(V)$ . Then*

$$\chi_R := \tau \circ \rho_R \quad (2.4)$$

*is an extremal character of  $S_\infty$ . On an  $n$ -cycle  $c_n$ ,  $n \geq 2$ , it evaluates to*

$$\chi_R(c_n) = d^{-n} \text{Tr}_{V^{\otimes n}}(R_1 \cdots R_{n-1}), \quad d = \dim V. \quad (2.5)$$

*Proof.* By standard properties of the trace,  $\chi_R$  is a normalized positive class function. To show that  $\chi_R$  is also extremal, we have to verify that it factorizes over permutations  $\sigma, \sigma' \in S_\infty$  with disjoint supports (Thm. 2.1).

Let  $\sigma, \sigma' \in S_\infty$  have disjoint supports. Taking into account that  $\chi_R$  is a class function, we may assume without loss of generality that  $\text{supp } \sigma \subset \{1, \dots, n\}$  and  $\text{supp } \sigma' \subset \{n+1, \dots, n+m\}$  for some  $n, m \in \mathbb{N}$ .

Setting  $N := n+m$ , we then have  $\rho_R^{(N)}(\sigma) = \rho_R^{(n)}(\sigma) \otimes 1^{\otimes m}$  and  $\rho_R^{(N)}(\sigma') = 1^{\otimes n} \otimes \rho_R^{(m)}(\sigma')$ . Using  $\text{Tr}_{V \otimes W}(A \otimes B) = \text{Tr}_V(A) \text{Tr}_W(B)$ , we arrive at

$$\begin{aligned} \chi_R(\sigma\sigma') &= d^{-N} \text{Tr}_{V^{\otimes N}}((\rho_R^{(n)}(\sigma) \otimes 1^{\otimes m})(1^{\otimes n} \otimes \rho_R^{(m)}(\sigma'))) \\ &= d^{-n} \text{Tr}_{V^{\otimes n}}(\rho_R^{(n)}(\sigma)) \cdot d^{-m} \text{Tr}_{V^{\otimes m}}(\rho_R^{(m)}(\sigma')) \\ &= \chi_R(\sigma)\chi_R(\sigma'), \end{aligned}$$

and the proof of extremality of  $\chi_R$  is finished.

For the second statement, we only need to note that  $\rho_R$  (1.2) maps the  $n$ -cycle  $\sigma_1\sigma_2 \cdots \sigma_{n-1}$  to  $R_1R_2 \cdots R_{n-1} \in \text{End}(V)^{\otimes n}$ .  $\square$

We will call the characters  $\chi_R$ ,  $R \in \mathcal{R}_0$ , *Yang-Baxter characters* of  $S_\infty$ . As we just demonstrated, every Yang-Baxter character is extremal. We will see in the next section that the converse is not true: not every extremal character is Yang-Baxter.

Using the representation theory of finite groups and the inductive limit definition of  $S_\infty$ , it follows from Prop. 2.2 that two R-matrices  $R, S \in \mathcal{R}_0$  are equivalent in the sense of Def. 1.1 if and only if they have the same character and the same dimension. (As we work with normalized characters, the dimension is not contained in the character.) Thus the dimension and the sequence of traces (2.5) (indirectly) characterize the equivalence classes  $\mathcal{R}_0/\sim$ .

Thoma not only found a criterion for characterizing extremal characters, but also gave a classification in terms of an infinite dimensional simplex.

<sup>3</sup>See Sec. 5.3 for different choices of states on  $\mathcal{E}_0$ .

**Theorem 2.3. [Tho64a]** Let  $\mathbb{T}$  denote the collection of all sequences  $\{\alpha_i\}_{i \in \mathbb{N}}$ ,  $\{\beta_i\}_{i \in \mathbb{N}}$  of real numbers such that

- i)  $\alpha_i \geq 0$  and  $\beta_i \geq 0$ ,
- ii)  $\alpha_i \geq \alpha_{i+1}$  and  $\beta_i \geq \beta_{i+1}$ ,
- iii)  $\sum_i \alpha_i + \sum_j \beta_j \leq 1$ .

For each  $(\alpha, \beta) \in \mathbb{T}$ , there exists a unique extremal character  $\chi$  of  $S_\infty$ . On an  $n$ -cycle, it takes the value

$$\chi(c_n) = \sum_i \alpha_i^n + (-1)^{n+1} \sum_i \beta_i^n, \quad n \geq 2. \quad (2.6)$$

We will call the parameters  $(\alpha, \beta) \in \mathbb{T}$  the *Thoma parameters* of a character.

As a consequence of these results, any  $R \in \mathcal{R}_0$  defines a point  $(\alpha, \beta) \in \mathbb{T}$ . Questions Q1) and Q3) from the Introduction are therefore closely connected to the problem of identifying the subset of all Thoma parameters of Yang-Baxter characters inside  $\mathbb{T}$ . This task is taken up in the following sections.

Question Q2), concerned with an explicit characterization of the equivalence relation  $\sim$ , amounts to extracting the Thoma parameters  $(\alpha, \beta)$  from an involutive R-matrix. In view of (2.5) and (2.6), the parameters  $(\alpha, \beta) \in \mathbb{T}$  corresponding to  $R \in \mathcal{R}_0$  are uniquely fixed by the system of equations

$$\sum_i \alpha_i^n + (-1)^{n+1} \sum_i \beta_i^n = d^{-n} \operatorname{Tr}_{V^{\otimes n}}(R_1 \cdots R_{n-1}), \quad n \geq 2. \quad (2.7)$$

We will develop tools to compute  $(\alpha, \beta)$  directly from  $R$  in Sections 3 and 4.1.

To conclude this section, let us list the Thoma parameters of the simple R-matrices encountered so far. Recall that the flip in any dimension  $d$  is denoted by  $F$ .

$R$	non-vanishing Thoma parameters
1	$\alpha_1 = 1$ , independent of $d$
-1	$\beta_1 = 1$ , independent of $d$
$F$	$\alpha_1 = \dots = \alpha_d = d^{-1}$
$-F$	$\beta_1 = \dots = \beta_d = d^{-1}$

Since the R-matrices  $R = 1$  and  $R = -1$  obviously give the trivial and alternating representation of  $S_\infty$ , respectively, the first two lines immediately follow from (2.7). The claimed parameters of  $\pm F$  can be verified by computing  $\operatorname{Tr}_{V^{\otimes n}}(F_1 \cdots F_{n-1}) = d$ .

## 2.2 Faithfulness

Given an R-matrix  $R \in \mathcal{R}_0$  of dimension  $d$ , the homomorphism  $\rho_R$  restricts to a representation  $\rho_R^{(n)}$  of  $S_n$  on  $V^{\otimes n}$ , which has dimension  $d^n$ . This observation expresses that Yang-Baxter representations are “small”, and leads to restrictions on the Thoma parameters of Yang-Baxter characters.

**Proposition 2.4.** *Let  $R \in \mathcal{R}_0$ .*

*i) As a group homomorphism,  $\rho_R$  is injective if and only if  $R \neq \pm 1$ .*

*ii) As an algebra homomorphism,  $\rho_R : \mathbb{C}[S_\infty] \rightarrow \mathcal{E}_0$  is not injective.*

*Proof.* *i)* This is a general property of  $S_\infty$ . Clearly, if  $R = \pm 1$  then  $\rho_R$  is not injective. Conversely, assume that  $\rho_R$  is not injective and  $\sigma \in S_\infty$  lies in the kernel, then  $\sigma$  also lies in the kernel of  $\rho|_{S_n}$  for  $n$  sufficiently large. But for  $n \geq 5$ , the only non-trivial proper normal subgroup of  $S_n$  is the alternating group  $A_n$ . Thus  $\ker \rho|_{S_n}$  contains at least  $A_n$ . This implies that the image of  $\rho$  is either trivial or  $\mathbb{Z}_2$ . In the case at hand, this means that  $\rho_R$  is injective if and only if  $R \neq \pm 1$ .

*ii)*  $\rho_R$  restricts to an algebra homomorphism  $\rho_R^{(n)} : \mathbb{C}[S_n] \rightarrow \text{End } V^{\otimes n}$ . As the dimensions of  $\mathbb{C}[S_n]$  and  $\text{End}(V^{\otimes n})$  are  $n!$  and  $d^{2n}$ , respectively, and  $n! > d^{2n}$  for  $n$  sufficiently large, it follows that  $\rho_R^{(n)}$  cannot be injective.  $\square$

The second part of this proposition implies that Yang-Baxter characters are never faithful. This observation allows us to make use of the following theorem due to Wassermann [Was81, Thm. III.6.5].

**Theorem 2.5. [Was81]** *Let  $\chi$  be an extremal character of  $S_\infty$  with Thoma parameters  $(\alpha, \beta) \in \mathbb{T}$ . Then  $\chi$  is faithful as a state of the group  $C^*$ -algebra  $C^*S_\infty$  if and only if either  $\sum_i \alpha_i + \sum_i \beta_i < 1$ , or  $\sum_i \alpha_i + \sum_i \beta_i = 1$  and infinitely many  $\alpha_i$  or  $\beta_i$  are non-zero.*

In combination with Prop. 2.4 *ii)*, this immediately implies the following result.

**Corollary 2.6.** *Let  $(\alpha, \beta) \in \mathbb{T}$  be the Thoma parameters of a Yang-Baxter character  $\chi_R$ . Then  $\sum_i \alpha_i + \sum_i \beta_i = 1$ , and only finitely many  $\alpha_i$  or  $\beta_i$  are non-zero.*

We can now give a first example of an extremal non-Yang-Baxter character, namely the Plancherel trace  $\chi(\sigma) = \delta_{\sigma, e}$ . By (2.6), the Plancherel trace has Thoma parameters  $\alpha = \beta = 0$  and therefore violates the condition  $\sum_i \alpha_i + \sum_i \beta_i = 1$ . Its GNS representation is the left regular representation, which is “too large” to be of Yang-Baxter form.

### 3 Yang-Baxter subfactors

The Thoma parameters of a Yang-Baxter character have further properties, in addition to the ones spelled out in Cor. 2.6. To extract these properties, and to derive a characterization of the equivalence relation  $\sim$ , we now switch to a setting involving von Neumann algebras. Specifically, we will consider subfactors [Jon83, JS97] arising from the subgroup

$$S_\infty^> \subset S_\infty, \quad S_\infty^> := \{\sigma \in S_\infty : \sigma(1) = 1\}. \quad (3.1)$$



Given an extremal character  $\chi$  of  $S_\infty$ , we may view it as a tracial state on the group  $C^*$ -algebra  $C^*S_\infty$  (we denote the state and the character by the same symbol).

The GNS data of  $(C^*S_\infty, \chi)$  will be denoted  $(\mathcal{H}_\chi, \Omega_\chi, \pi_\chi)$ , and the von Neumann algebra generated by the representation  $\mathcal{M}_\chi := \pi_\chi(C^*S_\infty)''$ . Since  $\chi$  is extremal,  $\mathcal{M}_\chi$  is a (finite) factor — it is trivial for the one-dimensional trivial and alternating representations, and hyperfinite of type  $\text{II}_1$  in all other cases.

In our situation of Yang-Baxter representations, we have the homomorphism  $\rho_R : \mathbb{C}[S_\infty] \rightarrow \mathcal{E}_0 = \bigcup_n \text{End } V^{\otimes n}$ . Proceeding to the GNS representation  $\pi_\tau$  of  $\mathcal{E}_0$  with respect to the trace  $\tau$ , we may weakly close  $\mathcal{E}_0$  to  $\mathcal{E}$  (a hyperfinite  $\text{II}_1$  factor), and obtain the subfactor

$$\mathcal{M}_R := \rho_R(\mathbb{C}[S_\infty])'' \subset \mathcal{E}. \quad (3.2)$$

Since  $\pi_\tau$  is faithful (in contrast to  $\rho_R$  and  $\pi_{\chi_R}$ , see Prop. 2.4), we suppress it in our notation and often write  $\rho_R$  instead of  $\pi_\tau \circ \rho_R$ . We can canonically identify  $\pi_\tau \circ \rho_R = \pi_{\chi_R}$ ,  $\Omega_\tau = \Omega_{\chi_R}$ ,  $\mathcal{M}_R = \mathcal{M}_{\chi_R}$ ,  $\mathcal{H}_{\chi_R} = \overline{\mathcal{M}_R \Omega_\tau}$ .

As an aside, let us mention that our equivalence relation  $R \sim S$  implies the unitary equivalence of the representations

$$R \sim S \implies \pi_\tau \circ \rho_R \cong \pi_\tau \circ \rho_S. \quad (3.3)$$

In fact,  $R \sim S$  implies  $\chi_R = \chi_S$  and hence  $\pi_{\chi_R} = \pi_{\chi_S}$  — since  $\pi_{\chi_R}$  can be identified with the restriction of  $\pi_\tau \circ \rho_R$  to  $\mathcal{H}_{\chi_R}$ , (3.3) follows.

The subgroup (3.1) generates the von Neumann algebra

$$\mathcal{N}_R := \rho_R(C^*S_\infty^>)'' \subset \mathcal{M}_R. \quad (3.4)$$

As  $S_\infty^> \cong S_\infty$ , this is a ( $\text{I}_1$  or  $\text{II}_1$ ) subfactor.

Gohm and Köstler [GK10] and Yamashita [Yam12] have independently analyzed the subfactor  $\mathcal{N}_\chi \subset \mathcal{M}_\chi$  in the setting of general (not necessarily Yang-Baxter) extremal characters. They found that it is irreducible if and only if the parameters  $(\alpha, \beta)$  have one of the following values:

- i)  $\alpha_1 = \dots = \alpha_d = d^{-1}$  for some  $d \in \mathbb{N}$ ,
- ii)  $\beta_1 = \dots = \beta_d = d^{-1}$  for some  $d \in \mathbb{N}$ ,
- iii)  $\alpha_i = 0$  and  $\beta_i = 0$  for all  $i$ .

By comparison with our examples of R-matrices at the end of the preceding section, we see that the relative commutant  $\mathcal{N}'_R \cap \mathcal{M}_R$  is trivial if and only if  $R$  is equivalent to one of the four R-matrices  $1, -1, F, -F$ , of arbitrary dimension  $d \in \mathbb{N}$ . As we pointed out earlier, the last possibility *iii*) is realized by the Plancherel trace, which is not Yang-Baxter.

To extract information about  $R$  from the subfactor (3.4), we consider the unique  $\tau$ -preserving conditional expectation onto its relative commutant,

$$E_R : \mathcal{M}_R \rightarrow \mathcal{N}'_R \cap \mathcal{M}_R. \quad (3.5)$$

The inclusion  $\mathcal{N}_R \subset \mathcal{M}_R$  is replicated on the level of the infinite tensor product  $\mathcal{E}$ : Here we consider the inclusion  $\mathbb{C} \otimes \text{End } V \otimes \text{End } V \otimes \cdots \subset \mathcal{E}$ , the relative commutant of which is  $\text{End } V$ , viewed as a subalgebra of  $\mathcal{E}$  via the embedding  $X \mapsto X \otimes 1 \otimes 1 \cdots$ . The corresponding  $\tau$ -preserving conditional expectation is

$$E : \mathcal{E} \rightarrow \text{End } V, \quad E = \text{id}_{\text{End } V} \otimes \tau \otimes \tau \otimes \cdots. \quad (3.6)$$

In the following arguments, it turns out to be better to use a twisted version of  $E$ , namely the *left* partial trace,

$$E_l := (\tau \otimes \text{id}_{\text{End } V}) \otimes \tau \otimes \tau \otimes \cdots : \mathcal{E} \rightarrow \text{End } V \otimes \mathbb{C} \otimes \mathbb{C} \otimes \cdots. \quad (3.7)$$

instead of the right partial trace (3.6). A posteriori, we will be able to conclude  $E_l(R_1) \cong E(R_1)$ , but this is not clear from the outset.

We want to show that the diagram

$$\begin{array}{ccc} \text{End } V & \xleftarrow{E_l} & \mathcal{E} \\ \uparrow & & \uparrow \\ \mathcal{N}'_R \cap \mathcal{M}_R & \xleftarrow{E_R} & \mathcal{M}_R \end{array} \quad (3.8)$$

is a commuting square. A priori, it is in particular not obvious that we have the inclusion  $\mathcal{N}'_R \cap \mathcal{M}_R \subset \text{End } V$  on the left hand side, but this will be shown below.

The key step is to show that  $E_R$  and  $E_l$  agree on  $R_1 = \rho_R(\sigma_1) \in \mathcal{M}_R$ . Following [Yam12], we consider the subgroups  $T_n = \{\sigma \in S_{n+1} : \sigma(1) = 1\} \subset S_\infty$  and the von Neumann algebras generated by them,  $\mathcal{N}_{R,n} := \rho_R(T_n)'' \subset \mathcal{M}_R$ . As  $T_n \subset T_{n+1}$ , this yields a descending chain of relative commutants,  $n \in \mathbb{N}$ ,

$$\mathcal{M}_R \supset (\mathcal{N}'_{R,n} \cap \mathcal{M}_R) \supset (\mathcal{N}'_{R,n+1} \cap \mathcal{M}_R) \supset (\mathcal{N}'_R \cap \mathcal{M}_R),$$

with corresponding conditional expectations  $E_{R,n} : \mathcal{M}_R \rightarrow \mathcal{N}'_{R,n} \cap \mathcal{M}_R$ . Since  $T_n$  is finite,  $E_{R,n}$  is simply given by averaging,

$$E_{R,n}(M) = \frac{1}{n!} \sum_{\sigma \in T_n} \rho_R(\sigma) M \rho_R(\sigma^{-1}), \quad M \in \mathcal{M}_R. \quad (3.9)$$

It is not hard to compute that for  $M = R_1$ , one gets [Yam12]

$$E_{R,n}(R_1) = \frac{1}{n} \sum_{j=2}^{n+1} \rho_R(\sigma_{1,j}). \quad (3.10)$$

**Lemma 3.1.**  $E_R(R_1) = E_l(R_1)$ .

*Proof.* The basic idea of the proof is to use the fact that  $E_{R,n} \rightarrow E_R$  as  $n \rightarrow \infty$  in the 2-norm given by  $\tau$ . That is, we need to show that (note that  $R_1 = R_1^*$ )

$$\begin{aligned} \tau(|E_{R,n}(R_1) - E_l(R_1)|^2) &= \tau(E_{R,n}(R_1)E_{R,n}(R_1)) \\ &\quad - 2\tau(E_{R,n}(R_1)E_l(R_1)) + \tau(E_l(R_1)^2) \end{aligned} \quad (3.11)$$

converges to zero as  $n \rightarrow \infty$ . The argument is thus similar to the one in [Yam12], but involves some extra twists in our setting.

In a first step, we claim that for any  $X \in \rho_R(\mathbb{C}[S_\infty])$ , we have

$$\lim_{n \rightarrow \infty} \tau(X E_{R,n}(R_1)) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=2}^{n+1} \tau\left(X \rho_R(\sigma_{1,j})\right) = \tau\left(X E_l(R_1)\right). \quad (3.12)$$

It is sufficient to show the claim for  $X = \rho_R(\sigma)$  with  $\sigma \in S_k$ ,  $k \in \mathbb{N}$ .

As  $\sigma_{1,j} = \sigma_{j-1} \cdots \sigma_2 \sigma_1 \sigma_2 \cdots \sigma_{j-1}$  (2.1b), it follows that  $\chi_R(\sigma \sigma_{1,j}) = \chi_R(\sigma \sigma_{1,k+1})$  for all  $j > k$ . With this observation, we have

$$\frac{1}{n} \sum_{j=2}^{n+1} \tau\left(\rho_R(\sigma) \rho_R(\sigma_{1,j})\right) = \frac{1}{n} \sum_{j=2}^k \chi_R(\sigma \cdot \sigma_{1,j}) + \frac{1}{n} \sum_{j=k+1}^{n+1} \chi_R(\sigma \cdot \sigma_{1,k+1}).$$

As  $n \rightarrow \infty$ , the first term vanishes and the second converges to  $\chi_R(\sigma \sigma_{1,k+1})$ . To compute this character value, we compare the cycle structures of  $\sigma$  and  $\sigma \cdot \sigma_{1,k+1}$ : Any cycle of  $\sigma$  that does not involve 1 also occurs in  $\sigma \cdot (1, k+1)$ . If the cycle involving 1 has length  $m$ , then this cycle is changed to a cycle of length  $m+1$  in  $\sigma \cdot \sigma_{1,k+1}$ . Thus  $\chi_R(\sigma \cdot \sigma_{1,k+1}) = \chi_R(\sigma) \cdot \chi_R(c_{m+1}) / \chi_R(c_m)$ .

By definition of  $E_l$  (here the difference to  $E$  enters), the right hand side of (3.12) is

$$\tau\left(\rho_R(\sigma) E_l(R_1)\right) = \tau\left((1 \otimes \rho_R(\sigma)) R_1\right) = \chi_R(\sigma' \cdot \sigma_1),$$

where  $\sigma' \in S_{k+1}$  is defined by  $\sigma'(1) = 1$ ,  $\sigma'(i) = \sigma(i-1)$ ,  $i > 1$ . Again we compare the cycle structures of  $\sigma$  and  $\sigma' \sigma_1$ : Any cycle not involving 1 in  $\sigma$  appears as a cycle not involving 2 in  $\sigma'$ . The cycle including 1 in  $\sigma$ , say of length  $m$ , yields an  $(m+1)$ -cycle in  $\sigma' \sigma_1$ . Thus  $\chi_R(\sigma' \cdot \sigma_1) = \chi_R(\sigma) \cdot \chi_R(c_{m+1}) / \chi_R(c_m)$ , in agreement with the previously computed limit, proving (3.12).

This result implies that  $E_l(R_1)$  is an element of  $\mathcal{M}_R$ . In fact, by the traciality of  $\tau$ , we have shown that for any  $X, Y \in \rho_R(\mathbb{C}[S_\infty])$ , we have the limit  $\tau(X E_{R,n}(R_1) Y) \rightarrow \tau(X E_l(R_1) Y)$ . As  $\rho_R(\mathbb{C}[S_\infty]) \Omega_\tau \subset \mathcal{H}_{\chi_R}$  is dense, and  $\|E_{R,n}(R_1)\|$  is uniformly bounded in  $n$ , it follows that  $E_{R,n}(R_1)$  converges weakly to  $E_l(R_1) \in \mathcal{M}_R$ . In particular,

$$\lim_{n \rightarrow \infty} \tau(E_{R,n}(R_1) E_l(R_1)) = \tau(E_l(R_1)^2).$$

Inserted into (3.11), we see that it remains to show  $t_{n,n} \rightarrow t_\infty := \tau(E_l(R_1)^2)$  as  $n \rightarrow \infty$ , where we have introduced  $t_{n,m} := \tau(E_{R,n}(R_1) E_{R,m}(R_1))$ .

To this end, note that the weak limit  $E_{R,n}(R_1) \rightarrow E_l(R_1)$  implies

$$\lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} t_{n,m} = t_\infty.$$

For showing that this coincides with the desired diagonal limit  $\lim_n t_{n,n}$ , we recall that for  $n \geq m$ , the conditional expectation  $E_{R,n}$  projects onto a smaller algebra than  $E_{R,m}$ , so that we have  $E_{R,n} \circ E_{R,m} = E_{R,n}$ . Hence, for  $n \geq m$ ,

$$t_{n,m} = \tau(E_{R,n}(E_{R,n}(R_1) E_{R,m}(R_1))) = \tau(E_{R,n}(R_1) E_{R,n}(E_{R,m}(R_1))) = t_{n,n},$$

i.e., we have  $t_{n,m} = t_{\max\{n,m\}}$ . This implies  $\lim_n t_{n,n} = \lim_n \lim_m t_{n,m}$  and concludes the proof.  $\square$

With  $E_l(R_1)$ , we now have concrete elements of the relative commutant  $\mathcal{N}'_R \cap \mathcal{M}_R$  at our disposal. For  $R = \pm 1$  or  $R = \pm F$ , these partial traces are trivial,  $E_l(\pm 1) = \pm d \text{id}_V$ ,  $E_l(\pm F) = \pm \text{id}_V$ , as can be computed directly or inferred from the above mentioned result on irreducibility of  $\mathcal{N}_R \subset \mathcal{M}_R$ .

However, for all R-matrices not equivalent to  $\pm 1, \pm F$ , we get non-trivial partial traces  $E_l(R_1)$ . In fact, it was shown in [GK11, Yam12] that  $E_R(R_1)$  generates the relative commutant  $\mathcal{N}'_R \cap \mathcal{M}_R$ . This implies that for  $R \not\sim \pm 1, \pm F$ , the expectation  $E_l(R_1)$  is not a multiple of the identity. Additionally, we can conclude that (3.8) is indeed a commuting square.

The partial trace  $E_l(R_1)$  of the R-matrix turns out to be a complete invariant for the equivalence relation  $\sim$ . This is a consequence of the next theorem, which follows from the work of Gohm and Köstler, and our Lemma 3.1. These authors prove it in a setting of noncommutative probability [GK10], building on their earlier work [GK09, Kös10] (see also [GK11]). In our situation, only certain aspects of [GK09, GK10, Kös10, GK11] are needed, and we give a shortened proof for the sake of self-containedness.

This proof makes use of the so-called partial shifts, defined as

$$\gamma_m(M) = \lim_{n \rightarrow \infty} R_{m+1} R_{m+2} \cdots R_n \cdot M \cdot R_n \cdots R_{m+2} R_{m+1}, \quad m \in \mathbb{N}. \quad (3.13)$$

These limits exist in the strong operator topology for any  $M \in \mathcal{M}_R$  [GK10, Prop. 2.13], and define  $\tau$ -preserving endomorphisms of  $\mathcal{M}_R$ . Clearly  $\gamma_m$  acts trivially on  $\mathcal{N}'_R \cap \mathcal{M}_R$ . Thus the conditional expectation onto the relative commutant is invariant,  $E_R \circ \gamma_m = E_R$ .

By explicit calculation based on (2.1b) and the Yang-Baxter equation, one shows that [GK10, Prop. 3.3]

$$\gamma_m(\rho_R(\sigma_{1,n})) = \begin{cases} \rho_R(\sigma_{1,n}) & n < m+1 \\ \rho_R(\sigma_{1,n+1}) & n \geq m+1 \end{cases}. \quad (3.14)$$

and in particular,

$$\gamma_1^p(R_1) = \rho_R(\sigma_{1,p+2}), \quad p \in \mathbb{N}. \quad (3.15)$$

As mentioned before, the relative commutant  $\mathcal{N}'_R \cap \mathcal{M}_R$  is contained in the fixed point algebra of  $\gamma_1$ . Gohm and Köstler proved that in fact, equality holds:  $\mathcal{M}_R^{\gamma_1} = \mathcal{N}'_R \cap \mathcal{M}_R$  [GK10, Thm. 3.6 (iii)].

**Proposition 3.2.** *Let  $c_n \in S_\infty$  be an  $n$ -cycle,  $n \geq 2$ . Then*

$$\chi_R(c_n) = \tau(E_l(R_1)^{n-1}). \quad (3.16)$$

*Proof.* For  $n = 2$ , the statement is a direct consequence of the definition of  $E_l$ . For the induction step, we consider the specific cycle  $c_{n+1} = c_n \sigma_{1,n+1}$  (2.1a). Writing  $C_n = \rho_R(c_n)$  as a shorthand, we note that  $\gamma_n(C_n) = C_n$  for (see (2.1a) and (3.14)). As  $E_R$  is invariant under  $\gamma_n$ , we obtain

$$E_R(C_{n+1}) = E_R(\gamma_n(C_n \rho_R(\sigma_{1,n+1}))) = E_R(C_n \cdot \rho_R(\sigma_{1,n+2})).$$

In the same manner, we can now insert the endomorphism  $\gamma_{n+1}$ , which also leaves  $C_n$  invariant, and maps  $\rho_R(\sigma_{1,n+2})$  to  $\rho_R(\sigma_{1,n+3})$ . Iteratively, this gives  $E_R(C_{n+1}) = E_R(C_n \rho_R(\sigma_{1,n+p})) = E_R(C_n \gamma_1^{n+p-2}(R_1))$ ,  $p \in \mathbb{N}$ .

Averaging over  $p$  yields for any  $N \in \mathbb{N}$

$$E_R(C_{n+1}) = E_R \left( C_n \cdot \frac{1}{N} \sum_{p=1}^N \gamma_1^p(\gamma_1^{n-2}(R_1)) \right).$$

We may now use the ergodic theorem [Pet83], stating here that for any  $M \in \mathcal{M}_R$ , the ergodic averages  $N^{-1} \sum_{p=1}^N \gamma_1^p(M)$  converge strongly to the conditional expectation  $E_R(M)$  onto the fixed point algebra  $\mathcal{M}_R^{\gamma_1} = \mathcal{N}'_R \cap \mathcal{M}_R$  as  $N \rightarrow \infty$  [Kös10, Thm.8.3]. As  $\gamma_1^{n-2}(R_1) \in \mathcal{M}_R$ , and  $E_R$  is continuous in the strong operator topology, we have  $E_R(C_{n+1}) = E_R(C_n \cdot E_R(\gamma_1^{n-2}(R_1))) = E_R(C_n) \cdot E_R(\gamma_1^{n-2}(R_1))$ . In view of the  $\gamma_1$ -invariance of  $E_R$  and Lemma 3.1, the last term simplifies to  $E_R(\gamma_1^{n-2}(R_1)) = E_R(R_1) = E_l(R_1)$ .

We thus have shown  $E_R(C_{n+1}) = E_R(C_n) \cdot E_l(R_1)$ , which implies  $E_R(C_n) = E_l(R_1)^{n-1}$  by induction. Evaluating in  $\tau$  then gives the claimed result.  $\square$

We have now extracted sufficient information from the subfactor setting, and return to our analysis of equivalence of R-matrices. At this point, it is better to switch to the usual partial trace of  $R$ , defined as

$$\text{ptr } R = \dim V \cdot E_l(R_1) = (\text{Tr}_V \otimes \text{id}_{\text{End } V})(R) \quad (3.17)$$

and viewed as an element of  $\text{End } V$  rather than  $\mathcal{E}$ . The key relation (3.16) can then be rewritten as

$$\chi_R(c_n) = d^{-n} \text{Tr}_V(\text{ptr}(R)^{n-1}). \quad (3.18)$$

**Theorem 3.3.** *Two R-matrices  $R, S \in \mathcal{R}_0$  are equivalent if and only if they have similar partial traces,  $\text{ptr } R \cong \text{ptr } S$ .*

*Proof.* If  $R \in \mathcal{R}_0(V)$  and  $S \in \mathcal{R}_0(W)$  have similar partial traces, then clearly  $\text{Tr}_V(\text{ptr}(R)^{n-1}) = \text{Tr}_W(\text{ptr}(S)^{n-1})$ . As similarity of the partial traces implies in particular that the dimensions  $\dim V = \dim W$  coincide, we conclude  $\chi_R = \chi_S$  from (3.18) and Thoma multiplicativity. Thus,  $\text{ptr}(R) \cong \text{ptr}(S) \Rightarrow R \sim S$ .

Conversely, if  $R \sim S$ , then these R-matrices have the same dimension and character, and hence  $\text{Tr}_V(\text{ptr}(R)^{n-1}) = \text{Tr}_W(\text{ptr}(S)^{n-1})$ ,  $n \geq 2$ , from (3.18). This implies that the selfadjoint endomorphisms  $\text{ptr}(R)$ ,  $\text{ptr}(S)$  have the same characteristic polynomial, and are therefore similar.  $\square$

As an immediate application, let us return to our earlier discussion of left and right partial traces. Since  $FRF \sim R$  for any  $R \in \mathcal{R}_0$ , Thm. 3.3 implies that  $\text{ptr}(R) \cong \text{ptr}(FRF)$ . But  $\text{ptr}(FRF)$  coincides with the *right* partial trace of  $R$ , namely  $(\text{id}_{\text{End } V} \otimes \text{Tr}_V)(R)$ . Thus, a posteriori, we also have  $E(R_1) \cong E_R(R_1)$ , where  $E$  is the conditional expectation (3.6).

Thm. 3.3 shows that the eigenvalues of  $\text{ptr}(R)$  (and their multiplicities) characterize the equivalence classes  $\mathcal{R}_0/\sim$ . Such spectral characterizations also

appear in the work of Okounkov on Thoma measures and Olshanski pairs [Oko95]. In our Yang-Baxter setting, the spectrum of  $\text{ptr}(R)$  has a very specific form, which will be the key to our classification of R-matrices in the next section.

As a second important consequence of Prop. 3.2, next we demonstrate that Yang-Baxter characters have *rational* Thoma parameters after stating a preparatory lemma.

**Lemma 3.4.** *Let  $\{x_i\}_i$  and  $\{y_j\}_j$  be two finite sequences of positive real numbers such that for all  $n \in \mathbb{N}$ ,*

$$\sum_i x_i^{2n+1} = \sum_j y_j^n. \quad (3.19)$$

*Then the  $x_i, y_j$  are rational.*

*Proof.* We order the sequences  $\{x_i\}_i, \{y_j\}_j$  non-increasingly and define  $\mu \in \mathbb{N}$  as the multiplicity of the maximal value of the first sequence, i.e.  $x_1 = \dots = x_\mu > x_{\mu+1}$ . Dividing (3.19) by  $x_1^{2n+1}$  yields

$$\sum_i \left(\frac{x_i}{x_1}\right)^{2n+1} = \frac{1}{x_1} \sum_j \left(\frac{y_j}{x_1^2}\right)^n.$$

In the limit  $n \rightarrow \infty$ , the left hand side converges to  $\mu$ . In this limit, the right hand side goes to infinity if  $y_1 > x_1^2$  and to 0 if  $y_1 < x_1^2$ . As  $0 < \mu < \infty$ , we conclude that  $y_1 = x_1^2$ , and define  $\nu \in \mathbb{N}$  as its multiplicity,  $y_1 = \dots = y_\nu > y_{\nu+1}$ . Then the right hand side has the limit  $\frac{\nu}{x_1}$  as  $n \rightarrow \infty$ , so that  $x_1 = \frac{\nu}{\mu}$  and  $y_1 = x_1^2$  are rational.

Inserting these values of  $x_1$  and  $y_1$  into (3.19), we find

$$\mu \left(\frac{\nu}{\mu}\right)^{2n+1} + \sum_{i>\mu} x_i^{2n+1} = \nu \left(\frac{\nu}{\mu}\right)^{2n} + \sum_{j>\nu} y_j^n,$$

and hence (3.19) also holds for the shorter sequences  $\{x_i\}_{i>\mu}$  and  $\{y_j\}_{j>\nu}$ . The claim now follows by induction.  $\square$

**Definition 3.5.**  $\mathbb{T}_{\text{YB}} \subset \mathbb{T}$  is defined as the subset of all  $(\alpha, \beta) \in \mathbb{T}$  satisfying:

- i) Only finitely many parameters  $\alpha_i, \beta_j$  are non-zero.
- ii)  $\sum_i \alpha_i + \sum_j \beta_j = 1$ .
- iii) All  $\alpha_i, \beta_j$  are rational.

**Theorem 3.6.** *The Thoma parameters of any Yang-Baxter character lie in  $\mathbb{T}_{\text{YB}}$ .*

*Proof.* Due to Cor. 2.6, the only property of Definition 3.5 that remains to be shown is *iii*). To do so, we express the character on the left hand side of (3.18) in terms of its Thoma parameters  $(\alpha, \beta)$  (2.6), and the traces on the right hand

side of (3.18) in terms of the non-zero eigenvalues  $t_j$  of  $\text{ptr } R$  (note that  $\text{ptr } R$  is selfadjoint, so the  $t_j$  are real). This yields

$$\sum_i \alpha_i^n + (-1)^{n+1} \sum_i \beta_i^n = d^{-n} \sum_j t_j^{n-1}.$$

Specializing to the case that  $n = 2m + 1$  is odd, we are in the situation of the preceding lemma with  $\{x_i\}_i = \{d\alpha_i, d\beta_i\}_i$  and  $y_j = t_j^2$ .  $\square$

Not only the Thoma parameters, but also the eigenvalues of  $\text{ptr}(R)$  are rational. In fact, we will see that each eigenvalue of  $\text{ptr } R$  is a non-zero integer, with multiplicities following a specific pattern. These facts will be discussed in the next section.

## 4 The structure of $\mathcal{R}_0/\sim$

### 4.1 Normal forms of involutive R-matrices

Our next aim is to prove that  $\mathbb{T}_{\text{YB}}$  parameterizes the set of all Yang-Baxter characters, that is, that every  $(\alpha, \beta) \in \mathbb{T}_{\text{YB}}$  is realized as the Thoma parameters of some R-matrix.

We will follow a procedure which has some analogy to building general group representations (of, say, a finite group) as direct sums of irreducibles. Yang-Baxter representations are reducible, but decomposing them gives representations which are no longer of Yang-Baxter form. Conversely, taking direct sums of Yang-Baxter representations is not compatible with the Yang-Baxter equation either.

To get around these problems, we introduce a binary operation  $\boxplus$  on R-matrices that on the level of the base spaces corresponds to taking direct sums, and respects the Yang-Baxter equation. Under various names, such operations have been considered in the literature before [Lyu87, Gur91, Hie93]. We present here the version that is most useful for the case at hand.

**Definition 4.1.** *Let  $V, W$  be finite dimensional vector spaces and let  $X \in \text{End}(V \otimes V)$ ,  $Y \in \text{End}(W \otimes W)$ . We define  $X \boxplus Y \in \text{End}((V \oplus W) \otimes (V \oplus W))$  as*

$$\begin{aligned} X \boxplus Y &= X \oplus Y \oplus F \quad \text{on} & (4.1) \\ (V \oplus W) \otimes (V \oplus W) &= (V \otimes V) \oplus (W \otimes W) \oplus ((V \otimes W) \oplus (W \otimes V)). \end{aligned}$$

In other words,  $X \boxplus Y$  acts as  $X$  on  $V \otimes V$ , as  $Y$  on  $W \otimes W$ , and as the flip on the ‘‘mixed tensors’’ involving factors from both,  $V$  and  $W$ . Note that the above definition works in the same way for infinite dimensional Hilbert spaces.

Before applying this operation to R-matrices, we collect its main properties. In particular, we note that  $\boxplus$  behaves well under taking the partial trace

$$\text{ptr} : \text{End}(U \otimes U) \rightarrow \text{End } U, \quad (4.2)$$

$$X \mapsto (\text{Tr}_U \otimes \text{id}_{\text{End } U})(X), \quad (4.3)$$

where  $U$  is any finite dimensional vector space.

**Lemma 4.2.** *Let  $V, W$  be finite dimensional vector spaces and  $X \in \text{End}(V \otimes V)$ ,  $Y \in \text{End}(W \otimes W)$ .*

- i)  $\boxplus$  is commutative and associative up to canonical isomorphism.*
- ii) If  $X$  and  $Y$  are unitary (respectively selfadjoint, involutive, invertible), then  $X \boxplus Y$  is unitary (respectively selfadjoint, involutive, invertible).*
- iii) If  $X$  commutes with the flip (on  $V \otimes V$ ) and  $Y$  commutes with the flip (on  $W \otimes W$ ), then  $X \boxplus Y$  commutes with the flip (on  $(V \oplus W) \otimes (V \oplus W)$ ).*
- iv)  $\text{ptr}(X \boxplus Y) = (\text{ptr } X) \oplus (\text{ptr } Y)$ . In particular,  $\text{Tr}(X \boxplus Y) = \text{Tr } X + \text{Tr } Y$ . The same formula holds for the right partial trace.*

*Proof.* *i)* The definition (4.1) is invariant under exchanging  $(X, V)$  with  $(Y, W)$ , that is  $X \boxplus Y = Y \boxplus X$ . Associativity follows by repeatedly evaluating the definition. Given finite dimensional vector spaces  $V^1, \dots, V^n$  and  $X^i \in \text{End}(V^i \otimes V^i)$ ,  $i = 1, \dots, n$ , one finds

$$\bigoplus_{i=1}^n X^i = X^1 \oplus \dots \oplus X^n \oplus F, \quad (4.4)$$

where on the right hand side, each  $X^i$  acts on  $V^i \otimes V^i$ , and  $F$  on the orthogonal complement of  $\bigoplus_i (V^i \otimes V^i)$  in  $(\bigoplus_i V^i)^{\otimes 2}$ .

*ii), iii)* These statements follow directly from the facts that  $F$  is unitary, selfadjoint, involutive, invertible, and the flip of  $(V \oplus W) \otimes (V \oplus W)$  leaves the three subspaces in the decomposition (4.1) invariant.

*iv)* Proving the claimed formula amounts to showing that the partial trace of  $FQ$  vanishes, where  $Q$  is the orthogonal projection onto  $(V \otimes W) \oplus (W \otimes V)$ . Let  $v_1, v_2 \in V$ , and let  $\{w_k\}$  be an orthonormal basis of  $W$ . Then the left partial trace satisfies  $\langle v_1, \text{ptr}(FQ)v_2 \rangle = \sum_k \langle w_k \otimes v_1, F(w_k \otimes v_2) \rangle$ , because  $Q$  vanishes on  $V \otimes V$ . But  $\langle w_k \otimes v_1, F(w_k \otimes v_2) \rangle = 0$  because  $V$  and  $W$  lie orthogonal to each other. The argument for the right partial trace is the same.  $\square$

We now apply  $\boxplus$  to R-matrices. The following result is known [Lyu87, Gur91, Hie93]. But since no proof seems to be available in the literature, we state it here with a proof.

**Proposition 4.3.** *Let  $R, \tilde{R} \in \mathcal{R}$ . Then  $R \boxplus \tilde{R} \in \mathcal{R}$ .*

*Proof.* Invertibility of  $R \boxplus \tilde{R}$  follows from the preceding lemma and the invertibility of  $R, \tilde{R}$ . The main point is to check that  $\hat{R} := R \boxplus \tilde{R}$  solves the Yang-Baxter equation on the third tensor power of  $V \oplus \tilde{V}$ , where  $V, \tilde{V}$  stand for the base spaces of  $R, \tilde{R}$ .

This space is the direct sum of eight orthogonal subspaces  $V_1 \otimes V_2 \otimes V_3$ , where each  $V_i$  is either  $V$  or  $\tilde{V}$ . By definition of  $\hat{R}$ , this operator leaves the subspaces  $V \otimes V$  and  $\tilde{V} \otimes \tilde{V}$  invariant, and exchanges  $V \otimes \tilde{V}$  with  $\tilde{V} \otimes V$ . This implies that  $\hat{R}_1 \hat{R}_2 \hat{R}_1$  and  $\hat{R}_2 \hat{R}_1 \hat{R}_2$  both decompose into the direct sum of their restrictions to the six subspaces  $V^{\otimes 3}$ ,  $V \otimes \tilde{V} \otimes V$ ,  $(V \otimes V \otimes \tilde{V}) \oplus (\tilde{V} \otimes V \otimes V)$ ,  $\tilde{V}^{\otimes 3}$ ,  $\tilde{V} \otimes V \otimes \tilde{V}$ , and  $(\tilde{V} \otimes \tilde{V} \otimes V) \oplus (V \otimes \tilde{V} \otimes \tilde{V})$ .



We have to show that  $\hat{R}_1\hat{R}_2\hat{R}_1$  and  $\hat{R}_2\hat{R}_1\hat{R}_2$  coincide on each of these subspaces, and by symmetry in  $V, \tilde{V}$  and  $R, \tilde{R}$ , it suffices to do so for the first three subspaces in the list. On  $V^{\otimes 3}$ ,  $\hat{R}_i\hat{R}_j\hat{R}_i$  acts as  $R_iR_jR_i$ ,  $i = 1, 2$ , and since  $R$  solves the Yang-Baxter equation, we immediately get  $\hat{R}_1\hat{R}_2\hat{R}_1|_{V^{\otimes 3}} = \hat{R}_2\hat{R}_1\hat{R}_2|_{V^{\otimes 3}}$ .

Inserting the definition of  $\hat{R}$ , one finds that on  $V \otimes \tilde{V} \otimes V$ ,  $\hat{R}_1\hat{R}_2\hat{R}_1$  acts as  $F_1R_2F_1$ , while  $\hat{R}_2\hat{R}_1\hat{R}_2$  acts as  $F_2R_1F_2$ . But these two operators coincide, as they are both equal to  $R_{1,3}$ , defined as acting as  $R$  on the outer two tensor factors, and trivially on the middle factor.

Similarly, for the last remaining subspace  $W := (V \otimes V \otimes \tilde{V}) \oplus (\tilde{V} \otimes V \otimes V)$  one finds  $\hat{R}_1\hat{R}_2\hat{R}_1|_W = (F_1F_2R_1 \oplus R_1F_2F_1)|_W$  and  $\hat{R}_2\hat{R}_1\hat{R}_2|_W = (R_2F_1F_2 \oplus F_2F_1R_2)|_W$ . But  $F_1F_2R_1 = R_2F_1F_2$  and  $F_2F_1R_2 = R_1F_2F_1$ , which finishes the proof.  $\square$

By Lemma 4.2 *ii*),  $\boxplus$  preserves involutivity, and thus also induces a binary operation on  $\mathcal{R}_0 \subset \mathcal{R}$ .

It is clear that variants of this operation are possible: A trivial change would be to use  $-F$  instead of  $F$  in the definition of  $\boxplus$ , but also more substantial variations exist [Hie93]. However, all these variations lead to R-matrices that are equivalent in the sense of Def. 1.1.

For characterizing equivalence classes of R-matrices, we next describe how  $\boxplus$  acts on the Yang-Baxter characters of  $S_\infty$  and their Thoma parameters.

**Proposition 4.4.** *Let  $R, \tilde{R} \in \mathcal{R}_0$  have dimensions  $d, \tilde{d}$ .*

*i) The characters of  $R, \tilde{R}$ , and  $R \boxplus \tilde{R}$  are related by ( $c_n$  an  $n$ -cycle,  $n \geq 2$ )*

$$\chi_{R \boxplus \tilde{R}}(c_n) = \frac{d^n}{(d + \tilde{d})^n} \chi_R(c_n) + \frac{\tilde{d}^n}{(d + \tilde{d})^n} \chi_{\tilde{R}}(c_n). \quad (4.5)$$

*ii) Let  $(\alpha, \beta)$  and  $(\tilde{\alpha}, \tilde{\beta})$  be the Thoma parameters of  $R$  and  $\tilde{R}$ , respectively. Then the Thoma parameters of  $R \boxplus \tilde{R}$  are the non-increasing arrangements of*

$$\begin{aligned} \{\hat{\alpha}_i\}_i &= \left\{ \frac{d}{d+\tilde{d}} \alpha_k, \frac{\tilde{d}}{d+\tilde{d}} \tilde{\alpha}_l, : k, l \in \mathbb{N} \right\}, \\ \{\hat{\beta}_i\}_i &= \left\{ \frac{d}{d+\tilde{d}} \beta_k, \frac{\tilde{d}}{d+\tilde{d}} \tilde{\beta}_l, : k, l \in \mathbb{N} \right\}. \end{aligned} \quad (4.6)$$

*Proof.* *i)* We denote the base spaces of  $R$  and  $\tilde{R}$  by  $V_+$  and  $V_-$ , respectively, and write  $\hat{R} := R \boxplus \tilde{R}$  and  $\hat{V} := V_+ \oplus V_-$ . Noting that the dimension of  $\hat{V}$  is  $d + \tilde{d}$ , equation (4.5) is equivalent to

$$\mathrm{Tr}_{\hat{V}^{\otimes n}}(\hat{R}_1 \cdots \hat{R}_{n-1}) = \mathrm{Tr}_{V_+^{\otimes n}}(R_1 \cdots R_{n-1}) + \mathrm{Tr}_{V_-^{\otimes n}}(\tilde{R}_1 \cdots \tilde{R}_{n-1}). \quad (4.7)$$

The trace on the left hand side is taken over  $\hat{V}^{\otimes n} = \bigoplus_{\varepsilon_1, \dots, \varepsilon_n} (V_{\varepsilon_1} \otimes \cdots \otimes V_{\varepsilon_n})$ , where the sum runs over  $\varepsilon_i = \pm$ ,  $i = 1, \dots, n$ . We claim that

$$(\hat{R}_1 \cdots \hat{R}_{n-1})V_{\varepsilon_1} \otimes \cdots \otimes V_{\varepsilon_n} \not\subset V_{\varepsilon_1} \otimes \cdots \otimes V_{\varepsilon_n} \Rightarrow \varepsilon_1 = \cdots = \varepsilon_n. \quad (4.8)$$

Note that (4.8) implies (4.7): If (4.8) holds, then the trace over  $\hat{V}^{\otimes n}$  simplifies to the sum of the trace over  $V_+^{\otimes n}$  and that over  $V_-^{\otimes n}$ . As  $\hat{R}$  acts as  $R$  and  $\tilde{R}$  on  $V_+ \otimes V_+$  and  $V_- \otimes V_-$ , respectively, (4.7) then follows.

To show (4.8), we consider the position of the image  $(\hat{R}_1 \cdots \hat{R}_{n-1})V_{\varepsilon_1} \otimes \cdots \otimes V_{\varepsilon_n}$  relative to  $V_{\varepsilon_1} \otimes \cdots \otimes V_{\varepsilon_n}$  for given  $\varepsilon_1, \dots, \varepsilon_n = \pm$ . Assume that  $\varepsilon_{n-1} \neq \varepsilon_n$ . Then the rightmost factor  $\hat{R}_{n-1}$ , acting non-trivially only on  $V_{\varepsilon_{n-1}} \otimes V_{\varepsilon_n}$ , simplifies to the flip by definition of  $\hat{R} = R \boxplus \tilde{R}$ . As all other factors  $\hat{R}_1 \cdots \hat{R}_{n-2}$  act trivially on the last tensor factor  $V_{\varepsilon_n}$ , this implies  $(\hat{R}_1 \cdots \hat{R}_{n-1})V_{\varepsilon_1} \otimes \cdots \otimes V_{\varepsilon_n} \subset \hat{V}^{\otimes n-1} \otimes V_{\varepsilon_{n-1}} \perp V_{\varepsilon_1} \otimes \cdots \otimes V_{\varepsilon_n}$ . Hence the non-orthogonality assumption in (4.8) implies  $\varepsilon_{n-1} = \varepsilon_n$ .

We next assume  $\varepsilon_{n-2} \neq \varepsilon_{n-1} = \varepsilon_n$ . In this situation, the rightmost factor  $\hat{R}_{n-1}$  maps the product of the last two tensor factors  $V_{\varepsilon_n} \otimes V_{\varepsilon_n}$  onto itself, so that we are left with the same situation as before, but with the number of tensor factors reduced by one. Inductively, we conclude that the non-orthogonality assumption in (4.8) implies  $\varepsilon_1 = \dots = \varepsilon_n$ .

*ii)* Define parameters  $\hat{\alpha}_i, \hat{\beta}_j$  by (4.6), ordered non-increasingly. Then  $0 \leq \hat{\alpha}_i, \hat{\beta}_j \leq 1$ , and for any  $n \in \mathbb{N}$ ,

$$\begin{aligned} \sum_i \hat{\alpha}_i^n + (-1)^{n+1} \sum_j \hat{\beta}_j^n &= \left( \frac{d}{d + \tilde{d}} \right)^n \left( \sum_i \alpha_i^n + (-1)^{n+1} \sum_j \beta_j^n \right) \\ &\quad + \left( \frac{\tilde{d}}{d + \tilde{d}} \right)^n \left( \sum_i \tilde{\alpha}_i^n + (-1)^{n+1} \sum_j \tilde{\beta}_j^n \right). \end{aligned}$$

Since  $(\alpha, \beta), (\tilde{\alpha}, \tilde{\beta}) \in \mathbb{T}$ , we have  $\sum_i \alpha_i + \sum_j \beta_j \leq 1$  and  $\sum_i \tilde{\alpha}_i + \sum_j \tilde{\beta}_j \leq 1$ , and therefore  $\sum_i \hat{\alpha}_i + \sum_j \hat{\beta}_j \leq 1$ . This shows that  $(\hat{\alpha}, \hat{\beta}) \in \mathbb{T}$ . In terms of characters, the above equation reads,  $n \in \mathbb{N}$ ,

$$\sum_i \hat{\alpha}_i^n + (-1)^{n+1} \sum_j \hat{\beta}_j^n = \frac{d^n}{(d + \tilde{d})^n} \chi_R(c_n) + \frac{\tilde{d}^n}{(d + \tilde{d})^n} \chi_{\tilde{R}}(c_n),$$

and by part *i)* and the uniqueness of the Thoma parameters of an extremal character, identifies  $(\hat{\alpha}, \hat{\beta})$  as the Thoma parameters of  $\chi_{\hat{R}}$ .  $\square$

By construction,  $\boxplus$  maps pairs of parameters in  $\mathbb{T}_{\text{YB}}$  into  $\mathbb{T}_{\text{YB}}$ , preserving the three properties of  $\mathbb{T}_{\text{YB}}$  (Def. 3.5). But given  $d, \tilde{d} > 0$ , (4.5) also makes sense as an operation on general extremal characters of  $S_\infty$ . We do not investigate this observation any further here.

After these preparations, we come to the definition of special “normal form R-matrices” as  $\boxplus$ -sums of identities and negative identities. We will write  $1_a$  for the identity on a vector space of dimension  $a^2$ , i.e.  $1_a \in \mathcal{R}_0(\mathbb{C}^a)$ .

**Definition 4.5.** *Let  $n, m \in \mathbb{N}_0$  with  $n + m \geq 1$ ,  $d^+ \in \mathbb{N}^n$  and  $d^- \in \mathbb{N}^m$ . The normal form R-matrix  $N$  with dimensions  $d^+, d^-$  is*

$$N := 1_{d_1^+} \boxplus \dots \boxplus 1_{d_n^+} \boxplus (-1_{d_1^-}) \boxplus \dots \boxplus (-1_{d_m^-}). \quad (4.9)$$

Any R-matrix of the type (4.9) will be called normal form R-matrix. Note that in view of Prop. 4.3,  $N$  is indeed an involutive R-matrix, of dimension  $|d^+| + |d^-|$ , where  $|d^\pm| = \sum_i d_i^\pm$ . We emphasize that  $N$  is not simply the identity: For example,  $1_1 \boxplus 1_1 = F$  is the flip of dimension 2.

**Lemma 4.6.** *Let  $N \in \mathcal{R}_0$  be the normal form R-matrix with dimensions  $d^+ \in \mathbb{N}^n, d^- \in \mathbb{N}^m$ . The Thoma parameters of  $\chi_N$  are*

$$\alpha_i = \frac{d_i^+}{d}, \quad i = 1, \dots, n, \quad \beta_j = \frac{d_j^-}{d}, \quad j = 1, \dots, m, \quad (4.10)$$

where  $d = |d^+| + |d^-|$ .

*Proof.* Recall that the identity  $1 \in \mathcal{R}_0(V)$  has  $\alpha_1 = 1$  as its only non-vanishing Thoma parameter, and the negative identity  $-1 \in \mathcal{R}_0(V)$  has  $\beta_1 = 1$  as its only non-vanishing Thoma parameter, independently of the dimension of  $V$ . From this observation and the fact that  $\boxplus$  adds dimensions, one can easily compute the Thoma parameters of  $N$  (4.9) by iterating Prop. 4.4 *ii*), with the claimed result  $\alpha = d^+/d$  and  $\beta = d^-/d$ .  $\square$

In Thm. 3.6 we had proven that the Thoma parameters of every Yang-Baxter character lie in  $\mathbb{T}_{\text{YB}}$ . Lemma 4.6 now implies the converse.

**Theorem 4.7.** *The Yang-Baxter characters of  $S_\infty$  are in one to one correspondence with  $\mathbb{T}_{\text{YB}}$  (Def. 3.5) via Thoma's formula (2.6).*

*Proof.* Let  $(\alpha, \beta) \in \mathbb{T}_{\text{YB}}$ . All that remains to be shown is that there is an R-matrix with these Thoma parameters. There exists  $d \in \mathbb{N}$  such that all  $d\alpha_i, d\beta_j$  are integer because the  $\alpha_i, \beta_j$  are rational and finite in number (Def. 3.5). The character of the normal form R-matrix  $N$  with dimensions  $d_i^+ = d\alpha_i, d_j^- = d\beta_j$  then has Thoma parameters  $(\alpha, \beta)$  by Lemma 4.6.  $\square$

This result also justifies the notation  $\mathbb{T}_{\text{YB}}$  as the ‘‘Yang-Baxter simplex’’, consisting of all Thoma parameters of Yang-Baxter characters. Thoma's simplex  $\mathbb{T}$ , viewed as a subset of  $[0, 1]^\infty \times [0, 1]^\infty$ , where  $[0, 1]^\infty$  is equipped with the product topology, is a compact metrizable space. It is noteworthy to point out that  $\mathbb{T}_{\text{YB}} \subset \mathbb{T}$  is a dense subset, cf. [BO17, Ch. 3].

At this stage, we know that every R-matrix  $R \in \mathcal{R}_0$  is equivalent to a normal form R-matrix. We briefly mention further properties of normal form R-matrices: Any normal form R-matrix  $N$  commutes with the flip because of Lemma 4.2 *iii*). Thus any involutive R-matrix is equivalent to an R-matrix which commutes with the flip, though this need not be true for an R-matrix not in normal form.

Furthermore, one can check that any normal form R-matrix  $N$  satisfies

$$N(1 \otimes \text{ptr } N)N = \text{ptr } N \otimes 1. \quad (4.11)$$

By Thm. 3.3, a normal form R-matrix  $N$  (of dimension  $d$ ) satisfies, as any involutive R-matrix,

$$\chi_N(c_n) = d^{-n} \text{Tr}_{V^{\otimes n}}(N_1 \cdots N_{n-1}) = d^{-n} \text{Tr}_V((\text{ptr } N)^{n-1}), \quad (4.12)$$

where  $c_n$  is an  $n$ -cycle,  $n \geq 2$ . With the exchange relation (4.11), it is a matter of explicit calculation to prove (4.12) directly for normal form R-matrices, without relying on subfactor theory.

## 4.2 Parameterization by pairs of Young diagrams

The correspondence in Thm. 4.7 classifies the family of Yang-Baxter *characters*, but it does not classify  $\mathcal{R}_0/\sim$  because the dimension of the base space is not recorded in the Thoma parameters. However, it is now easy to incorporate the dimension as well: Given  $R \in \mathcal{R}_0$  with Thoma parameters  $(\alpha, \beta)$  and dimension  $d$ , we switch to the *rescaled Thoma parameters*

$$a_i := d\alpha_i, \quad b_i := d\beta_i. \quad (4.13)$$

By Lemma 4.6, the  $a_i, b_i$  are integers summing to  $d$ . We can therefore view  $(a, b)$  as an ordered pair of integer partitions, or, equivalently, Young diagrams. Denoting the set of all Young diagrams (with an arbitrary number of boxes) by  $\mathbb{Y}$ , we arrive at the following theorem.

### Theorem 4.8.

- i)  $\mathcal{R}_0/\sim$  is in one to one correspondence with  $\mathbb{Y} \times \mathbb{Y} \setminus \{(\emptyset, \emptyset)\}$  via mapping  $[R]$  to the pair  $(a, b)$  (4.13). Classes of  $R$ -matrices of dimension  $d$  correspond to pairs of Young diagrams with  $d$  boxes in total.*
- ii) Let  $R \in \mathcal{R}_0$ . The eigenvalues of  $\text{ptr } R$  lie in  $\{\pm 1, \pm 2, \dots, \pm d\}$  and for each eigenvalue  $\lambda$ , there exists  $n_\lambda \in \mathbb{N}$  such that its multiplicity is  $n_\lambda \cdot |\lambda|$ . Define an integer partition  $a$  as the ordered set of positive eigenvalues, in which  $\lambda$  is repeated  $n_\lambda$  times, and analogously for  $b$  and the negative eigenvalues. Then  $R$  corresponds to  $(a, b)$  via the bijection in part i).*

*Proof.* *i)* If  $R, S \in \mathcal{R}_0$  are equivalent, they have the same dimension  $d$  and the same Thoma parameters  $(\alpha, \beta)$  and hence the same rescaled parameters (4.13). Conversely, if  $R, S \in \mathcal{R}_0$  have the same parameters  $(a, b)$ , they have the same dimension  $d = \sum_i (a_i + b_i)$  and therefore the same Thoma parameters, i.e.  $R \sim S$ . This also shows the claim about the dimension, and that all pairs of Young diagrams with the exception of  $(\emptyset, \emptyset)$  occur.

*ii)* We may switch from  $R$  to its normal form  $N$  (with dimensions  $d^\pm$ ), which has the same partial trace  $\text{ptr } N \cong \text{ptr } R$ . Repeated application of Lemma 4.2 *iv)* shows  $\text{ptr } N = \bigoplus_i \text{ptr}(1_{d_i^+}) \oplus \bigoplus_i \text{ptr}(-1_{d_i^-})$ . But  $\text{ptr}(\pm 1_{d_i^\pm}) = \pm d_i^\pm \text{id}_{i, \pm}$ , where  $\text{id}_{i, \pm}$  is the identity matrix on  $\mathbb{C}^{d_i^\pm}$ . Hence the eigenvalues of  $\text{ptr } R$  are  $\pm d_i^\pm$ . Defining  $n_{d_i^\pm}$  as the number of times  $d_i^\pm$  occurs in  $d^\pm$ , we see that  $\pm d_i^\pm$  has multiplicity  $n_{d_i^\pm} \cdot d_i^\pm$ ,  $i = 1 \dots, k$ .

In view of (4.10) and (4.13), the rescaled Thoma parameters of  $N$  are exactly  $a_i = d_i^+$ ,  $b_i = d_i^-$ . As  $d_i^\pm$  occurs  $n_{d_i^\pm}$  times in this list, the proof is finished.  $\square$

To illustrate the correspondence with ordered pairs of Young diagrams, let us list all normal forms of dimension two in terms of box sums and diagrams:

$1_2$	$-1_2$	$1_1 \boxplus 1_1$	$-1_1 \boxplus -1_1$	$1_1 \boxplus -1_1$
$(\square, \emptyset)$	$(\emptyset, \square)$	$(\begin{array}{ c } \hline \square \\ \hline \end{array}, \emptyset)$	$(\emptyset, \begin{array}{ c } \hline \square \\ \hline \end{array})$	$(\square, \square)$

From left to right, these R-matrices (in  $\text{End}(\mathbb{C}^4)$ ) are: 1) the identity, 2) the negative identity, 3) the flip, 4) equivalent to the negative flip, and 5)

$$1_1 \boxplus -1_1 = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & -1 \end{pmatrix}.$$

As a higher-dimensional example, consider  $(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}, \begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix})$ . This R-matrix has dimension 8 (the number of boxes) and Thoma parameters  $\alpha = (\frac{3}{8}, \frac{1}{8})$ ,  $\beta = (\frac{1}{4}, \frac{1}{4})$ .

The rescaled Thoma parameters are also useful for describing the  $\boxplus$  operation introduced in Sect. 4.1. We have already seen that  $\boxplus$  gives  $\mathcal{R}_0$  the structure of an abelian semigroup and preserves equivalence, i.e., descends to the quotient  $\mathcal{R}_0/\sim$ . Recalling the effect of  $\boxplus$  on the level of Thoma parameters (Prop. 4.4 *ii*), it becomes apparent that for the rescaled parameters, we have

$$(a, b) \boxplus (a', b') = (a \cup a', b \cup b'), \quad (4.14)$$

where  $a \cup a'$  denotes the partition whose parts are the union of those of  $a$  and  $a'$ .

Another operation on  $\mathcal{R}$  is the tensor product of R-matrices. For  $R \in \mathcal{R}(V)$ ,  $S \in \mathcal{R}(W)$ , we define  $R \boxtimes S \in \mathcal{R}(V \otimes W)$  by

$$R \boxtimes S = F_2(R \otimes S)F_2 : V \otimes W \otimes V \otimes W \rightarrow V \otimes W \otimes V \otimes W, \quad (4.15)$$

where  $F_2$  exchanges the second and third tensor factors. It is evident that  $\boxtimes$  preserves the Yang-Baxter equation and involutivity, i.e. it defines a product on  $\mathcal{R}$  and  $\mathcal{R}_0$ .

**Lemma 4.9.** *Let  $R, R' \in \mathcal{R}_0$  have rescaled Thoma parameters  $(a, b)$  and  $(a', b')$ , respectively. Then the rescaled Thoma parameters of  $R \boxtimes R'$  are the non-increasing arrangements of*

$$\begin{aligned} \{\hat{a}_{ij}\} &= \{a_i a'_j, b_i b'_j\}, \\ \{\hat{b}_{ij}\} &= \{a_i b'_j, b_i a'_j\}. \end{aligned} \quad (4.16)$$

*Proof.* With  $d, d'$  the dimensions of  $R, R'$ , we have on an  $n$ -cycle,  $n \geq 2$ ,

$$\begin{aligned} (d \cdot d')^n \chi_{R \boxtimes R'}(c_n) &= \text{Tr}_{(V \otimes W)^{\otimes n}}((R \boxtimes R')_1 \cdots (R \boxtimes R')_{n-1}) \\ &= \text{Tr}_{V^{\otimes n}}(R_1 \cdots R_{n-1}) \text{Tr}_{W^{\otimes n}}(R'_1 \cdots R'_{n-1}) \\ &= d^n \chi_R(c_n) \cdot (d')^n \chi_{R'}(c_n) \\ &= \left( \sum_i a_i^n + (-1)^{n+1} \sum_j b_j^n \right) \left( \sum_k (a'_k)^n + (-1)^{n+1} \sum_l (b'_l)^n \right) \\ &= \sum_{i,k} (a_i a'_k)^n + \sum_{j,l} (b_j b'_l)^n + (-1)^{n+1} \left( \sum_{i,l} (a_i b'_l)^n + \sum_{j,k} (b_j a'_k)^n \right), \end{aligned}$$

and the claim follows.  $\square$

It follows that  $\boxtimes$  defines an associative commutative product on  $\mathcal{R}_0/\sim$  for which the class  $[1_1] = (\square, \emptyset)$  (consisting of the identity R-matrix in dimension  $d = 1$ ) is the unit, that is,  $\mathcal{R}_0/\sim$  has a second unital abelian semigroup structure.

From the description of  $\boxtimes$  and  $\boxplus$  in terms of the rescaled Thoma parameters, it is evident that they satisfy the distributive law

$$([R] \boxplus [S]) \boxtimes [T] = ([R] \boxtimes [T]) \boxplus ([S] \boxtimes [T]), \quad R, S, T \in \mathcal{R}_0. \quad (4.17)$$

These operations give  $\mathcal{R}_0/\sim$  the structure of a semiring, sometimes also referred to as a “rig” (ring without negatives). Additionally, the multiplication rules for rescaled Thoma parameters in Lemma 4.9 can be generalized to a  $\lambda$ -operation. This  $\lambda$ -operation is most easily described using symmetric polynomials and we therefore postpone it until Sect. 5.2. The consequences of the ring and  $\lambda$  structures of  $\mathcal{R}_0/\sim$  will be an interesting topic of further study.

As yet another operation on  $\mathcal{R}$ , we briefly mention the cabling procedure known from the braid groups, applied to the Yang-Baxter equation by Wenzl [Wen90]: Given any  $p \in \mathbb{N}$ , one can form “cabling powers”  $R^{c(p)}$ , which lie in  $\mathcal{R}$  (or  $\mathcal{R}_0$ ) if  $R$  does. We do not give details here because it turns out that  $R^{c(p)} \sim R^{\boxtimes p}$  for all  $R \in \mathcal{R}_0$ ,  $p \in \mathbb{N}$ .

## 5 Yang-Baxter representations

Our basic Def. 1.1 of equivalence of R-matrices refers only to the  $S_n$ -representations  $\rho_R^{(n)}$ . We have seen already that  $R \sim S$  implies unitary equivalence of the GNS representations  $\pi_\tau \circ \rho_R \cong \pi_\tau \circ \rho_S$  (3.3). Now we investigate the implications of  $R \sim S$  for the homomorphisms  $\rho_R, \rho_S$ .

### 5.1 R-matrices and $K$ -theory

In this section we extend the previously defined  $\rho_R$  to a  $*$ -homomorphism of  $C^*$ -algebras,  $\rho_R: C^*S_\infty \rightarrow \mathcal{E}^\infty$ , where  $\mathcal{E}^\infty$  is the  $C^*$ -algebraic counterpart of the algebra  $\mathcal{E}$  from Section 3. On  $K$ -theory the map  $\rho_R$  will induce a ring homomorphism  $\rho_{R*}: K_0(C^*S_\infty) \rightarrow \mathbb{Z}[\frac{1}{d}]$  with  $d = \dim(V)$ . The equivalence relation introduced in Def. 1.1 will then translate into the approximate unitary equivalence of the corresponding  $*$ -homomorphisms. In fact, when the invariant  $\rho_{R*}$  is composed with the canonical inclusion  $\mathbb{Z}[\frac{1}{d}] \subset \mathbb{R}$  it is an indecomposable finite trace on  $K_0(C^*S_\infty)$  in the sense of Kerov and Vershik and we recover [VK83, Thm. 2.3] from the Yang-Baxter equation. For the basic facts about UHF-algebras that we use we refer the reader to [RS02].

Let  $R \in \mathcal{R}_0(V)$ , let  $d = \dim(V)$  and denote the associated unitary representation of  $S_n$  by  $\rho_R^{(n)}$ . We obtain the following sequence of  $*$ -homomorphisms, which we will continue to denote  $\rho_R^{(n)}$ :

$$\rho_R^{(n)}: C^*(S_n) = \mathbb{C}[S_n] \rightarrow \text{End}(V^{\otimes n})$$

Let  $\mathcal{E}^\infty$  be the  $C^*$ -algebra obtained as the infinite tensor product of the algebras  $\text{End}(V)$ , i.e. as the  $C^*$ -algebraic inductive limit

$$\mathcal{E}^\infty = \varinjlim_n \text{End}(V^{\otimes n})$$

taken over the maps sending  $T$  to  $T \otimes \text{id}_V$ . This is an infinite UHF-algebra with  $K_0(\mathcal{E}^\infty) \cong \mathbb{Z}[\frac{1}{d}]$  and  $K_1(\mathcal{E}^\infty) \cong 0$ . The algebra  $\mathcal{E}^\infty$  has a unique trace that restricts to the normalized trace on  $\text{End}(V^{\otimes n})$ . It induces an explicit isomorphism  $\tau_*: K_0(\mathcal{E}^\infty) \rightarrow \mathbb{Z}[\frac{1}{d}]$  as follows: Let  $p, q \in M_N(\mathcal{E}^\infty)$  be projections. Then

$$\tau_*([p] - [q]) = (\text{Tr}_N \otimes \tau)(p) - (\text{Tr}_N \otimes \tau)(q) ,$$

where  $\text{Tr}_N \otimes \tau: M_N(\mathcal{E}^\infty) = M_N(\mathbb{C}) \otimes \mathcal{E}^\infty \rightarrow \mathbb{C}$  is induced by the non-normalized trace  $\text{Tr}_N$  tensored with  $\tau$ . This is in fact a ring isomorphism. To understand the ring structure on  $K_0(\mathcal{E}^\infty)$  note that  $\mathcal{E}^\infty$  is strongly self-absorbing [TW07, Ex. 1.14]. In particular, there is an isomorphism  $\psi: \mathcal{E}^\infty \otimes \mathcal{E}^\infty \rightarrow \mathcal{E}^\infty$  and any two such isomorphisms are homotopic. Let  $p_i \in M_{N_i}(\mathcal{E}^\infty)$  for  $i \in \{1, 2\}$  be projections and let  $[p_i] \in K_0(\mathcal{E}^\infty)$  be the corresponding  $K$ -theory classes. Let  $\psi': M_{N_1}(\mathcal{E}^\infty) \otimes M_{N_2}(\mathcal{E}^\infty) \rightarrow M_{N_1 N_2}(\mathcal{E}^\infty)$  be the isomorphism induced by  $\psi$ . Then we have  $[p_1] \cdot [p_2] = [\psi'(p_1 \otimes p_2)]$ . It follows from the uniqueness of the normalized trace on  $\mathcal{E}^\infty$  that

$$(\text{Tr}_{N_1} \otimes \tau) \otimes (\text{Tr}_{N_2} \otimes \tau) = (\text{Tr}_{N_1 N_2} \otimes \tau) \circ \psi' ,$$

which implies  $\tau_*([p_1] \cdot [p_2]) = \tau_*([p_1]) \cdot \tau_*([p_2])$ .

Note that  $\mathcal{E}^\infty \subset \mathcal{E}$  with  $\mathcal{E}$  as in Section 3. The inductive limit of the representations  $\rho_R^{(n)}: C^*(S_n) \rightarrow \text{End}(V^{\otimes n})$  provides us with a  $*$ -homomorphism

$$\rho_R: C^*S_\infty \rightarrow \mathcal{E}^\infty . \quad (5.1)$$

The  $K$ -theory of  $C^*S_\infty$  was studied by Kerov and Vershik in [VK83]. In particular, they obtained that  $K_0(C^*S_\infty)$  is isomorphic to a quotient of the ring of symmetric functions. As an abelian group it is therefore spanned by projections  $p_\lambda \in K_0(C^*S_\infty)$ , that are labeled by partitions  $\lambda = [\lambda_1, \dots, \lambda_k]$  of natural numbers  $n \in \mathbb{N}$ . The map  $\rho_R$  induces a group homomorphism

$$\rho_{R*}: K_0(C^*S_\infty) \rightarrow K_0(\mathcal{E}^\infty) \quad (5.2)$$

in  $K$ -theory. Using the ring isomorphism induced by the unique trace on  $\mathcal{E}^\infty$  we will identify  $K_0(\mathcal{E}^\infty)$  with  $\mathbb{Z}[\frac{1}{d}]$ . The following lemma shows that  $\rho_{R*}$  remembers the equivalence class of  $R$ .

**Lemma 5.1.** *Let  $\lambda$  be a partition of  $n \in \mathbb{N}$ . We will identify  $\lambda$  with the corresponding irreducible representation of  $S_n$ . On the projection  $p_\lambda \in C^*S_\infty$  associated to  $\lambda$  the value of  $\rho_{R*}$  is given by*

$$\rho_{R*}([p_\lambda]) = \frac{1}{d^n} \langle \lambda, \rho_R^{(n)} \rangle ,$$

where  $\langle \lambda, \mu \rangle$  denotes the multiplicity of the irreducible representation  $\lambda$  in the representation  $\mu$ .

*Proof.* We have  $\rho_{R*}([p_\lambda]) = \tau(\rho_R(p_\lambda))$ . Let  $\tau_n: \text{End}(V^{\otimes n}) \rightarrow \mathbb{C}$  be the normalized trace. Since  $p_\lambda \in C^*(S_n) \subset C^*S_\infty$  and the inclusion  $\text{End}(V^{\otimes n}) \rightarrow \mathcal{E}^\infty$  preserves the normalized trace, we obtain

$$\tau(\rho_R(p_\lambda)) = \tau_n(\rho_R^{(n)}(p_\lambda)) = \frac{1}{d^n} \text{Tr}_{V^{\otimes n}}(\rho_R^{(n)}(p_\lambda)) .$$

Let  $V_R = V^{\otimes n}$  be the representation space of  $\rho_R^{(n)}$ . The decomposition into its irreducible components gives

$$V_R \cong \bigoplus_{\mu \in \text{Irrep}(S_n)} \text{hom}_{C^*(S_n)}(V_\mu, V_R) \otimes V_\mu ,$$

where the action on the left is via  $\rho_R^{(n)}$  and on the right acts only on the second tensor factor  $V_\mu$  via  $\mu$ . Observe that  $p_\lambda V_\mu$  is zero for  $\lambda \neq \mu$  and 1-dimensional for  $\lambda = \mu$ . Hence,

$$\begin{aligned} \text{Tr}_{V^{\otimes n}}(\rho_R^{(n)}(p_\lambda)) &= \dim(\rho_R^{(n)}(p_\lambda)V_R) = \dim(\text{hom}_{C^*(S_n)}(V_\lambda, V_R) \otimes p_\lambda V_\lambda) \\ &= \dim(\text{hom}_{C^*(S_n)}(V_\lambda, V_R)) = \langle \lambda, \rho_R^{(n)} \rangle . \end{aligned} \quad \square$$

From this we obtain two useful additional characterizations of the equivalence relation from Def. 1.1, one of them  $K$ -theoretic, the other one  $C^*$ -algebraic. For the second one we need the following equivalence relation [RS02, Def. 1.1.15]:

**Definition 5.2.** *Let  $\varphi, \psi: A \rightarrow B$  be  $*$ -homomorphisms between separable unital  $C^*$ -algebras  $A$  and  $B$ . We call them approximately unitarily equivalent if there is a sequence of unitaries  $u_n \in B$  with the property that for all  $a \in A$  we have*

$$\lim_{n \rightarrow \infty} \|\varphi(a) - u_n \psi(a) u_n^*\| = 0 .$$

We denote this by  $\varphi \approx_u \psi$ .

**Theorem 5.3.** *Let  $R, S \in \mathcal{R}_0(V)$ . The following are equivalent:*

- i)  $R \sim S$ ,*
- ii)  $\rho_{R^*} = \rho_{S^*}$ ,*
- iii)  $\rho_R \approx_u \rho_S$ .*

*Proof.* The equivalence of *i)* and *ii)* is a consequence of Lemma 5.1 and the fact that  $\rho_R^{(n)}$  and  $\rho_S^{(n)}$  are unitarily equivalent if and only if the multiplicities of their irreducible subrepresentations agree.

To see that *ii)* and *iii)* are equivalent, note that the  $C^*$ -algebras  $C^*S_\infty$  and  $\mathcal{E}^\infty$  are both AF-algebras. The statement then follows from [RS02, Prop. 1.3.4].  $\square$

The  $K$ -group  $K_0(C^*S_\infty)$  is in fact a ring: Let  $\lambda$  be a partition of  $n \in \mathbb{N}$  and let  $\mu$  be a partition of  $m \in \mathbb{N}$ . Denote by  $p_\lambda, p_\mu \in C^*S_\infty$  the associated projections. Let  $\iota_{n,m}: C^*(S_n) \otimes C^*(S_m) \rightarrow C^*(S_{n+m})$  be the  $*$ -homomorphism induced by the inclusion  $S_n \times S_m \rightarrow S_{n+m}$ , where  $S_n$  permutes the first  $n$  elements and  $S_m$  the last  $m$  elements. The product  $[p_\lambda] \cdot [p_\mu]$  is then defined to be the class of the projection  $\iota_{n,m}(p_\lambda \otimes p_\mu) \in C^*(S_{n+m}) \subset C^*S_\infty$  in  $K_0(C^*S_\infty)$ . With respect to this ring structure we make the following observation:

**Proposition 5.4.** *Let  $R \in \mathcal{R}_0(V)$ . Then the associated  $K$ -theory invariant*

$$\rho_{R^*}: K_0(C^*S_\infty) \rightarrow \mathbb{Z}[\frac{1}{d}]$$

*is a ring homomorphism.*



*Proof.* Let  $\lambda, \mu$  be partitions of  $n, m \in \mathbb{N}$  respectively. Let  $p_\lambda \in C^*(S_n), p_\mu \in C^*(S_m)$  be the corresponding projections. Since the representations  $\rho_R^{(n)}$  arise from the same R-matrix, we have  $\rho_R^{(n+m)} \circ \iota_{n,m} = \rho_R^{(n)} \otimes \rho_R^{(m)}$ . Hence, we obtain

$$\rho_{R^*}([p_\lambda] \cdot [p_\mu]) = [\rho_R^{(n+m)} \circ \iota_{n,m}(p_\lambda \otimes p_\mu)] = [\rho_R^{(n)}(p_\lambda) \otimes \rho_R^{(m)}(p_\mu)]$$

and after application of the isomorphism  $\tau_*: K_0(\mathcal{E}^\infty) \rightarrow \mathbb{Z}[\frac{1}{d}]$  induced by the trace:

$$\begin{aligned} \tau_*(\rho_{R^*}([p_\lambda] \cdot [p_\mu])) &= \tau_{n+m}(\rho_R^{(n)}(p_\lambda) \otimes \rho_R^{(m)}(p_\mu)) \\ &= \tau_n(\rho_R^{(n)}(p_\lambda)) \tau_m(\rho_R^{(m)}(p_\mu)) = \tau_*(\rho_{R^*}([p_\lambda])) \cdot \tau_*(\rho_{R^*}([p_\mu])) \end{aligned}$$

where  $\tau_r: \text{End}(V^{\otimes r}) \rightarrow \mathbb{C}$  denotes the (normalized) trace on the matrix algebra and we used that the inclusion  $\text{End}(V^{\otimes r}) \subset \mathcal{E}^\infty$  is trace preserving.  $\square$

Finite traces on  $K_0(C^*S_\infty)$  have been studied by Kerov and Vershik in [VK83] and  $\rho_{R^*}$  can be seen as a refinement of such a trace taking values in  $\mathbb{Z}[\frac{1}{d}]$ . In particular, we recover the multiplicativity proven in [VK83, Thm. 2.3] in Prop. 5.4. From the point of view of  $C^*$ -algebras it is surprising that we obtain a ring homomorphism on  $K$ -theory that is induced by a  $*$ -homomorphism.

## 5.2 R-matrices, $K$ -theory and symmetric functions

Before discussing the connections between R-matrices,  $K$ -theory and symmetric functions in greater detail, we first collect some facts about symmetric functions to fix notation. We refer readers unfamiliar with symmetric functions to Macdonald's book [Mac95].

The ring of symmetric functions,  $\Lambda$ , admits numerous free generators. Here, the most important are:

- i) Elementary symmetric functions:

$$\mathbf{e}_k = \sum_{1 \leq i_1 < i_2 < \dots < i_k} x_{i_1} x_{i_2} \cdots x_{i_k}, \quad k \geq 1.$$

- ii) Complete symmetric functions:

$$\mathbf{h}_k = \sum_{1 \leq i_1 \leq i_2 \leq \dots \leq i_k} x_{i_1} x_{i_2} \cdots x_{i_k}, \quad k \geq 1.$$

- iii) Power sums:

$$\mathbf{p}_k = \sum_{i \geq 1} x_i^k.$$

The ring of symmetric functions also admits many interesting bases usually indexed by partitions of integers. For example the above three sets of generators each define a basis by having the basis vector associated to a partition  $\lambda = [\lambda_1, \lambda_2, \dots]$  be  $f_\lambda = f_{\lambda_1} f_{\lambda_2} \cdots$ , where  $f$  is either  $\mathbf{e}, \mathbf{h}$  or  $\mathbf{p}$  and one defines  $f_0 = 1$ .

An additional important basis is given by Schur functions  $\mathbf{s}_\lambda$ , which in terms of elementary and complete symmetric functions are given by the following determinantal formulae:

$$\mathbf{s}_\lambda = \det (\mathbf{h}_{\lambda_i - i + j})_{1 \leq i, j \leq n} = \det (\mathbf{e}_{\lambda'_i - i + j})_{1 \leq i, j \leq m},$$

where  $n \geq \ell(\lambda)$ ,  $\lambda'$  is the partition conjugate to  $\lambda$  and  $m \geq \ell(\lambda')$ .

The ring of symmetric functions also admits a ring involution  $\omega : \Lambda \rightarrow \Lambda$  which on the generators and bases defined above acts as

$$\omega(\mathbf{e}_k) = \mathbf{h}_k, \quad \omega(\mathbf{h}_k) = \mathbf{e}_k, \quad \omega(\mathbf{p}_k) = (-1)^{k+1} \mathbf{p}_k, \quad \omega(\mathbf{s}_\lambda) = \mathbf{s}_{\lambda'}.$$

Finally, we will also make use of the coproduct  $\Delta : \Lambda \rightarrow \Lambda \otimes \Lambda$  which maps a symmetric function  $f(x)$  to the same function  $f(x, y)$  but with the alphabet of variables split into two alphabets.

The images of the sets of generators defined above are then

$$\begin{aligned} \Delta(\mathbf{e}_k) &= \sum_{q=0}^k \mathbf{e}_q(x) \mathbf{e}_{k-q}(y), \quad \Delta(\mathbf{h}_k) = \sum_{q=0}^k \mathbf{h}_q(x) \mathbf{h}_{k-q}(y), \\ \Delta(\mathbf{p}_k) &= \mathbf{p}_k(x) + \mathbf{p}_k(y). \end{aligned}$$

The corresponding formulae for Schur functions are more involved but can be derived from their determinantal expressions in terms of elementary or complete symmetric functions.

The elementary symmetric functions and their coproducts can be used to define a  $\lambda$ -operation on rescaled Thoma parameters  $(a, b)$  in the following way: Denote the  $\lambda^n$  operation on  $(a, b)$  by  $\lambda^n(a, b) = (\lambda^n a, \lambda^n b)$ , where  $\lambda^n a$  and  $\lambda^n b$  are the non-decreasing arrangements of

$$\begin{aligned} \lambda^n a &= \{\text{monomial summands of } \mathbf{e}_n(a, b) \text{ with even number of factors from } b\}, \\ \lambda^n b &= \{\text{monomial summands of } \mathbf{e}_n(a, b) \text{ with odd number of factors from } b\}. \end{aligned}$$

For example if  $(a, b) = ([a_1, a_2], [b_1])$ ,

$$\begin{aligned} \mathbf{e}_0(a, b) &= 1, \quad \mathbf{e}_1(a, b) = a_1 + a_2 + b_1, \quad \mathbf{e}_2(a, b) = a_1 a_2 + a_1 b_1 + a_2 b_1, \\ \mathbf{e}_3(a, b) &= a_1 a_2 b_1, \quad \mathbf{e}_n(a, b) = 0, \quad n \geq 4. \end{aligned}$$

Thus,

$$\begin{aligned} \lambda^0(a, b) &= ([1], \emptyset), \quad \lambda^1(a, b) = ([a_1, a_2], [b_1]), \quad \lambda^2(a, b) = ([a_1 a_2], [a_1 b_1, a_2 b_1]), \\ \lambda^3(a, b) &= (\emptyset, [a_1 a_2 b_1]), \quad \lambda^n(a, b) = (\emptyset, \emptyset), \quad n \geq 4. \end{aligned}$$

**Lemma 5.5.** *The operation  $\lambda^n$  is a  $\lambda$ -operation.*

*Proof.* In order for  $\lambda^n(a, b)$  to be a  $\lambda$ -operation it must satisfy  $\lambda^0(a, b) = (\square, \emptyset)$ ,  $\lambda^1(a, b) = (a, b)$  and  $\lambda^n((a, b) \boxplus (c, d)) = \boxplus_{q=0}^n \lambda^q(a, b) \boxtimes \lambda^{n-q}(c, d)$ . The first two properties follow directly from the definition of  $\mathbf{e}_0$  and  $\mathbf{e}_1$ , while the last property follows from the action of the coproduct  $\Delta$  on elementary symmetric functions.  $\square$

Let  $I \subset \Lambda$  be the ideal generated by  $\mathbf{e}_1 - 1$  and let  $\widehat{\Lambda} = \Lambda/I$ . Kerov and Vershik point out that the homomorphism  $\theta: \widehat{\Lambda} \rightarrow K_0(C^*S_\infty)$  fixed by  $\theta(\mathbf{s}_\lambda) = [p_\lambda]$  is in fact a ring isomorphism [VK83]. Using this identification we can now completely determine the K-theory invariant  $\rho_{R^*}$  in terms of the Thoma parameters of  $R$ .

**Theorem 5.6.** *Let  $R \in \mathcal{R}_0(V)$ . Let  $(\alpha, \beta)$  be the Thoma parameters of  $R$ . Then we have*

$$\rho_{R^*}(\theta(\mathbf{e}_k)) = [(1 \otimes \omega) \circ \Delta(\mathbf{e}_k)](\alpha, \beta)$$

*Proof.* The generating function  $g_R$  associated to the trace  $\varphi = \tau_* \circ \rho_{R^*} \circ \theta: \widehat{\Lambda} \rightarrow \mathbb{Z}[\frac{1}{d}] \subset \mathbb{R}$  is given by

$$g_R(z) = \sum_{l=0}^{\infty} \varphi(\mathbf{e}_l) z^l$$

as described in [VK83, eq. (11)] and is related to the Thoma parameters  $(\alpha, \beta)$  as follows [VK83, eq. (12)] (note that  $\gamma = 0$  and  $N = \max\{n, m\}$  in our case):

$$g_R(z) = \prod_{i=1}^N \frac{1 + \alpha_i z}{1 - \beta_i z}.$$

Hence, the statement follows from the following computation and comparison of coefficients with  $g_R(z)$ :

$$\begin{aligned} & \sum_{l=0}^{\infty} [(1 \otimes \omega) \circ \Delta(\mathbf{e}_l)](\alpha, \beta) z^l \\ &= \sum_{l=0}^{\infty} \sum_{i+j=l} \mathbf{e}_i(\alpha_1, \dots, \alpha_n) \mathbf{h}_j(\beta_1, \dots, \beta_m) z^l \\ &= \left( \sum_{i=0}^{\infty} \mathbf{e}_i(\alpha_1, \dots, \alpha_n) z^i \right) \left( \sum_{j=0}^{\infty} \mathbf{h}_j(\beta_1, \dots, \beta_m) z^j \right) \\ &= \prod_{i=1}^n (1 + \alpha_i z) \prod_{j=1}^m \frac{1}{1 - \beta_j z} = \prod_{i=1}^N \frac{1 + \alpha_i z}{1 - \beta_i z}. \quad \square \end{aligned}$$

An immediate consequence of the above theorem is that applying  $\rho_{R^*}(\theta(-))$  to a power sum  $\mathbf{p}_\lambda$  is the same as evaluating the class function  $\chi_R$  at a group element of cycle shape  $\lambda$ . Moreover, Lemma 5.1, Theorem 5.6 and the fact that  $\theta(\mathbf{s}_\lambda) = [p_\lambda]$  can now be used to easily derive explicit formulae for the multiplicities of irreducible representations of  $S_n$  in  $\rho_R^{(n)}$ .

**Proposition 5.7.** *Let  $R \in \mathcal{R}_0$  with rescaled Thoma parameters  $(a, b)$ , then the multiplicity of the  $S_n$  representation associated to a partition  $\lambda$  of  $n$  in  $\rho_R^{(n)}$  is*

$$\langle \lambda, \rho_R^{(n)} \rangle = [(1 \otimes \omega) \circ \Delta(\mathbf{s}_\lambda)](a, b).$$

*Further let  $\ell(a), \ell(b)$  be the respective lengths of  $a$  and  $b$ . Then  $\langle \lambda, \rho_R^{(n)} \rangle = 0$  if and only if the Young diagram of  $\lambda$  contains a rectangle of height  $\ell(a) + 1$  and*



In this section, we briefly discuss how this construction extends to a much larger class of extremal characters when we change  $\tau$  to a different product state on  $\mathcal{E}_0$ : All extremal characters with Thoma parameters summing to 1 can, together with their GNS representations, be expressed in terms of R-matrices and product states. The essential difference to our Yang-Baxter setting is that instead of the canonical trace  $\tau$ , a different product state is used.

Let  $R \in \mathcal{R}_0(V)$  and  $Z \in \text{End}(V)$  such that  $[R, Z \otimes Z] = 0$ . We consider the product state  $\omega_Z := \bigotimes_{n \geq 1} \text{Tr}_V(Z \cdot)$  on  $\mathcal{E}_0$ , which is tracial in restriction to  $\rho_R(\mathbb{C}[S_\infty])$ . Composed with  $\rho_R$ , we thus get a character  $\omega_Z^R := \omega_Z \circ \rho_R$  of  $S_\infty$ . As in Prop. 2.2, one shows that  $\omega_Z^R$  is extremal.

We want to show that any extremal  $S_\infty$ -character with Thoma parameters  $(\alpha, \beta)$  satisfying  $\sum_i (\alpha_i + \beta_i) = 1$  is of this form. To this end, consider two Hilbert spaces  $V_1, V_2$ , such that  $\dim V_1$  and  $\dim V_2$  equal the number of non-vanishing  $\alpha$ 's and  $\beta$ 's, respectively (which might be countably infinite). Let us fix orthonormal bases  $\{e_i\}_i$  and  $\{f_j\}_j$  of  $V_1$  and  $V_2$ , and trace class operators  $A \in \mathcal{B}(V_1), B \in \mathcal{B}(V_2)$  fixed by  $Ae_i = \alpha_i \cdot e_i, Bf_j = \beta_j \cdot f_j$ .

**Lemma 5.8.** *In the notation introduced above, consider the Hilbert space  $W := V_1 \oplus V_2$  and the (now possibly infinite-dimensional) R-matrix  $F \boxplus -F \in \mathcal{R}_0(W)$ . Then*

$$\omega_{A \oplus B}^{F \boxplus -F} := \bigotimes_{n \geq 1} \text{Tr}_W((A \oplus B) \cdot) \circ \rho_{F \boxplus -F} : S_\infty \rightarrow \mathbb{C} \quad (5.3)$$

is an extremal character of  $S_\infty$ . Its  $\alpha$ -parameters are the eigenvalues of  $A$ , and its  $\beta$ -parameters are the eigenvalues of  $B$ .

*Proof.* In view of the simple structure of  $F \boxplus -F$ , it is easy to see that this operator commutes with  $Z \otimes Z$ , where  $Z := A \oplus B$ . Thus  $\omega_{A \oplus B}^{F \boxplus -F}$  is indeed an extremal character, and it remains to compute its Thoma parameters. Using the orthogonality  $V_1 \perp V_2$  and the direct sum structure of  $Z = A \oplus B$ , one shows in close analogy to Prop. 4.4 (see, in particular, (4.7)), that for any  $n$ -cycle,

$$\omega_{A \oplus B}^{F \boxplus -F}(c_n) = \omega_A^F(c_n) + \omega_B^{-F}(c_n). \quad (5.4)$$

Furthermore,

$$\begin{aligned} \omega_A^F(c_n) &= \sum_{i_1, \dots, i_n} \alpha_{i_1} \cdots \alpha_{i_n} \langle e_{i_1} \otimes \dots \otimes e_{i_n}, e_{i_2} \otimes \dots \otimes e_{i_n} \otimes e_{i_1} \rangle = \sum_i \alpha_i^n, \\ \omega_B^{-F}(c_n) &= (-1)^{n+1} \sum_{j_1, \dots, j_n} \beta_{j_1} \cdots \beta_{j_n} \langle f_{j_1} \otimes \dots \otimes f_{j_n}, f_{j_2} \otimes \dots \otimes f_{j_n} \otimes f_{j_1} \rangle \\ &= (-1)^{n+1} \sum_j \beta_j^n. \end{aligned}$$

These two terms sum to the value of the extremal character with Thoma parameters  $(\alpha, \beta)$ .  $\square$

We next describe the GNS representation of  $\omega_{A \oplus B}^{F \boxplus -F}$ , which turns out to be closely related to R-matrices as well. In the notation introduced above, let  $V := W \otimes W$  and

$$R := (F \boxplus -F) \boxtimes 1 \in \mathcal{R}_0(V), \quad (5.5)$$

where  $F \in \mathcal{R}_0(V_1)$ ,  $-F \in \mathcal{R}_0(V_2)$ , and  $1 \in \mathcal{R}_0(W)$ . In  $V$ , we fix the unit vector

$$\xi := \sum_i \sqrt{\alpha_i} e_i \otimes e_i + \sum_j \sqrt{\beta_j} f_j \otimes f_j \in W \otimes W = V. \quad (5.6)$$

This vector determines inclusions  $V^{\otimes n} \rightarrow V^{\otimes(n+1)}$  by tensoring with  $\xi$  from the right, and we denote the corresponding inductive limit Hilbert space  $\bigotimes_{n \geq 1}^\xi V$ .

**Proposition 5.9.** *Let  $\chi$  be an extremal  $S_\infty$ -character with Thoma parameters  $(\alpha, \beta)$  satisfying  $\sum_i (\alpha_i + \beta_i) = 1$ . Then the GNS data  $(\pi_\chi, \mathcal{H}_\chi, \Omega_\chi)$  can be described in terms of the previously introduced  $V, R$ , and  $\xi$  as*

$$\Omega_\chi = \bigotimes_{n \geq 1} \xi, \quad \mathcal{H}_\chi = \overline{\rho_R(\mathbb{C}[S_\infty])\Omega_\chi}, \quad \pi_\chi = \rho_R. \quad (5.7)$$

*Proof.* Observe that for  $w_1, w_2, w_3, w_4 \in W$ , the R-matrix (5.5) acts according to

$$R(w_1 \otimes w_2 \otimes w_3 \otimes w_4) = \pm w_3 \otimes w_2 \otimes w_1 \otimes w_4, \quad (5.8)$$

where the sign is negative if  $w_1$  and  $w_3$  lie in  $V_2$ , and positive if at least one of these vectors lies in  $V_1$ .

Let  $w_1, \dots, w_n, u_1, \dots, u_n \in W$ , and  $\sigma \in S_n$ . Then this action of  $R$  implies

$$\rho_R(\sigma) \bigotimes_{k=1}^n (w_k \otimes u_k) = \pm \bigotimes_{k=1}^n (w_{\sigma^{-1}(k)} \otimes u_k), \quad (5.9)$$

with the sign depending on the number of vectors  $w_k$  lying in  $V_2$ . From here one verifies

$$\langle \bigotimes_{n \geq 1} \xi, \rho_R(\sigma) \bigotimes_{n \geq 1} \xi \rangle = \chi(\sigma) \quad (5.10)$$

by following [BO17, Prop. 10.5, Prop. 10.6]. □

The representation in the above proposition is known (see the original literature [Ols90, Was81] or the monograph [BO17], where also the relation to spherical representations of  $S_\infty$  is discussed), but takes a particularly simple formulation in terms of our operations  $\boxplus$  and  $\boxtimes$ .

## 6 Examples

In this section, we discuss two special classes of involutive R-matrices.

### 6.1 R-matrices of diagonal type

As a simple class of examples which exist in any dimension, we consider involutive R-matrices of *diagonal type*. An R-matrix  $R \in \mathcal{R}_0$  is said to be diagonal if it is

of the form  $R = DF$ , with  $F$  the flip, and for some orthonormal basis  $\{e_i\}_i$  of  $V$ , the matrix  $D \in \text{End}(V \otimes V)$  is diagonal in the corresponding tensor basis, i.e.

$$D(e_i \otimes e_j) = \lambda_{ij} e_i \otimes e_j, \quad i, j = 1, \dots, d, \quad (6.1)$$

where  $\lambda_{ij} \in \mathbb{C}$ . It is easy to check that such  $R$  solve the Yang-Baxter equation. An R-matrix is said to be of diagonal type if it is equivalent to a diagonal one.

The R-matrix  $R = DF$ ,  $R(e_i \otimes e_j) = \lambda_{ji} e_j \otimes e_i$ , is unitary and involutive if and only if

$$|\lambda_{ij}| = 1, \quad \lambda_{ji} = \lambda_{ij}^{-1}, \quad i, j = 1, \dots, d. \quad (6.2)$$

In particular, we have  $\lambda_{ii} = \pm 1$  for each  $i \in \{1, \dots, d\}$ , and we introduce the parameter  $\ell \in \{0, \dots, d\}$  as the number of  $\lambda_{ii}$ 's that are equal to  $+1$ . This parameter is uniquely fixed by the rank  $r$ , defined as the multiplicity of the eigenvalue  $+1$  of  $R$ . In fact, the trace of  $R$  is

$$2\ell - d = \sum_{i=1}^d \lambda_{ii} = \text{Tr}(R) = 2r - d^2. \quad (6.3)$$

As  $\ell$  ranges over  $\{0, \dots, d\}$ , the rank  $r$  ranges over

$$\frac{1}{2}d(d-1) \leq r \leq \frac{1}{2}d(d+1). \quad (6.4)$$

Thus diagonal involutive R-matrices of dimension  $d$  and rank  $r$  exist if and only if (6.4) is satisfied.

**Proposition 6.1.**

*i) Let  $R \in \mathcal{R}_0$  be of diagonal type, with dimension  $d$ , rank  $r$ , and  $\ell := r - \frac{1}{2}d(d-1)$ . Then*

$$R \sim \bigoplus_{i=1}^{\ell} 1 \oplus \bigoplus_{j=1}^{d-\ell} (-1), \quad (6.5)$$

*and the non-vanishing Thoma parameters of  $R$  are  $\alpha_1 = \alpha_2 = \dots = \alpha_{\ell} = \beta_1 = \dots = \beta_{d-\ell} = d^{-1}$ .*

*ii) Any two involutive R-matrices of diagonal type with the same dimension and rank are equivalent.*

*Proof.* *i)* In the basis defining  $D$ , one has

$$\langle e_i, \text{ptr}(R) e_j \rangle = \sum_k \langle e_i \otimes e_k, D e_k \otimes e_j \rangle = \lambda_{ij} \delta_{ij},$$

which shows  $\text{ptr}(R) = 1_{\ell} \oplus (-1_{d-\ell})$ . The claim now follows from Thm. 3.3 *ii)* and Thm. 4.8 *ii)*.

*ii)* The character depends only on  $d$  and  $\ell$ , and the rank  $r$  determines  $\ell$  uniquely.  $\square$

In terms of diagrams, diagonal R-matrices have the form

$$\left( \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \right), \quad (6.6)$$

with  $\ell$  boxes in the left and  $d - \ell$  boxes in the right column. Yang-Baxter characters of diagonal R-matrices appear in the analysis of the statistics of superselection sectors in quantum field theory [DHR71, Prop. 6.10].

## 6.2 Temperley-Lieb R-matrices

As a second class of examples, we consider solutions coming from representations of the Temperley-Lieb algebra [TL71]. Given an involutive R-matrix  $R \in \mathcal{R}_0$ , we denote its spectral projection onto eigenvalue  $+1$  by  $P$ , i.e.  $P = \frac{1}{2}(R + 1)$ . One computes

$$\frac{1}{8}(R_1 R_2 R_1 - R_2 R_1 R_2) = \left( P_1 P_2 P_1 - \frac{1}{4} P_1 \right) - \left( P_2 P_1 P_2 - \frac{1}{4} P_2 \right), \quad (6.7)$$

and this vanishes by the Yang-Baxter equation. If both terms on the right hand side vanish individually,

$$P_1 P_2 P_1 = \frac{1}{4} P_1, \quad P_2 P_1 P_2 = \frac{1}{4} P_2, \quad (6.8)$$

then  $R$  is said to be of *Temperley-Lieb type*. This terminology is justified by the close relation of (6.8) to the defining relations of the Temperley-Lieb algebra: Recall that given  $q \in \mathbb{C}$ , the Temperley-Lieb algebra  $\mathcal{T}(q)$  is the unital  $*$ -algebra over  $\mathbb{C}$  with generators  $T_k$ ,  $k \in \mathbb{N}$ , and the relations

$$\begin{aligned} T_k^2 &= q T_k, & T_k^* &= T_k, \\ T_k T_m &= T_m T_k, & |k - m| &\geq 2, \\ T_k T_m T_k &= T_k, & |k - m| &= 1. \end{aligned}$$

Given an orthogonal projection  $P \in \text{End}(V \otimes V)$  satisfying (6.8), setting  $T_k := 2P_k$ ,  $k = 1, \dots, n - 1$ , defines a representation of the Temperley-Lieb algebra  $\mathcal{T}(q)$  with  $q = 2$ .

Our equivalence relation  $\sim$  preserves the property of being of Temperley-Lieb type, as follows from the lemma below.

**Lemma 6.2.** *An R-matrix  $R \in \mathcal{R}_0$  is of Temperley-Lieb type if and only if  $\rho_R^{(3)}$  does not contain the trivial representation of  $S_3$ .*

*Proof.* Let  $p_3 \in \mathbb{C}[S_3]$  be the projection given by the trivial representation of  $S_3$ , represented as

$$\rho_R(p_3) = \frac{1}{6}(R_1 R_2 R_1 + R_1 R_2 + R_2 R_1 + R_1 + R_2 + 1). \quad (6.9)$$

Clearly,  $\rho_R^{(3)}$  does not contain the trivial representation if and only if  $\rho_R(p_3) = 0$ . Inserting  $R = 2P - 1$  into (6.9) gives by straightforward calculation

$$\rho_R(p_3) = \frac{4}{3} \left( P_1 P_2 P_1 - \frac{1}{4} P_1 \right) = \frac{4}{3} \left( P_2 P_1 P_2 - \frac{1}{4} P_2 \right).$$

Thus  $\rho_R(p_3) = 0$  is equivalent to the Temperley-Lieb relations (6.8).  $\square$



See Fig. 1 for an example of a Temperley-Lieb type R-matrix where the trivial representation of  $S_3$  does not appear in  $\rho_R^{(3)}$ .

Yang-Baxter representations of the Temperley-Lieb algebra with general parameter  $q$ , i.e., representations in which the tensor structure  $T_k = 1^{\otimes(k-1)} \otimes T \otimes 1 \otimes \dots$  is required for the generators of  $\mathcal{T}(q)$ , have recently been studied by Bytsko. He found various inequalities between  $q$ , the dimension  $d$ , and the rank  $r = \text{Tr}_{V \otimes V}(P)$  that are necessary for such representations to exist [Byt15b, Byt15a].

For the special value  $q = 2$ , we can give a necessary *and sufficient* condition on  $d$  and  $r$  for Yang-Baxter representations of  $\mathcal{T}(2)$  with these parameters to exist, and classify such representations to equivalence.

**Proposition 6.3.**

*i) Yang-Baxter representations of the Temperley-Lieb algebra  $\mathcal{T}(2)$  with rank  $r$  and dimension  $d$  exist if and only if*

$$d^2 - 4r = k^2 \tag{6.10}$$

*for some  $k \in \mathbb{N}_0$ . Two such representations are equivalent if and only if they have the same dimension and rank.*

*ii) Let  $R$  be an involutive R-matrix of Temperley-Lieb type with dimension  $d$  and rank  $r$ . Then its non-vanishing Thoma parameters are*

$$\beta_1 = \frac{1}{2} \left( 1 + \sqrt{1 - \frac{4r}{d^2}} \right), \quad \beta_2 = \frac{1}{2} \left( 1 - \sqrt{1 - \frac{4r}{d^2}} \right). \tag{6.11}$$

*Proof.* We first show the second part *ii*). Let  $R \in \mathcal{R}_0$  have Thoma parameters  $(\alpha, \beta)$ . According to the preceding Lemma, the trivial representation of  $S_3$ , corresponding to the Young diagram  $\square\square\square$ , does not occur in  $\rho_R^{(3)}$  if and only if  $R$  is of Temperley-Lieb type. We can thus conclude from Prop. 5.7 that (equivalence classes of) Temperley-Lieb R-matrices are in one to one correspondence with those  $(\alpha, \beta) \in \mathbb{T}_{\text{YB}}$  that have  $\beta_1, \beta_2$  as their only non-vanishing entries.

We have  $\beta_1 + \beta_2 = 1$ , and on a two-cycle, we get

$$\chi_R(c_2) = -\beta_1^2 - \beta_2^2 = \frac{\text{Tr}_{V \otimes V}(R)}{d^2} = \frac{2r - d^2}{d^2}. \tag{6.12}$$

Solving the resulting quadratic equation proves (6.11).

*i)* A Yang-Baxter representation of  $\mathcal{T}(2)$  of dimension  $d$  and rank  $r$  exists if and only if a Temperley-Lieb R-matrix with the same parameters exists. Since the rescaled Thoma parameters are integers, we know that  $k := d(\beta_1 - \beta_2)$  is an integer. In view of (6.11),  $k = \sqrt{d^2 - 4r}$ . This shows that (6.10) is necessary for the existence of a representation with dimension  $d$  and rank  $r$ .

Conversely, if (6.10) holds for some  $k \in \mathbb{N}_0$ , then (6.11) defines a Temperley-Lieb R-matrix with dimension  $d$  and rank  $r$ . In terms of diagrams,

$$R = \left( \emptyset, \begin{array}{|c|c|c|c|c|c|} \hline \square & \square & \square & \square & \square & \square \\ \hline \end{array} \right), \tag{6.13}$$

consisting of  $d$  boxes distributed over (one or) two rows on the right, with the difference in row lengths equal to  $k$ .

The last statement follows because the Thoma parameters (6.11) depend only on  $d$  and  $r$ .  $\square$

Let us point out that our Temperley-Lieb R-matrices have non-vanishing  $\beta$ -parameters (instead of  $\alpha$ 's) because we required the Temperley-Lieb relation for the spectral projection onto eigenvalue  $+1$ . If we used the spectral projection onto eigenvalue  $-1$  instead,  $\alpha$  and  $\beta$  would be exchanged.

## Acknowledgements

We thank Vaughan Jones for stimulating discussions and for bringing the references [Gur86, Gur91] to our attention. We also thank James Borger for interesting discussions regarding  $\lambda$ -operations on Thoma parameters. Furthermore, the hospitality of the Isaac Newton Institute for Mathematical Sciences during the program ‘‘Operator algebras: subfactors and their applications’’, supported by EPSRC Grant Number EP/K032208/1, is gratefully acknowledged.

## References

- [AAR01] E. Abdalla, M. Abdalla, and K. Rothe. *Non-perturbative methods in two-dimensional quantum field theory*. World Scientific, 2001.
- [AL17] S. Alazzawi and G. Lechner. Inverse Scattering and Locality in Integrable Quantum Field Theories. *Comm. Math. Phys.*, 2017. 10.1007/s00220-017-2891-0.
- [Bax72] R. J. Baxter. Partition Function of the Eight-Vertex Lattice Model. *Ann. Phys.*, 70:193–228, 1972.
- [BO17] A. Borodin and G. Olshanski. *Representations of the Infinite Symmetric Group*. Cambridge University Press, 2017.
- [Byt15a] A. G. Bytsko. Tensor space representations of Temperley–Lieb algebra and generalized permutation matrices. *J. Math. Phys.*, 56(8):083503, 2015.
- [Byt15b] A. G. Bytsko. Tensor space representations of Temperley-Lieb algebra via orthogonal projections of rank  $r \geq 1$ . *J. Math. Phys.*, 56:083502, 2015.
- [CP94] V. Chari and A. Pressley. *A Guide to Quantum Groups*. Cambridge University Press, 1994.
- [DHR71] S. Doplicher, R. Haag, and J. Roberts. Local observables and particle statistics. I. *Comm. Math. Phys.*, 23:199–230, 1971.
- [DVL17] M. Dubois-Violette and G. Landi. Noncommutative products of Euclidean spaces. *Preprint: arXiv 1706.06930v1*, 2017.
- [EI16] T. Enomoto and M. Izumi. Indecomposable characters of infinite dimensional groups associated with operator algebras. *J. Math. Soc. Japan*, 68(3):1231–1270, 2016.
- [GK09] R. Gohm and C. Köstler. Noncommutative Independence from the Braid Group  $B_\infty$ . *Comm. Math. Phys.*, 289(2):435–482, 2009.
- [GK10] R. Gohm and C. Köstler. Noncommutative Independence from Characters of the Infinite Symmetric Group  $S_\infty$ . *Preprint arXiv:1005.5726*, 2010.

- [GK11] R. Gohm and C. Köstler. Noncommutative Independence in the Infinite Braid and Symmetric Group. *Banach Center Publications*, 96:193–206, 2011.
- [Gur86] D. J. Gurevich. The Yang-Baxter equation and a generalization of formal Lie theory. *Sov. Math. Dokl.*, 33(3), 1986.
- [Gur91] D. J. Gurevich. Algebraic aspects of the quantum Yang-Baxter equation. *Leningrad Math. J.*, 2(4), 1991.
- [Hie92] J. Hietarinta. All solutions to the constant quantum Yang-Baxter equation in two dimensions. *Phys. Lett. A*, 165(3):245–251, 1992.
- [Hie93] J. Hietarinta. The upper triangular solutions to the three-state constant quantum Yang-Baxter equation. *J. Phys. A: Math. Gen.*, 26(23):7077, 1993.
- [HL17] S. Hollands and G. Lechner.  $SO(d,1)$ -invariant Yang-Baxter operators and the dS/CFT-correspondence. *to appear in Comm. Math. Phys.*, 2017.
- [Jim89] M. Jimbo. Introduction to the Yang-Baxter equation. *Int. J. Mod. Phys. A*, 4(15):3759–3777, 1989.
- [Jon83] V. F. R. Jones. Index for subfactors. *Invent. Math.*, 72(1):1–25, 1983.
- [Jon87] V. Jones. Hecke algebra representations of braid groups and link polynomials. *Ann. Math.*, 126:335–388, 1987.
- [JS97] V. Jones and V.S. Sunder. *Introduction to Subfactors*. Cambridge University Press, 1997.
- [KL02] L. Kauffman and S. Lomonaco. Quantum entanglement and topological entanglement. *New Journal of Physics*, 4(1):73, 2002.
- [Kös10] C. Köstler. A noncommutative extended de Finetti theorem. *J. Funct. Anal.*, 258(4):1073–1120, 2010.
- [Lyu87] V. V. Lyubashenko. Vectorsymmetries. *Seminar on Supermanifolds*, 14, 1987.
- [LYZ<sup>+</sup>00] J. Lu, M. Yan, Y.-C. Zhu, et al. On the set-theoretical Yang-Baxter equation. *Duke Math. J.*, 104(1):1–18, 2000.
- [Mac95] I. G. Macdonald. *Symmetric Functions and Hall Polynomials*. Oxford Mathematical Monographs. Clarendon Press, Oxford, 2nd edition, 1995.
- [Oko95] A. Okounkov. *On the representations of the infinite symmetric group*. PhD thesis, Moscow State University, 1995. arXiv: math/9803037v1.
- [Ols90] G. Olshanski. Unitary representations of  $(G, K)$ -pairs connected with the infinite symmetric group  $S(\infty)$ . *Leningrad Math. J.*, 1(4):983–1014, 1990.
- [OO98] A. Okounkov and G. Olshanski. Shifted Schur Functions. *St. Petersburg Math. J.*, 9, 1998.
- [Pet83] K. Petersen. *Ergodic theory*. Cambridge University Press, 1983.
- [RS02] M. Rørdam and E. Størmer. *Classification of nuclear  $C^*$ -algebras. Entropy in operator algebras*, volume 126 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2002. Operator Algebras and Non-commutative Geometry, 7.
- [Tho64a] E. Thoma. Die unzerlegbaren, positiv-definiten Klassenfunktionen der abzählbar unendlichen, symmetrischen Gruppe. *Math. Z.*, 85:40–61, 1964.
- [Tho64b] E. Thoma. Über unitäre Darstellungen abzählbarer, diskreter Gruppen. *Math. Ann.*, 153:111–138, 1964.
- [TL71] H. N. V. Temperley and E. H. Lieb. Relations between the 'percolation' and 'colouring' problem and other graph-theoretical problems associated with regular planar lattices: some exact results for the 'percolation' problem. *Proc. Roy. Soc. Lond. A*, 322:251–280, 1971.

- [Tur88] V. G. Turaev. The Yang-Baxter equation and invariants of links. *Invent. Math.*, 92(3):527–553, 1988.
- [TW07] A. S. Toms and W. Winter. Strongly self-absorbing  $C^*$ -algebras. *Trans. Amer. Math. Soc.*, 359(8):3999–4029, 2007.
- [VK81] A. M. Vershik and S. V. Kerov. Characters and factor-representations of the infinite symmetric group. *Sov. Math. Dokl.*, 23(2):389–392, 1981.
- [VK<sup>+</sup>82] A. M. Vershik, S. V. Kerov, et al. Asymptotic theory of characters of the symmetric group. *Funct. Anal. Appl.*, 15:246–255, 1982.
- [VK83] A. M. Vershik and S. V. Kerov. The  $K$ -functor (Grothendieck group) of the infinite symmetric group. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 123:126–151, 1983. Differential geometry, Lie groups and mechanics, V.
- [Was81] A. Wassermann. *Automorphic actions of compact groups on operator algebras*. PhD thesis, University of Pennsylvania, 1981. available from <http://search.proquest.com/docview/303177398>.
- [Wen90] H. Wenzl. Representations of braid groups and the quantum Yang-Baxter equation. *Pac. J. Math.*, 145(1):153–180, 1990.
- [Yam12] M. Yamashita. On subfactors arising from asymptotic representations of symmetric groups. *Proc. Amer. Math. Soc.*, 140:249–261, March 2012.
- [Yan67] C.-N. Yang. Some exact results for the many body problems in one dimension with repulsive delta function interaction. *Phys. Rev. Lett.*, 19:1312–1314, 1967.

Gandalf Lechner: [LechnerG@cardiff.ac.uk](mailto:LechnerG@cardiff.ac.uk)

Ulrich Pennig: [PennigU@cardiff.ac.uk](mailto:PennigU@cardiff.ac.uk)

Simon Wood: [WoodSi@cardiff.ac.uk](mailto:WoodSi@cardiff.ac.uk)