

KNOT THEORY AND CLUSTER ALGEBRAS

VÉRONIQUE BAZIER-MATTE AND RALF SCHIFFLER

ABSTRACT. We establish a connection between knot theory and cluster algebras via representation theory. To every knot diagram (or link diagram), we associate a cluster algebra by constructing a quiver with potential. The rank of the cluster algebra is $2n$, where n is the number of crossing points in the knot diagram. We then construct $2n$ indecomposable modules $T(i)$ over the Jacobian algebra of the quiver with potential. For each $T(i)$, we show that the submodule lattice is isomorphic to the corresponding lattice of Kauffman states. We then give a realization of the Alexander polynomial of the knot as a specialization of the F -polynomial of $T(i)$, for every i . Furthermore, we conjecture that the collection of the $T(i)$ forms a cluster in the cluster algebra whose quiver is isomorphic to the opposite of the initial quiver, and that the resulting cluster automorphism is of order two.

1. INTRODUCTION

We establish a connection from cluster algebras and representation theory to knot theory. Let K be a knot diagram (or link diagram) with n crossings. A segment of K is a segment of the strand from one crossing point to the next. We associate to K a quiver Q with $2n$ vertices, one for each segment of K , as well as a potential W . We then construct $2n$ indecomposable representations $T(i)$, each of which encodes the Alexander polynomial of the link.

To be more precise, each crossing point p of the diagram K gives rise to an oriented cycle w_p of length four in the quiver Q and each region R in K gives rise to an oriented cycle w_R in Q whose length is the number of segments at the region R . The potential W is the difference of the sum of the crossing point cycles and the sum of the region cycles.

Denote by B the Jacobian algebra of the quiver with potential over an algebraically closed field. Then B is a non-commutative algebra which may be infinite dimensional. The representations $T(i)$ are finite-dimensional B -modules. We construct them explicitly as representations of the quiver Q by specifying a finite dimensional vector space at every vertex and a linear map for every arrow in Q . This construction is a representation theoretic analogue of the construction of the Kauffman states in [14]. The direct sum $T = \oplus T(i)$ is called the link diagram module of K .

Let \mathcal{A} be the cluster algebra with principal coefficients of the quiver Q as defined in [12]. \mathcal{A} is a commutative algebra with a special combinatorial structure. It is defined as a subring of a field of rational functions by constructing a set of

The second author was supported by the NSF grants DMS-1800860 and DMS-2054561. This work was partially supported by a grant from the Simons Foundation. The authors would like to thank the Isaac Newton Institute for Mathematical Sciences for support and hospitality during the programme Cluster Algebras and Representation Theory when work on this paper was undertaken. This work was supported by: EPSRC Grant Number EP/R014604/1.

generators, the cluster variables, via a recursive method called mutation that is determined by the quiver Q . Each mutation step connects two sets of $2n$ cluster variables and these sets are called the clusters of \mathcal{A} .

The cluster variables are Laurent polynomials in two sets of indeterminates x_i and y_i , for $i = 1, 2, \dots, 2n$, with positive integer coefficients [11, 16]. Moreover, their specialization, obtained by setting all x_i equal to 1 is a polynomial, called the F -polynomial [12].

It was shown in [9] that F -polynomials can also be computed from modules over our Jacobian algebra B . If M is a B -module then its F -polynomial is

$$F_M = \sum_{\mathbf{e}} \chi(\mathrm{Gr}_{\mathbf{e}}(M)) \mathbf{y}^{\mathbf{e}},$$

where the sum runs over all dimension vectors $\mathbf{e} = (e_i)_{i=1,2,\dots,2n}$ of submodules of M and $\mathbf{y}^{\mathbf{e}} = y_1^{e_1} y_2^{e_2} \dots y_{2n}^{e_{2n}}$. Moreover, $\mathrm{Gr}_{\mathbf{e}}(M)$ is the quiver Grassmannian of M , meaning the variety of all submodules of M whose dimension vector is \mathbf{e} , and χ denotes the Euler characteristic. In general, this Euler characteristic is very hard to compute, because it is known that every projective variety can be realized as a quiver Grassmannian.

We show that the F -polynomials of our B -modules $T(i)$ have a much simpler formula, since every submodule is uniquely determined by its dimension vector. Therefore

$$F_{T(i)} = \sum_{L \subset T(i)} \mathbf{y}^{\dim L},$$

where the sum runs over all submodules L of $T(i)$. We write $F_{T(i)}|_t$ for the specialization of the F -polynomial at

$$y_j = \begin{cases} -t & \text{if segment } j \text{ runs from an undercrossing to an overcrossing;} \\ -t^{-1} & \text{if segment } j \text{ runs from an overcrossing to an undercrossing;} \\ -1 & \text{if segment } j \text{ connects two overcrossings or two undercrossings.} \end{cases} \quad (1)$$

The Alexander polynomial Δ_K of an oriented link diagram K is an important polynomial invariant of the link. It is a Laurent polynomial in one variable t with integer coefficients. Introduced by Alexander in [1], it has several equivalent definitions, see for example [18]. In this paper, we follow Kauffman's approach that realizes the Alexander polynomial as a state sum [14]. More recently, the Alexander polynomial has been generalized in the work of Osváth and Szabó [23], as well as Rasmussen [25], on knot Floer homology.

We are now ready to state our main result. Recall that a link is prime if it cannot be decomposed as the connected sum of two non-trivial links.

Theorem 1.1. *Let K be a diagram of a prime link. Then, for every segment i of K , the Alexander polynomial of K is equal to the specialized F -polynomial of the B -module $T(i)$. That is*

$$\Delta_K = F_{T(i)}|_t.$$

We point out that the quiver Q , and hence the algebra B and the cluster algebra \mathcal{A} , is not an invariant of the link, because Q depends on the choice of the diagram K . For one, the number of vertices in Q is equal to the number of segments in K , which is not invariant under Reidemeister moves. Moreover, the definition of Q

does not take into account the difference between an overpass and an underpass in K . This difference is only recovered in the specialization (1).

The key step in the proof is the following result which is of interest in its own right.

Theorem 1.2. *The lattice of Kauffman states of K relative to a segment i is isomorphic to the lattice of submodules of the B -module $T(i)$.*

An interesting question is how the different $T(i)$ are related to each other. We conjecture the following. Recall that a B -module T is called a tilting module if the projective dimension of T is at most one, $\text{Ext}_B^1(T, T) = 0$ and there is an exact sequence $0 \rightarrow B \rightarrow T^0 \rightarrow T^1$ with T^0, T^1 in the additive closure of T . Also recall that, given a quiver Q , its opposite quiver Q^{op} is obtained by reversing the direction of all arrows.

- Conjecture 1.3.**
- (a) The module $T = \oplus_i T(i)$ is a B -tilting module.
 - (b) The Gabriel quiver of the endomorphism algebra of T is isomorphic to the quiver Q^{op} .
 - (c) The $2n$ cluster variables in \mathcal{A} corresponding to T form a cluster.
 - (d) There is a permutation σ of order two such that the mapping that sends the initial cluster variable x_i to the cluster variable corresponding to $T(\sigma(i))$ is a cluster automorphism of order two in the sense of [2].

Evidence for the conjecture has been obtained in previous work by David Whiting and the second author in [27]. They considered a very special family of links, namely 2-bridge links whose continued fraction has at most two parameters. For a slightly simpler quiver than our quiver Q , they constructed some of the modules $T(i)$ and showed that their direct sum $\oplus T(i)$ can be completed to a tilting module that satisfies the conditions in the conjecture.

As an application, we use a well-known property of the Alexander polynomial to show the following result that is related to the rank-unimodality conjecture of [20].

Theorem 1.4. *Let M be a module of Dynkin type \mathbb{A}_n and \mathcal{L} the submodule lattice of M . Then*

$$\sum_{L \in \mathcal{L}} (-1)^{h(L)} = \begin{cases} \pm 1 & \text{if } |\mathcal{L}| \text{ is odd;} \\ 0 & \text{if } |\mathcal{L}| \text{ is even,} \end{cases}$$

where $h(L) = \dim L = \sum_{j \in Q_0} \dim L_j$ is the total dimension of the submodule L .

Relation to other work. A first connection between cluster algebras and knot theory was given by Kyungyong Lee and the second author in [17] in the special case of 2-bridge links. The authors realized another invariant, the Jones polynomial, as a specialization of a cluster variable in a cluster algebra of Dynkin type \mathbb{A} . This result was based on an ad hoc construction using the fact that both the 2-bridge links and the cluster variables of type \mathbb{A} are parametrized by continued fractions. We now can see this correspondence as a special case of our general construction, as explained in section 8. This provides a more conceptual explanation for the results in [17]. However, we do not know how to generalize the Jones polynomial specialization to arbitrary links.

Nagai and Terashima used ancestral triangles constructed from continued fractions to give a formula for the cluster variables of type \mathbb{A} and then defined a specialization that produces the Alexander polynomial of the corresponding 2-bridge link, see [22]. Our specialization is a generalization of theirs.

In [6], Cohen, Dasbach and Russel gave a realization of the Alexander polynomial for arbitrary knots as a sum over perfect matchings of the bipartite graph whose vertices are given by the crossing points and the regions of the diagram. Their graph can be recovered from our quiver by the methods used for plabic graphs, see for example [10]. In the case of 2-bridge knots, their graph is equivalent to the snake graph associated to the continued fraction in [5] and in that case their formula seems to be a special case of the cluster variable expansion formula of [21] and therefore may be related to ours as well. However, in their approach, the weight of a perfect matching is given by edge weights, which in the cluster algebra setup corresponds to x -variables, whereas we use the y -variables instead. For arbitrary knots, it is unclear if their formula corresponds to a cluster variable.

All of the articles above consider a single segment of the link to produce a formula for the invariant. In our approach, we rather aim at a conceptual understanding of the collection of the $2n$ objects given by all of the segments of the link inside the cluster algebra and in the module category of the Jacobian algebra.

The paper is organized as follows. After fixing the notation and recalling certain facts and terminology in section 2, we review Kauffman's construction of the state poset and the state polynomial in section 3. In section 4, we define our quiver with potential and its Jacobian algebra B . The link module $T = \bigoplus_i T(i)$ is constructed in section 5. Section 6 is devoted to the proof of the lattice isomorphism in Theorem 1.2. Then Theorem 1.1 is proved in section 7. We end the paper with the special case of 2-bridge links and the proof of Theorem 1.4 in section 8.

Acknowledgments. We thank Dylan Rupel for pointing out a typo in an earlier version of this article.

2. PRELIMINARIES

We recall basic notions and results from knot theory and cluster algebras.

2.1. Knots and links. A *knot* is a subset of \mathbb{R}^3 that is homeomorphic to a circle. A *link with r components* is a subset of \mathbb{R}^3 that is homeomorphic to r disjoint circles. Thus a knot is a link with one component. Links are considered up to ambient isotopy. A link is said to be *prime* if it is not the connected sum of two nontrivial links.

A *link diagram* K is a projection of the link into the plane, that is injective except for a finite number of double points that are called *crossing points*. In addition, the diagram carries the information at each crossing point which of the two strands is on top and which is below. A diagram is called *alternating* if traveling along a strand alternates between overcrossings and undercrossings. A link is called *alternating* if it has an alternating diagram. A link is said to be *oriented* if for each component a direction of traveling along the strand is fixed.

A *curl* is a monogon in the diagram. We usually assume without loss of generality that our link diagrams are without curls, because one can always remove them (by a Reidemeister I move) without changing the link.

Throughout this paper, we assume that all links are prime and all link diagrams have a finite number of crossing points.




FIGURE 1. Skein relations for the Alexander polynomial

2.1.1. *The Alexander polynomial.* The Alexander polynomial Δ of an oriented link is a polynomial invariant of the link $\Delta \in \mathbb{Z}[t^{\pm\frac{1}{2}}]$ that can be defined in terms of homology, see [18, Chapter 6]. For the original definition of Alexander, see [1]. The Alexander polynomial is defined up to multiplication by a signed power of t .

In [7], Conway showed that the Alexander polynomial $\Delta(K)$ of an oriented link K can be defined recursively as follows. The Alexander polynomial of the unknot is 1, and whenever three oriented links K_-, K_+ and K_0 are the same except in the neighborhood of a point, where they are as shown in Figure 1, then

$$\Delta_{K_+} - \Delta_{K_-} = (t^{1/2} - t^{-1/2})\Delta_{K_0}.$$

This property also provides a normalization of the Alexander polynomial, but we will not use it here.

The Alexander polynomial Δ has the following properties, see for example [18, Chapter 6].

- (i) For any link, $\Delta(t) \doteq \Delta(t^{-1})$, where the symbol \doteq means “equal up to a signed power of t ”.
- (ii) $\Delta(1) = \pm 1$ for any knot, and $\Delta(1) = 0$ for any link with at least 2 components.
- (iii) For any knot

$$\Delta \doteq a_0 + a_1(t^{-1} + t) + a_2(t^{-2} + t^2) + \dots$$

with a_0 odd.

- (iv) If a knot has genus g then $2g \geq \text{breadth}(\Delta)$, where the breadth is the difference between the maximal and the minimal degree of the polynomial.

Kauffman gave a description of the Alexander polynomial as a state sum. This approach is crucial for us and we review it in section 3.

Let us close this subsection by mentioning a recent breakthrough in a closely related question. In 1982, Freedman showed in [13] that a knot in the 3-sphere is *topologically* slice if its Alexander polynomial is trivial. A famous pair of knots with 11 crossings, the Kinoshita-Terasaka knot and the Conway knot are the smallest non-trivial knots for which the Alexander polynomial is trivial. In particular, both knots are topologically slice. The Kinoshita-Terasaka knot is also known to satisfy the stronger property of being *smoothly* slice. Recently Lisa Piccirillo solved a long-standing open problem in [24] by proving that the Conway knot is *not* a smoothly slice knot. For an illustration of the quiver of the Conway knot see Example 9.3.

2.2. Cluster algebras. In this section, we recall the definition of a skew-symmetric cluster algebra with principal coefficients following [12, 26]

Let \mathbb{P} be the free abelian group on generators y_1, \dots, y_n written multiplicatively. Let $\mathbb{Z}\mathbb{P}$ be the ring of Laurent polynomials in the variables y_1, \dots, y_n and let $\mathbb{Q}\mathbb{P}$ denote its field of fractions. Denote by $\mathcal{F} = \mathbb{Q}\mathbb{P}(x_1, \dots, x_n)$ the field of rational

functions in n variables and coefficients in $\mathbb{Q}\mathbb{P}$. We also define an auxiliary addition \oplus by

$$\prod_j y_j^{a_j} \oplus \prod_j y_j^{b_j} = \prod_j y_j^{\min(a_j, b_j)}. \quad (2)$$

The cluster algebra is determined by the choice of an initial seed $(\mathbf{x}, \mathbf{y}, Q)$, which consists of the following data.

- Q is a finite connected quiver without loops $\circ \curvearrowright$ and 2-cycles $\circ \rightleftarrows \circ$, and with n vertices;
- $\mathbf{y} = (y_1, \dots, y_n)$ is the n -tuple of generators of \mathbb{P} , called *initial coefficient tuple*;
- $\mathbf{x} = (x_1, \dots, x_n)$ is the n -tuple of variables of \mathcal{F} , called *initial cluster*.

The cluster algebra $\mathcal{A} = \mathcal{A}(\mathbf{x}, \mathbf{y}, Q)$ is the $\mathbb{Z}\mathbb{P}$ -subalgebra of \mathcal{F} generated by so-called *cluster variables*, and these cluster variables are constructed from the initial seed by a recursive method called *mutation*. A mutation transforms a seed $(\mathbf{x}, \mathbf{y}, Q)$ into a new seed $(\mathbf{x}', \mathbf{y}', Q')$. Given any seed there are n different mutations μ_1, \dots, μ_n , one for each vertex of the quiver, or equivalently, one for each cluster variable in the cluster.

The *seed mutation* μ_k in direction k transforms $(\mathbf{x}, \mathbf{y}, Q)$ into the seed $\mu_k(\mathbf{x}, \mathbf{y}, Q) = (\mathbf{x}', \mathbf{y}', Q')$ defined as follows:

- \mathbf{x}' is obtained from \mathbf{x} by replacing one cluster variable by a new one, $\mathbf{x}' = \mathbf{x} \setminus \{x_k\} \cup \{x'_k\}$, and x'_k is defined by the following *exchange relation*

$$x_k x'_k = \frac{1}{y_k \oplus 1} \left(y_k \prod_{i \rightarrow k} x_i + \prod_{i \leftarrow k} x_i \right) \quad (3)$$

where the first product runs over all arrows in Q that end in k and the second product runs over all arrows that start in k .

- $\mathbf{y}' = (y'_1, \dots, y'_n)$ is a new coefficient n -uple, where

$$y'_j = \begin{cases} y_k^{-1} & \text{if } j = k; \\ y_j \prod_{k \rightarrow j} y_k (y_k \oplus 1)^{-1} \prod_{k \leftarrow j} (y_k \oplus 1) & \text{if } j \neq k. \end{cases}$$

Note that one of the two products is always trivial, hence equal to 1, since Q has no oriented 2-cycles. Also note that \mathbf{y}' depends only on \mathbf{y} and Q .

- The quiver Q' is obtained from Q in three steps:
 - (1) for every path $i \rightarrow k \rightarrow j$ add one arrow $i \rightarrow j$,
 - (2) reverse all arrows at k ,
 - (3) delete 2-cycles.

Mutations are involutions, that is, $\mu_k \mu_k(\mathbf{x}, \mathbf{y}, Q) = (\mathbf{x}, \mathbf{y}, Q)$. Note that Q' only depends on Q , that \mathbf{y}' depends on \mathbf{y} and Q , and that \mathbf{x}' depends on the whole seed $(\mathbf{x}, \mathbf{y}, Q)$.

Let \mathcal{X} be the set of all cluster variables obtained by finite sequences of mutations from $(\mathbf{x}, \mathbf{y}, Q)$. Then the cluster algebra $\mathcal{A} = \mathcal{A}(\mathbf{x}, \mathbf{y}, Q)$ is the $\mathbb{Z}\mathbb{P}$ -subalgebra of \mathcal{F} generated by \mathcal{X} .

By definition, the elements of \mathcal{A} are polynomials in \mathcal{X} with coefficients in $\mathbb{Z}\mathbb{P}$, so $\mathcal{A} \subset \mathbb{Z}\mathbb{P}[\mathcal{X}]$. On the other hand, $\mathcal{A} \subset \mathcal{F}$, so the elements of \mathcal{A} are also rational functions in x_1, \dots, x_n with coefficients in $\mathbb{Q}\mathbb{P}$.

2.2.1. *Laurent phenomenon and positivity.* We have the following important results.

Theorem 2.1 (Laurent Phenomenon). [11] *Let $u \in \mathcal{X}$ be any cluster variable. Then*

$$u = \frac{f(x_1, \dots, x_n)}{x_1^{d_1} \dots x_n^{d_n}}$$

where $f \in \mathbb{Z}\mathbb{P}[x_1, \dots, x_n]$, $d_i \in \mathbb{Z}$.

Theorem 2.2 (Positivity). [16] *The coefficients of the Laurent polynomials in Theorem 2.1 are positive in the sense that $f \in \mathbb{Z}_{\geq 0}\mathbb{P}[x_1, \dots, x_n]$.*

2.2.2. *F-polynomials.* Let u be any cluster variable in the cluster algebra $\mathcal{A} = \mathcal{A}(\mathbf{x}, \mathbf{y}, Q)$. By the two theorems above, we can write u as a positive Laurent polynomial in the initial cluster as $u = \mathcal{L}_u \in \mathbb{Z}_{\geq 0}[x_1, \dots, x_n, y_1, \dots, y_n]$. Then the *F-polynomial* of x is defined as the evaluation of \mathcal{L}_u at $x_1 = \dots = x_n = 1$. Thus

$$F_x = \mathcal{L}_u(1, \dots, 1, y_1, \dots, y_n).$$

2.3. **Quivers with potential.** In this subsection, we recall an alternative approach to *F-polynomials* using quivers with potential. Let Q be a finite quiver. Following [8] we let the *vertex span* of Q be the commutative algebra R over \mathbb{C} with basis the constant paths e_i , $i \in Q_0$ and multiplication $e_i e_j = \delta_{i,j} e_i$. Furthermore, the *arrow span* of Q is the R -bimodule A with \mathbb{C} -basis the set of arrows Q_1 and R -bimodule structure $e_i A e_j = \bigoplus_{\alpha: j \rightarrow i} \mathbb{C} \alpha$.

The *complete path algebra* of Q then is $R\langle\langle A \rangle\rangle = \prod_{d=0}^{\infty} A^{\otimes R^d}$, with \mathfrak{m} -adic topology given by the two-sided ideal $\mathfrak{m} = \prod_{d=1}^{\infty} A^{\otimes R^d}$. The elements of $R\langle\langle A \rangle\rangle$ are (possibly infinite) \mathbb{C} -linear combinations of paths in Q .

A *potential* $W = \sum_{c \in \mathfrak{m}_{\text{cyc}}} \lambda_c c$ on Q is a (possibly infinite) linear combination of cyclic paths in $R\langle\langle A \rangle\rangle$. The *cyclic derivative* ∂_α , for $\alpha \in Q_1$ is defined on a non-constant cyclic path $\alpha_1 \dots \alpha_d$ by

$$\partial_\alpha(\alpha_1 \dots \alpha_d) = \sum_{p: \alpha_p = \alpha} \alpha_{p+1} \dots \alpha_d \alpha_1 \dots \alpha_{p-1},$$

and extended linearly to the whole potential.

The *Jacobian algebra* $\text{Jac}(Q, W)$ of the quiver with potential is defined as the quotient $R\langle\langle A \rangle\rangle / J(W)$, where $J(W)$ is the closure of the two-sided ideal generated by all cyclic derivatives $\partial_\alpha W$, with $\alpha \in Q_1$.

For every finitely generated module M over the Jacobian algebra, Derksen, Weyman and Zelevinsky introduced its *F-polynomial* in [9] as

$$F_M = \sum_{\mathbf{e}} \chi(\text{Gr}_{\mathbf{e}}(M)) \prod_{i \in Q_0} y_i^{e_i}, \tag{4}$$

where the sum is over all dimension vectors $\mathbf{e} = (e_i)_{i \in Q_0}$ and $\chi(\text{Gr}_{\mathbf{e}}(M)) \in \mathbb{Z}$ is the Euler characteristic of the quiver Grassmannian of all submodules $N \subset M$ of dimension vector \mathbf{e} .

Furthermore, they introduced the notion of mutations of (decorated) representations and showed that if μ is a mutation sequence that transforms the zero module into M then the *F-polynomial* of M is equal to the *F-polynomial* of the cluster variable obtained by the same mutation sequence from the initial seed in the cluster algebra $\mathcal{A}(\mathbf{x}, \mathbf{y}, Q)$.

3. KAUFFMAN STATES

In this section, we recall Kauffman's realization of the Alexander polynomial as a state sum.

3.1. Poset of Kauffman states. Consider an oriented link and fix a diagram K without curls. Denote by n the number of crossings. Then, there are $n + 2$ regions and $2n$ segments. We chose a segment and label it 1 and then, label other segments following the orientation of the string by $2, 3, \dots, 2n$. In this paper, a pair (x, R) of a crossing point x and a region R such that x is incident to R is called an *arrow*.

To define a Kauffman state, we chose a segment $i = 1, 2, \dots, n$ and label the adjacent regions R_i and R'_i . A *Kauffman state* is a set of arrows (x, R) , called *markers*, such that:

- each crossing point is used in exactly one marker;
- each region except for R_i, R'_i is used in exactly one marker.

The regions R_i, R'_i are used in no marker.

A state \mathcal{S}' is obtained from a state \mathcal{S} by a *counterclockwise transposition* at a segment j if \mathcal{S}' is obtained from \mathcal{S} by switching two markers at the segment j as in Figure 2.





FIGURE 2. Kauffman counterclockwise transposition from \mathcal{S} to \mathcal{S}' .

More precisely, let x, y be the endpoints of the segment j and let R_1, R_2 be the adjacent regions at j such that, going clockwise around x , we go from R_1 to R_2 crossing j . Then, \mathcal{S} contains the markers $(x, R_2), (y, R_1)$, \mathcal{S}' contains the markers $(x, R_1), (y, R_2)$ and the other markers in \mathcal{S} and \mathcal{S}' are the same.

We define a partial order on the set of all Kauffman states by $\mathcal{S}_1 < \mathcal{S}_2$ if there is a sequence of counterclockwise transpositions that transforms \mathcal{S}_1 into \mathcal{S}_2 . Kauffman proved that the resulting poset is a lattice whose maximal element is a state that admits no counterclockwise transposition and is therefore called the *clocked state* in [14]. Similarly, the minimal element is called the *counterclocked state* in [14]. We will refer to these states as the *maximal* and the *minimal* state.

Example 3.1. Let's use the following labeling for the segments of the figure-eight knot.








FIGURE 3. Lattice of Kauffman states of the figure-eight knot

Figure 3 shows the lattice of Kauffman states for the figure-eight knot with regards to segment 1.

3.2. The State polynomial. Following Kauffman, we define the *weight* $w(x, R)$ of an arrow (x, R) as shown in the following two cases.



In this case, $w(x, R_1) = B$, $w(x, R_2) = 1$, $w(x, R_3) = W$ and $w(x, R_4) = 1$.



In this case, $w(x, R_1) = W$, $w(x, R_2) = 1$, $w(x, R_3) = B$ and $w(x, R_4) = 1$.

$$\begin{array}{cccc}
\begin{array}{c} \leftarrow \left| \begin{array}{ccc} B & j & 1 \\ 1 & & B \end{array} \right| \rightarrow \\ w(j) = B^{-2} \\ w(j) \mapsto t \end{array} & \begin{array}{c} \leftarrow \left| \begin{array}{ccc} \uparrow W & j & 1 \\ 1 & & W \end{array} \right| \rightarrow \\ w(j) = W^{-2} \\ w(j) \mapsto t^{-1} \end{array} & \begin{array}{c} \leftarrow \left| \begin{array}{ccc} \uparrow W & j & 1 \\ 1 & & B \end{array} \right| \rightarrow \\ w(j) = W^{-1}B^{-1} \\ w(j) \mapsto 1 \end{array} & \begin{array}{c} \leftarrow \left| \begin{array}{ccc} B & j & 1 \\ 1 & & W \end{array} \right| \rightarrow \\ w(j) = W^{-1}B^{-1} \\ w(j) \mapsto 1 \end{array} \\
\begin{array}{c} \leftarrow \left| \begin{array}{ccc} 1 & j & 1 \\ W & & B \end{array} \right| \rightarrow \\ w(j) = WB^{-1} \\ w(j) \mapsto t \end{array} & \begin{array}{c} \leftarrow \left| \begin{array}{ccc} 1 & j & 1 \\ \downarrow B & & W \end{array} \right| \rightarrow \\ w(j) = W^{-1}B \\ w(j) \mapsto t^{-1} \end{array} & \begin{array}{c} \leftarrow \left| \begin{array}{ccc} 1 & j & 1 \\ \downarrow B & & B \end{array} \right| \rightarrow \\ w(j) = 1 \end{array} & \begin{array}{c} \leftarrow \left| \begin{array}{ccc} 1 & j & 1 \\ W & & W \end{array} \right| \rightarrow \\ w(j) = 1 \end{array} \\
\begin{array}{c} \leftarrow \left| \begin{array}{ccc} B & j & W \\ 1 & & 1 \downarrow \end{array} \right| \rightarrow \\ w(j) = WB^{-1} \\ w(j) \mapsto t \end{array} & \begin{array}{c} \leftarrow \left| \begin{array}{ccc} \uparrow W & j & B \\ 1 & & 1 \downarrow \end{array} \right| \rightarrow \\ w(j) = W^{-1}B \\ w(j) \mapsto t^{-1} \end{array} & \begin{array}{c} \leftarrow \left| \begin{array}{ccc} \uparrow W & j & W \\ 1 & & 1 \downarrow \end{array} \right| \rightarrow \\ w(j) = 1 \end{array} & \begin{array}{c} \leftarrow \left| \begin{array}{ccc} B & j & B \\ 1 & & 1 \downarrow \end{array} \right| \rightarrow \\ w(j) = 1 \end{array} \\
\begin{array}{c} \leftarrow \left| \begin{array}{ccc} 1 & j & W \\ W & & 1 \downarrow \end{array} \right| \rightarrow \\ w(j) = W^2 \\ w(j) \mapsto t \end{array} & \begin{array}{c} \leftarrow \left| \begin{array}{ccc} 1 & j & B \\ \downarrow B & & 1 \downarrow \end{array} \right| \rightarrow \\ w(j) = B^2 \\ w(j) \mapsto t^{-1} \end{array} & \begin{array}{c} \leftarrow \left| \begin{array}{ccc} 1 & j & W \\ \downarrow B & & 1 \downarrow \end{array} \right| \rightarrow \\ w(j) = WB \\ w(j) \mapsto 1 \end{array} & \begin{array}{c} \leftarrow \left| \begin{array}{ccc} 1 & j & B \\ W & & 1 \downarrow \end{array} \right| \rightarrow \\ w(j) = WB \\ w(j) \mapsto 1 \end{array}
\end{array}$$

FIGURE 4. Possible values and specializations at $W = t^{\frac{1}{2}}$ and $B = t^{-\frac{1}{2}}$ for the weight of a segment j

The *weight* $w(\mathcal{S})$ of a state \mathcal{S} is defined as

$$w(\mathcal{S}) = \prod_{(x,R) \in \mathcal{S}} w(x,R).$$

The *state polynomial* is the sum of the weights of all states \mathcal{S}

$$\sum_{\mathcal{S}} \sigma(\mathcal{S})w(\mathcal{S}),$$

where $\sigma(\mathcal{S}) = (-1)^b$ with b is the exponent of B in $w(\mathcal{S})$.

Theorem 3.2 ([14]). *The Alexander-Conway polynomial of a diagram L is equal to the specialization of the state polynomial at $W = t^{\frac{1}{2}}$, $B = t^{-\frac{1}{2}}$.*

If a state \mathcal{S}' is obtained from a state \mathcal{S} by a counterclockwise transposition at a segment j , then we denote the *weight ratio* between \mathcal{S}' and \mathcal{S} by $w(j)$. Thus,

$$w(j) = \frac{w(\mathcal{S}')}{w(\mathcal{S})}.$$

Note that $w(j)$ depends only on the segment j and not on the state \mathcal{S} and \mathcal{S}' . The possible values for $w(j)$ and its specialization at $W = t^{\frac{1}{2}}$, $B = t^{-\frac{1}{2}}$ are shown in Figure 4.




FIGURE 5. The quiver of the figure-eight knot of Example 3.1

4. THE JACOBIAN ALGEBRA OF A LINK DIAGRAM

Let K be a reduced diagram of an oriented prime link without curls. Denote by n the number of crossings and label the segments $1, 2, \dots, 2n$ as in section 3.1. We shall use the notation K_0 for the set of crossing points, K_1 for the set of segments, and K_2 for the set of regions (including the unbounded region) of $\mathbb{R}^2 \setminus K$.

We are going to construct a quiver with potential and consider its Jacobian algebra.

We define the quiver Q as follows. The set of vertices Q_0 is the set of segments of K . Thus $Q_0 = K_1$. The set of arrows Q_1 is the set of arrows of K introduced in section 3.1, more precisely, there is an arrow $i \rightarrow j$ in Q if and only if

- the segments i and j of Q meet at a crossing point p ;
- the segments i and j bound the same region R ;
- going clockwise around p , we cross i then R then j .

For example the quiver of the figure-eight knot in Example 3.1 is shown in Figure 5.

The planar link diagram K induces a planar embedding of Q , and since K has no curls, Q has no loops. On the other hand, Q may have 2-cycles, see however section 4.1.

The quiver Q has the following two types of chordless cycles. For each crossing point $p \in K_0$, we obtain a 4-cycle ω_p and for each region R bounded by r segments, we obtain an r -cycle ω_R . Each arrow (p, R) lies in exactly two chordless cycles ω_p and ω_R .

We define a potential W as

$$W = \sum_{p \in K_0} \omega_p - \sum_{R \in K_2} \omega_R.$$

In the example in Figure 5, the potential is

$$W = \alpha_1\alpha_2\alpha_3\alpha_4 + \beta_1\beta_2\beta_3\beta_4 + \gamma_1\gamma_2\gamma_3\gamma_4 + \delta_1\delta_2\delta_3\delta_4 - \alpha_1\delta_2\beta_3 - \alpha_2\gamma_3\delta_1 - \alpha_3\beta_1\gamma_2 - \beta_2\delta_3\gamma_1 - \alpha_4\beta_4 - \gamma_4\delta_4$$

where the first row consists of the four 4-cycles of the four crossing points in K and the second row consists of the four 3-cycles and two 2-cycles of the six regions in K .

Definition 4.1. The algebra $B = \text{Jac}(Q, W)$ is called the (completed) *Jacobian algebra of the link diagram K* .

Remark 4.2. (a) The quiver Q and the algebra B are not invariants of the link. For example, the second Reidemeister move will change the number of vertices in Q .

(b) The quiver does not see the difference between an overpass and an underpass in K .

4.1. Removal of 2-cycles. Each bigon in the link diagram gives rise to a 2-cycle in the quiver. We can replace the quiver with potential (Q, W) by a quiver with potential (Q', W') without 2-cycles as follows. The quiver Q' is obtained from Q by removing all 2-cycles. The potential W' is obtained from W as follows. For every bigon R , given by two segments i, j that cross each other in two crossing points p_1, p_2 , we replace $\omega_{p_1} + \omega_{p_2} - \omega_R$ by the 6-cycle $(\partial_{(p_1, R)}\omega_{p_1})(\partial_{(p_2, R)}\omega_{p_2})$ obtained by joining the two 4-cycles ω_{p_1} and ω_{p_2} . This identification on all 2-cycles induces an isomorphism of algebras

$$B = \text{Jac}(Q, W) \cong \text{Jac}(Q', W').$$

This realization of the algebra B by a quiver without loops and 2-cycles will be important when we describe the connection to cluster algebras.

In our running example, there are two bigons formed by the pairs of segments $(4, 8)$ and $(2, 6)$ in Example 3.1 and these give rise to two 2-cycles $\alpha_4\beta_4$ and $\gamma_4\delta_4$ in the quiver in Figure 5. The above reduction produces the potential

$$\begin{aligned} W' = & \alpha_1\alpha_2\alpha_3\beta_1\beta_2\beta_3 + \gamma_1\gamma_2\gamma_3\delta_1\delta_2\delta_3 \\ & - \alpha_1\delta_2\beta_3 - \alpha_2\gamma_3\delta_1 - \alpha_3\beta_1\gamma_2 - \beta_2\delta_3\gamma_1. \end{aligned}$$

Note that $\partial_{\alpha_4}W = -\beta_4 + \alpha_1\alpha_2\alpha_3$ and thus, in the Jacobian algebra $\text{Jac}(Q, W)$, the arrow β_4 is equal to the path of length three $\alpha_1\alpha_2\alpha_3$. Similarly α_4 is equal to $\beta_1\beta_2\beta_3$.

5. THE LINK DIAGRAM MODULE T

Let K be a curl free diagram of a prime link and let $B = \text{Jac}(Q, W)$ be its Jacobian algebra. In this section, we associate a B -module $T = \bigoplus_{i \in K_1} T(i)$ to K , where each $T(i)$ is an indecomposable summand. The $T(i)$ are constructed explicitly as representations of the quiver Q .

5.1. A partition of K_1 . Let i be a fixed segment of K . We shall define a partition of the set of all segments $K_1 = \sqcup_{d \geq 0} K(d)$ and use it later to define the representation $T(i)$. The sets $K(d)$ depend on the choice of the segment i , but, in the interest of simplicity, our notation does not reflect this dependency. This should not create confusion, since i is fixed here. The construction is recursive and the case $d = 0$ is slightly different from the cases $d > 0$. But first let us run through the construction in the following example.

5.1.1. *An example.* Consider the knot diagram K illustrated in the top picture in Figure 6. This is the knot 10_{66} in the Rolfsen table. We choose the segment $i = 1$. The set $K(0)$ is the set of all edges that share a region with the segment $i = 1$, including i itself. This set is shown in red in the second picture in the figure. Thus we have $K(0) = \{1, 15, 11, 5, 13, 14\}$. We think of this set as a union of two paths both starting and ending with the segment 1. The first path $w_{L,0}$ starts on segment 1 in the direction given by the orientation of the knot and turns left at each crossing point until it comes back to 1. Thus $w_{L,0} = 1, 15, 11, 5, 13, 1$. The second path $w_{R,0}$ also starts on segment 1 in the same direction, but it turns right at each crossing point. Thus $w_{R,0} = 1, 14, 1$.

The set $K(1)$ is constructed in two steps. First, we remove the set $K(0)$ from K , and then we define the set $K'(1)$ as the set of all segments that are incident to the unbounded region of $K \setminus K(0)$. This set is shown in red in the third picture in Figure 6. Note that there are precisely two crossing points p_1 and q_1 that are incident to exactly one segment of $K'(1)$. Again, we can think of this set as the union of two paths, but this time they start at p_1 and end at q_1 . The first path $w_{L,1}$ makes a left turn at every crossing point. Thus $w_{L,1} = 2, 19, 10, 16, 7, 4, 12, 6, 3, 20$. The second path $w_{R,1}$ makes a right turn at every crossing point. Thus $w_{R,1} = 2, 20$.

In our example there are two crossing points x_1 and x_2 that are of degree 4 in $K'(1)$. In this situation, the set $K(1)$ is strictly larger than $K'(1)$. It is shown in the last picture in the figure and is defined as follows. The path $w_{L,1}$ goes through each of the points x_1, x_2 exactly twice. Let $D(x_i)$ denote the domain in the plane bounded by the subpath of $w_{L,1}$ from x_i to x_i . Thus $D(x_1)$ is bounded by the path $19, 10, 16, 7, 4, 12, 6, 3$ and $D(x_2)$ is bounded by the path $4, 12, 6$. The domain $D(x_2)$ actually consist of a single region of the diagram. On the other hand, the domain $D(x_1)$ contains 5 regions of K . We let $R(x_i)$ be the unique region of K inside $D(x_i)$ that is incident to x_i . Then $R(x_2) = D(x_2)$ and $R(x_1)$ is the region bounded by the segments $19, 9, 17, 7, 3$.

Then $K(1)$ is defined as the union of $K'(1)$ with the segments of the regions $R(x_i)$. Thus we need to add the segments 9 and 17 to our set. We are now done with the case $d = 1$.

The set $K(2)$ is again defined in two steps, but the second step will be trivial. First let $K'(2)$ be the set of all segments that are incident to the unbounded region of $K \setminus (K(0) \cup K(1))$. Thus $K'(2) = \{18, 8\}$. There are no crossing points of degree 4 in this set, and therefore we have $K(2) = K'(2)$.

5.1.2. *The general case for $d = 0$.* For a general link, define

$$K'(0) = K(0) = \{j \in K_1 \mid j \text{ and } i \text{ bound the same region of } K\} \cup \{i\},$$

and let $\overline{K'(0)}$ be the closure of $K'(0)$; here closure means that the set also contains the endpoints of the segments.

We can describe $K(0)$ as the union of two paths given by the boundaries of the two regions incident to i . We describe these paths below in a way that will generalize to an iteration of this procedure to $d > 0$. The set $K(0)$ can be described as the union of the segments along two paths

$$\begin{aligned} w_{L,0}: p_0 &\xrightarrow{w_{p,0}} p'_0 \xrightarrow{w'_{L,0}} q'_0 \xrightarrow{w_{q,0}} q_0 \\ w_{R,d}: p_0 &\xrightarrow{w_{p,0}} p'_0 \xrightarrow{w'_{R,0}} q'_0 \xrightarrow{w_{q,0}} q_0 \end{aligned}$$




FIGURE 6. An example of the construction of the partition of the segments of the knot into disjoint subsets $K(d)$. The quiver of this diagram is shown in Figure 11.




FIGURE 7. Proof of Lemma 5.1. At the point p , there is an even number of edges on either side of the curve γ in the two pictures on the left, and an odd number in the picture on the right.

as follows. Let p_0 and q_0 be the endpoints of the segment i . Since K has no curls, we have $p_0 \neq q_0$. Define $p'_0 = q_0$ and $q'_0 = p_0$ and let $w_{p,0} = w_{q,0}$ be the segment i . At p'_0 and at every subsequent crossing point, the path $w_{L,0}$ turns left, and therefore $w_{L,0}$ is the boundary of the region to the left of the segment i from p_0 to q_0 . Similarly $w_{R,0}$ is the boundary of the region to the right of the segment i . These are exactly the segments in $K'(0)$.

Note that the two points p_0, q_0 are of degree 3 in $\overline{K(0)}$ and all other crossing points in K have degree 0 or 2 in $\overline{K(0)}$.

Lemma 5.1. *The two subpaths $w'_{L,0}$ and $w'_{R,0}$ do not share a crossing point besides p'_0 and q'_0 .*

Proof. Let $R_1, R_2 \in K_2$ denote the two regions at i . Suppose there exists $p \in K_0$ such that p is not an endpoint of i and $p \in R_1 \cap R_2$. Then we can draw a closed curve γ from p to p that runs through R_1, R_2 , crosses i once, and does not cross any other segment of K . We consider two cases, depending on the local configuration of the four segments at p relative to γ , see Figure 7.

Suppose first that there is an even number of these four segments on either side of γ . This case is illustrated in the two pictures on the left of the figure. Note that at either endpoint of the segment i , there are three loose segments, so that in total there is an odd number of loose segments on either side of γ , and (using the Jordan curve theorem) it is thus impossible to pair them up without crossing γ in order to form a link.

Therefore, out of the four segments at p , there must be an odd number on either side of γ . This case is illustrated in the right picture of the figure. Then, on one side, there is only one segment; call it j . Moving γ slightly away from p toward the segment j , we obtain a closed curve γ' that crosses only two segments i and j . This shows that K is the connected sum of two links and thus not prime, a contradiction. \square

5.1.3. *The general case for $d \geq 1$.* We define $K(d)$ recursively.

Definition 5.2. Assume $K(e)$ is defined for all $e < d$. Let $K'(d)$ be the set of all segments j in $K_1 \setminus (\cup_{e < d} K(e))$ for which there exists a segment $k \in K(d-1)$ such that j and k bound the same region of K . Let $\overline{K'(d)}$ be the closure of $K'(d)$.

Definition 5.3. (a) A crossing point $p \in K_0$ is called an *external point in $K'(d)$* if exactly one of its incident segments lies in $K'(d)$. A segment $j \in K_1$ is called




FIGURE 8. The domain $D(x_1)$ and the region $R(x_1)$ associated to an interior point $x_1 \in K(d)$. The domain $D(x_1)$ is bounded by the red curve and the region $R(x_1)$ is shaded. It is bounded above by the black segments, one of which is labeled j_1 . The domain $D(x_2)$ for the second interior point x_2 is only the part of $D(x_1)$ that lies above the point x_2 .

external in $K'(d)$ if $j \in K'(d)$ and exactly one the endpoints of j is an external point in $K'(d)$.

(b) A crossing point p is called an *internal point* in $K'(d)$ if all four segments at p lie in $K'(d)$. Note that for $d = 0$ there are no internal crossing points since our diagram K has no curls. If there exists a non-constant path w starting and ending at x that uses only segments of $K'(d)$, we let $D(x)$ be the bounded domain enclosed by w in the plane. Then $D(x)$ is a union of regions of $\overline{K'(d)}$. Let $R(x) \in K_2$ be the unique region in $D(x)$ that contains x .

An example is shown in Figure 8. In that figure, there are four interior points x_1, \dots, x_4 . The domain $D(x_1)$ is bounded by the red subcurve w and the region $R(x_1)$ is shaded.

Now let j be a segment of K . We define $\epsilon_d(j) \in \{0, 1\}$ for $d \geq 1$ as follows.

$$\epsilon_d(j) = \begin{cases} 1 & \text{if there exists an internal point } x \text{ in } K'(d) \text{ such that } j \text{ lies} \\ & \text{in the interior of } D(x) \text{ and the region } R(x) \in K_2 \text{ contains the} \\ & \text{segment } j \text{ and the point } x; \\ 0 & \text{otherwise.} \end{cases} \quad (5)$$

Definition 5.4. For $d \geq 1$, let

$$K(d) = K'(d) \cup \{j \in K_1 \mid \epsilon_d(j) = 1\}.$$

For example, the segment j_1 in Figure 8 satisfies the first condition for $x = x_1$. Thus $\epsilon_d(j_1) = 1$ in this case. The set $K(d)$ contains every segment of the red curve and every segment of the black curves bounding $R(x_1)$ and $R(x_2)$.

Lemma 5.5. *Let $d \geq 1$.*

(a) *Each connected component C of $\overline{K'(d)}$ is either a single path w from p_d to q_d or the union of two paths*

$$\begin{aligned} w_{L,d}: p_d &\xrightarrow{w_{p,d}} p'_d \xrightarrow{w'_{L,d}} q'_d \xrightarrow{w_{q,d}} q_d \\ w_{R,d}: p_d &\xrightarrow{w_{p,d}} p'_d \xrightarrow{w'_{R,d}} q'_d \xrightarrow{w_{q,d}} q_d \end{aligned}$$

from p_d to q_d , where p_d and q_d are external points in $K'(d)$, the initial and terminal subpaths $w_{p,d}$ and $w_{q,d}$ are the same in both paths, $\deg p'_d = \deg q'_d = 3$ in C , $p'_d \neq q'_d$ and $w'_{L,d}$ (respectively $w'_{R,d}$) is obtained by turning left (respectively right) at every crossing point in $K'(d)$.

If no such points p'_d, q'_d exist then $w_{L,d} = w_{R,d}$ and $K'(d)$ is a single path from p_d to q_d .

(b) *All other crossing points, besides p_d, p'_d, q_d, q'_d , have degree 0, 2 or 4 in C . In particular, C has exactly two external points p_d and q_d , and moreover, $p_d, q_d \in \overline{K'(d)} \cap \overline{K'(d-1)}$ and $p_d, q_d \notin \overline{K'(e)}$, with $e \neq d, d-1$.*

(c) *The paths $w'_{L,d}$ and $w'_{R,d}$ do not share a crossing point besides p'_d and q'_d .*

Proof. (a) Suppose first $d = 1$. The external points are $p_1 = p'_0$ and $q_1 = q'_0$, and there are no other external points in $K'(1)$. If there are points of degree 3 in $\overline{K'(1)}$, we let p'_1 be the point of degree 3 closest to p_1 , and let q'_1 be the point of degree 3 closest to q_1 . Note that every connected component has an even number of points of degree 3, because of parity, and that there are at most two because there are only two external points.

Let $w_{p,1}$ be the unique path from p_1 to p'_1 in $K'(1)$ that

- does not use an edge twice;
 - is of maximal length;
 - turns left or right at every point of degree 4 in $\overline{K'(d)}$.
- (6)

Such a path exists and is unique by the following argument. The starting point p_1 is of degree 1 in C , so the first step is uniquely determined. At every point of degree 2, the incoming edge leaves only one choice for the outgoing edge. At a point of degree 4, there are a priori two possibilities, turn left or turn right, but only one of these will produce a path of maximal length. Similarly, Let $w_{q,1}$ be the unique path from q'_1 to q_1 in $K'(1)$ that respects conditions 6.

Then the paths $w_{L,1}, w_{R,1}$ form the boundary of the regions on the left and right of the path $(w_{q,1} \cup w_{p,1})$ in $(K \setminus K(0)) \cup \{i\}$. These are exactly the segments in $K'(1)$. This completes the proof of (a) for $d = 1$.

(b) The degree formulas follow directly from (a). In particular p_1 and q_1 are the only external points in C . Furthermore, three of the segments at p_1 lie in $K(0)$, and the remaining segment, which is the first segment of $w_{p,1}$, lies in $K'(1)$. Thus




FIGURE 9. Proof of Lemma 5.5 part (c).

$p_0 \in \overline{K'(1)} \cap \overline{K'(0)}$ and $p_0 \notin K'(e)$, with $e \neq 0, 1$. The proof for q_1 is similar. This also implies that all other points have degree 0, 2 or 4 in $\overline{K'(1)}$.

(c) Suppose $w'_{L,1}$ and $w'_{R,1}$ share a point $x \neq p'_1, q'_1$, see Figure 9. Let D denote the domain in the plane bounded by the segments of the paths $w_{L,1}$ and $w_{R,1}$ from p'_1 to x . Let ℓ denote the segment at p'_1 that does not belong to $w_{L,1}$ or $w_{R,1}$. Since $w_{L,1}$ turns left at p'_1 and $w_{R,1}$ turns right, the segment ℓ lies in D . Following the link, starting at p'_1 in direction of ℓ , we must reach a point y where we leave D . Let ℓ' denote the segment outside D right after y . In K , this segment ℓ' bounds the same region as a segment on the path $w'_{R,1}$, and therefore ℓ' must lie in $K(0) \cup K(1)$. However, ℓ' cannot lie in $K(0)$, because $K(0)$ has no external segments. On the other hand, ℓ' cannot be in $K'(1)$, because (a) implies that every segment of $K'(1)$ lies on one of the two paths $w_{L,1}, w_{R,1}$. This is a contradiction, and thus the two paths $w'_{L,1}, w'_{R,1}$ cannot have the point x in common. This completes the proof for $d = 1$.

For $d > 1$, the proof is similar, with the additional feature that now we may have components that arise from internal points in $K'(d-1)$. Indeed, every internal point x of $K'(d-1)$, such that there is a segment j bounding the region $R(x)$ with $\epsilon_{d-1}(j) = 1$, determines two crossing points $p'_d(x)$ and $q'_d(x)$ as the unique points in $R(x) \setminus \{x\}$ that lie on the boundary of $D(x)$, and such that the path $w = w_{L,d-1}$ or $w_{R,d-1}$ is of the form

$$w: p_d \text{ --- } x \text{ --- } p'_d(x) \text{ --- } q'_d(x) \text{ --- } x \text{ --- } q_d,$$

see Figure 8. Therefore every internal point x of $K'(d-1)$ gives rise to a connected component of $K(d)$ in which the points $p_d = p'_{d-1}(x)$ and $q_d = q'_{d-1}(x)$ are the unique external points. For all other components, we have $p_d = p'_{d-1}$ and $q_d = q'_{d-1}$. Note that p_d and q_d have degree 3 in $\overline{K'(d-1)}$ and thus degree 1 in $\overline{K'(d)}$. Hence they are external points.

The rest of the proof of (a) is analogous to the case $d = 1$. The point p'_d is the point of degree 3 closest to p_d , and q'_d is the point of degree 3 closest to q_d . The paths $w_{p,d}$ and $w_{q,d}$ are the unique paths from p_d to p'_d and from q_d to q'_d that satisfy the conditions (6). Moreover the paths $w_{L,d}, w_{R,d}$ form the boundaries of the regions to the left and right of the path

$$w_{q,d} \dots w_{q,2} w_{q,1} \overset{i}{w_{p,1}} w_{p,2} \dots w_{p,d}$$

in $(K \setminus \cup_{e=0}^{d-1} K(e)) \cup \{w_{q,d-1}, \dots, w_{q,1}, i, w_{p,1}, \dots, w_{p,d-1}\}$. Thus $w_{L,d}$ and $w_{R,d}$ consist of the edges of $K'(d)$.

The proof of (b) is analogous to the case $d = 1$ except that the point p_d may be equal to a point $p'_{d-1}(x)$ for some interior point x in $K'(d)$.

In the proof of (c), the only difference to the proof in case $d = 1$ is that now the segment ℓ' in Figure 9 cannot lie in $K(d-1)$, because otherwise y would be an external point of $K(d-1)$ that lies in $\overline{K(d)}$, a contradiction to (b). \square

Remark 5.6. Each interior point x of $K'(d-1)$ whose region $R(x)$ contains a segment j with $\epsilon_{d-1}(j) = 1$ gives rise to a connected component of $K'(d)$.

Proposition 5.7. *For every segment $i \in K_1$, we have constructed a partition*

$$K_1 = \sqcup_{d \geq 0} K(d).$$

Proof. This follows directly from the construction. \square

We are now ready to define the dimension vector of the representation $T(i)$.

Definition 5.8. Let $K_1 = \sqcup_{d \geq 0} K(d)$ be the partition with respect to a segment $i \in K_1$. For every segment $j \in K_1$, we define $d(i)_j = d$ if $j \in K(d)$.

In the example of Figure 6, we have $d(1)_j = 1$ if $j = 2, 3, 4, 6, 7, 9, 10, 12, 16, 17, 19, 20$; $d(1)_j = 2$ for $j = 8, 18$, and $d(1)_j = 0$ for all other j .

Our next result says that the dimension difference at adjacent vertices is at most one.

Proposition 5.9. *Let $i, j, k \in K_1$. If there is an arrow $j \rightarrow k \in Q$ then*

$$|d(i)_j - d(i)_k| \leq 1.$$

Proof. Let $d = d(i)_j$. Thus $j \in K(d)$. The existence of the arrow $j \rightarrow k$ implies that j and k bound the same region in K . If $k \notin \cup_{e \leq d} K(e)$ then Definition 5.2 implies that $k \in K'(d+1)$, thus $k \in K(d+1)$ and $d(i)_j - d(i)_k = -1$. If $k \in K(d) \cup K(d-1)$ then $d(i)_k \in \{d, d-1\}$ and there is nothing to show. Finally suppose $k \in K(e)$, with $e \leq d-2$. Then Definition 5.2 implies that j lies in $K'(e+1)$ unless it already lies in $\cup_{e' < e} K(e')$. In both cases, we have $d(i)_j \leq e+1 \leq d-1$, a contradiction. \square

5.2. Properties of $K(d)$. It will be convenient to use the following terminology. Given two segments $i, j \in K_1$, a curve in \mathbb{R}^2 is called a *dimension curve* from j to i if it starts at a point on segment j , ends at a point on segment i and does not go through a crossing point of K .

Let $\dim^\circ(i, j)$ be the minimal number of crossings between the segments of K and a dimension curve from segment j to segment i . We call a segment $j \in K_1$ an *interior segment of $K(d)$* if $K(d)$ contains all the segments on the boundary of the two regions incident to j in K .

Note that the segment i is an interior segment of $K(0)$. The following lemma says that there are no other interior segments.

Lemma 5.10. *If $d \geq 1$ then $K(d)$ has no interior segments.*

Proof. Suppose a segment j belongs to two regions R_1, R_2 in K and each segment in $R_1 \cup R_2$ lies in $K(d)$, see Figure 10. Since $d \geq 1$, the dimension curve of j must cross a segment k of $R_1 \cup R_2$, so it has one more crossing than the dimension curve of the segment k . Since $j, k \in K(d)$ this means that $\epsilon_d(j) = 1$. Thus there exists an internal point x in $K(d)$ satisfying the condition (5). In particular, one of the two regions at j , say R_1 , contains x and j . Thus R_1 is the region $R(x)$ of condition (5). The other region of K at j is the region R_2 and both lie entirely in $K(d)$. Since $j \notin K'(d)$, the segments of $R_1 \cup R_2$ that do lie in $K'(d)$ all lie in the same region in




FIGURE 10. Proof of Lemma 5.10

$K'(d)$. In particular, no segment k of $R_1 \cup R_2$ shares a region with another internal point $x' \neq x$ such that k lies in $D(x')$. Therefore each segment k of $R_2 \setminus \{j\}$ satisfies $\epsilon_d(k) = 0$. Thus $k \in K'(d)$, which implies $k \in D(x)$.

Now consider an endpoint y of j . Three of the segments incident to y lie in $R_1 \cup R_2$ and the fourth segment ℓ doesn't, see Figure 10. Therefore ℓ either lies on the boundary of $D(x)$ and hence $\dim^\circ(i, \ell) = d$, or ℓ lies outside $D(x)$ and hence $\dim^\circ(i, \ell) = d - 1$. The latter case is impossible by Lemma 5.5(b), because ℓ would be an external segment of $K'(d - 1)$, but ℓ has an endpoint in $\overline{K'(d)}$ (and not in $\overline{K'(d - 2)}$).

Thus the segment ℓ lies on the boundary of $D(x)$. Since ℓ is not in R_1 , there exists a point $y' \in K_0$ such that two of its incident segments lie in R_1 and one, call it h , lies on the boundary of $D(x)$ between x and y as in the figure. Denote by ℓ' the fourth segment at y' . Note that it must lie inside $D(x)$, because otherwise y' would be an external point of $K'(d - 1)$ that does not lie in $\overline{K'(d - 2)}$, contradicting Lemma 5.5(b). Since R_1 is a region in K , the segment ℓ must lie in the connected component C of $D(x) \setminus R_1$ that contains y' . Following the link starting at y' in direction ℓ' , we must reach a point y'' , where we leave the component C . Let ℓ'' be the segment outside C right after y'' . Then ℓ'' is an external segment of $K'(d - 1)$ with external point y'' not in $\overline{K(d - 2)}$, again a contradiction to Lemma 5.5(b). \square

It will be convenient to consider the following dual graph. For an illustration, see Example 5.16

Definition 5.11. (a) Let $Q(d)$ be the full subquiver of Q on the vertices j such that $j \in K(d)$.

(b) Let $G(d)$ be the graph with vertex set the set of chordless cycles in Q that also lie in $Q(d)$, and two chordless cycles are connected in $G(d)$ by an edge if they share an arrow in $Q(d)$.

Notice that the graph $G(d)$ has two types of vertices, the crossing point vertices and the region vertices. The first type corresponds to the chordless 4-cycles ω_p , with $p \in K_0$, and the second type corresponds to the chordless cycles ω_R , with $R \in K_2$.

Corollary 5.12. *In $G(d)$, the degree of a crossing point vertex is at most 2.*

Proof. If a crossing point vertex x has three adjacent regions in $G(d)$, hence in $K(d)$, then $K(d)$ has an interior segment, contradicting Lemma 5.10. \square

Definition 5.13. Let R be a region of K such that each segment of R lies in $K(d)$. By Lemma 5.5, one of the two paths $w = w_{L,d}$ or $w_{R,d}$ encloses the region R . Let $x(R)$ be the first crossing point of the region R on the path w . We call $x(R)$ the *root* of the region R .

Recall that a *leaf* in a graph is a vertex of degree one.

Lemma 5.14. (a) *The mapping $R \mapsto x(R)$ is a bijection between the sets of region vertices of $G(d)$ and crossing point vertices of $G(d)$.*

(b) *Every connected component of $G(d)$ has a unique vertex that is a leaf and a crossing point vertex.*

Proof. (a) Since the path w starts and ends at vertices outside R it must go through the point $x(R)$ twice, in the sense that it contains all four segments at $x(R)$. Thus $x(R)$ is an internal point of $K'(d)$ and therefore a vertex of $G(d)$. This shows that the mapping is well-defined.

The mapping is injective by definition. Now let x be any crossing point vertex in $G(d)$. Then all four segments at x lie in $K(d)$. This implies that x is an internal point of $K'(d)$, because the endpoints of the segments j with $\epsilon_d(j)$ have at most degree 3 in $K(d)$. Lemma 5.5 then implies that the four segments at x all lie on one path $w = w_{L,d}$ or $w = w_{R,d}$, and thus w goes through x twice. Then w is of the form

$$w: p_d \text{ --- } x \text{ ---}_{w(x)} \text{ --- } x \text{ --- } q_d$$

and the subpath $w(x)$ forms the boundary of the domain $D(x)$ in condition (5). There is a unique region $R(x)$ of K that lies within $D(x)$ and contains x . By definition of $K(d)$, all segments of R lie in $K(d)$. Thus R is a region vertex of $G(d)$ and $x(R) = x$. This shows that the mapping is surjective.

(b) Let C be a connected component of $G(d)$. By Lemma 5.5, there is a path $w = w_{L,d}$ or $w = w_{R,d}$ that encloses all regions of the region vertices of C . Let x be the first point on w that is a crossing point vertex of C . Then x is a leaf of C .

To show that there is no other crossing point that is a leaf, note first that every crossing point vertex of $G(d)$ is of degree at most 2 in $G(d)$, by Corollary 5.12. Now we proceed by induction on the number of region vertices. If there is only one region vertex R in C then $C = R \text{ --- } x(R)$ and we are done. Suppose there is more than one region vertex. Take a leaf ℓ . If $\ell = R$ is a region vertex, then $C \setminus \{\ell\}$ has $x(R)$ as a leaf and thus $C \setminus \{\ell, x(R)\}$ is connected, and by induction it has no other crossing point vertex that is a leaf than x . On the other hand, if C contains no leaf that is a region vertex, then there are more crossing point vertices than region vertices in C , which is impossible by part (a). \square

Remark 5.15. We don't know if $G(d)$ is a forest.

5.3. Definition of the link diagram module T . Let \mathbb{k} be an algebraically closed field. Let K be an oriented diagram without curls of a prime link with n crossings. Let (Q, W) be the associated quiver with potential and $B = \text{Jac}(Q, W)$ its Jacobian algebra. Let I_d denote the identity matrix of rank d . We define the *link diagram module*

$$T = \bigoplus_{i \in K_1} T(i)$$

of K as follows.

For each segment i of K , the direct summand $T(i) = (T(i)_j, T(i)_\alpha)_{j \in Q_0, \alpha \in Q_1}$ is the representation of Q given by

$$T(i)_j = \mathbb{k}^{d(i)_j},$$

for each vertex j , where $d(i)_j$ is the dimension defined in Definition 5.8; and for each arrow $\alpha: j \rightarrow \ell$, we define the corresponding linear map

$$T(i)_\alpha = \begin{cases} \left(\begin{array}{c|c} 0 & \\ \vdots & I_{d-1} \\ 0 & \end{array} \right) & \text{if } d(i)_j = d(i)_\ell + 1 = d \\ \left(\begin{array}{c} I_{d-1} \\ \hline 0 \quad \dots \quad 0 \end{array} \right) & \text{if } d(i)_j + 1 = d(i)_\ell = d \end{cases}$$

and if $d(i)_j = d(i)_\ell = d$ then

$$T(i)_\alpha = \begin{cases} \left(\begin{array}{c|c} 0 & \\ \vdots & I_{d-1} \\ 0 & \hline 0 & 0 \quad \dots \quad 0 \end{array} \right) & \text{if } \alpha \text{ is of the form } (x, R(x)) \text{ with} \\ & x \text{ an internal point of } K'(d) \\ I_d & \text{otherwise.} \end{cases}$$

Because of Proposition 5.9, there are no other possibilities for the dimensions and thus $T(i)_\alpha$ is well-defined.

Example 5.16. The quiver Q of the knot 10_{66} in Figure 6 is shown in the top picture and the representation $T(1)$ in the bottom picture of Figure 11. The quiver $Q(1)$ is the full subquiver on the vertices $2, 3, 4, 6, 7, 9, 10, 12, 16, 17, 19, 20$. It contains four chordless cycles, the two crossing point cycles w_{x_1}, w_{x_2} the two region cycles $w_{R(x_1)}, w_{R(x_2)}$, where we use the notation of Figure 6. Therefore its dual graph is

$$w_{x_1} \text{ --- } w_{R(x_1)} \text{ --- } w_{x_2} \text{ --- } w_{R(x_2)} .$$

6. KAUFFMAN STATES AND SUBMODULES OF THE LINK DIAGRAM MODULE

We keep the notation of the previous sections. Again we choose a segment $i \in K_1$ and consider the Kauffman states and the B -module $T(i)$. Our goal is now to prove that the lattice of Kauffman states of a link K relative to a segment i is isomorphic to the lattice of submodules of the direct summand $T(i)$ of the corresponding link diagram module T .

6.1. The state module $M(\mathcal{S})$. Let \mathcal{S} be a Kauffman state. We will define a B -module $M(\mathcal{S}) = (M_j, M_\alpha)_{j \in Q_0, \alpha \in Q_1}$. Consider a sequence \mathfrak{s} of counterclockwise transpositions that transforms the minimal Kauffman state into the state \mathcal{S} . Then we define

$$M_j = \mathbb{k}^{e_j},$$





FIGURE 11. The quiver Q and the representation $T(1)$ for the knot diagram of Figure 6.

where e_j is the number of occurrences of j in \mathfrak{s} . The order in which the transpositions at j occur determines a basis for M_j , which we call the *basis induced by* \mathfrak{s} .

Next we define the linear maps of the representation M . In the remainder of this section, we use the following matrices


$$J_\ell = \left(\begin{array}{c|ccc} 0 & & & \\ \vdots & & I_\ell & \\ 0 & & & \\ \hline 0 & 0 & \cdots & 0 \end{array} \right) \quad V_\ell = \left(\begin{array}{c|ccc} 0 & & & \\ \vdots & & I_\ell & \\ 0 & & & \end{array} \right) \quad H_\ell = \left(\begin{array}{ccc} & I_\ell & \\ \hline 0 & \cdots & 0 \end{array} \right)$$

where I_ℓ denotes the identity matrix of size ℓ . We point out that J_ℓ is a Jordan block of size $\ell + 1$ with eigenvalue 0, and that $H_\ell V_\ell = J_\ell$ and $V_\ell H_\ell = J_{\ell-1}$.

Every crossing point $p \in K_0$ induces a subsequence $\mathfrak{s}(p)$ of \mathfrak{s} consisting of all occurrences of the transpositions at the four segments incident to p . Let a, b, c, d denote these four segments in counterclockwise order around p such that a is the first entry in $\mathfrak{s}(p)$. Then $\mathfrak{s}(p)$ is of one of the following forms

$$(abcd)^\ell, (abcd)^\ell a, (abcd)^\ell ab, (abcd)^\ell abc, \quad (7)$$

for some $\ell \geq 0$. Let



be the corresponding 4-cycle in the quiver Q , and let $w_p = \delta\gamma\beta\alpha \in B$.

Since every arrow of Q lies in a unique 4-cycle induced by a crossing point, it suffices to define the linear maps of the representation M on these four arrows $\alpha, \beta, \gamma, \delta$. There are four cases depending on the sequence $\mathfrak{s}(p)$.

(i) If $\mathfrak{s}(p) = (abcd)^\ell$ then $e_a = e_b = e_c = e_d = \ell$ and

$$M_\delta = J_{\ell-1} \quad M_\gamma = M_\beta = M_\alpha = I_\ell.$$

(ii) If $\mathfrak{s}(p) = (abcd)^\ell a$ then $e_a = \ell + 1, e_b = e_c = e_d = \ell$ and

$$M_\delta = V_\ell \quad M_\gamma = M_\beta = I_\ell, \quad M_\alpha = H_\ell$$

(iii) If $\mathfrak{s}(p) = (abcd)^\ell ab$ then $e_a = e_b = \ell + 1, e_c = e_d = \ell$ and

$$M_\delta = V_\ell \quad M_\gamma = I_\ell, \quad M_\beta = H_\ell \quad M_\alpha = I_{\ell+1}.$$

(iv) If $\mathfrak{s}(p) = (abcd)^\ell abc$ then $e_a = e_b = e_c = \ell + 1, e_d = \ell$ and

$$M_\delta = V_\ell \quad M_\gamma = H_\ell \quad M_\beta = M_\alpha = I_{\ell+1}.$$

Definition 6.1. The B -module $M(\mathcal{S})$ is called the *state module* associated to the Kauffman state \mathcal{S} .

Remark 6.2. In all four cases (i)-(iv) above the composition of the four matrices along the cycle w_p is equal to $J_{\ell-1}$. Thus the action of w_p on $M(\mathcal{S})$ is given by this matrix.

From the construction of the state module, we have the following results.

Lemma 6.3. *Let $M(\mathcal{S}) = (M_x, M_\alpha)_{x \in Q_0, \alpha \in Q_1}$. Then for every arrow $\alpha: j \rightarrow k$, we have $|\dim M_j - \dim M_k| \leq 1$.*

Lemma 6.4. *If the state \mathcal{S}' is obtained from the state \mathcal{S} by applying the transposition at a segment a then the module $M(\mathcal{S}')$ is obtained from $M(\mathcal{S})$ by*

- (i) *increasing the dimension at vertex a by one;*
- (ii) *increasing the rank of the map on each arrow $\alpha: a \rightarrow \bullet$ starting at a by one.*

The dimension at the other vertices and the rank on the other arrows do not change.

We also note the following for future reference.

Lemma 6.5. *Let \mathcal{S} be a state and \mathfrak{s} a sequence of transpositions that transforms the minimal state into \mathcal{S} . If*

$$w = a_0 \xrightarrow{\alpha_1} a_1 \xrightarrow{\alpha_2} \dots \xrightarrow{\alpha_{t-1}} a_t \xrightarrow{\alpha_t} a_0$$

is a chordless cycle in Q then the subsequence of \mathfrak{s} of all occurrences of transpositions at vertices of w is of the form

$$a_j \dots a_2 a_1 a_0 (a_t a_{t-1} \dots a_1 a_0)^\ell a_t a_{t-1} \dots a_k$$

for some j, k and ℓ . In particular, the order in \mathfrak{s} is opposite to the order in w .

Proof. We have already proved this result in equation (7) in the case where $w = w_p$ is the chordless 4-cycle given by a crossing point $p \in K_0$. It suffices to show the result in the case where $w = w_R$ is the chordless cycle of a region $R \in K_2$. The transposition at a_i is defined by moving the markers counterclockwise around the endpoints of a_i , but it can also be seen as moving the markers in the clockwise direction along the segment a_i , see Figure 2. Thus the proof for the region cycle w_R is dual to the proof for the crossing point cycle w_p . \square

As an immediate consequence we have the following.

Corollary 6.6. *If $\alpha: a \rightarrow d$ is an arrow in Q such that the transpositions at a and d occur consecutively in \mathfrak{s} then d comes before a .* \square

6.2. Lattice isomorphism. We start with the following result.

Proposition 6.7. *Let K be a link diagram without curls and let $i \in K_1$ be a segment. Let $\mathcal{S}, \mathcal{S}'$ be two Kauffman states relative to i . Then*

- (a) *$M(\mathcal{S})$ is a B -module.*
- (b) *$M(\mathcal{S}) \not\cong M(\mathcal{S}')$ if $\mathcal{S} \neq \mathcal{S}'$.*
- (c) *If \mathcal{S} is the minimal Kauffman state then $M(\mathcal{S}) = 0$.*
- (d) *If \mathcal{S} is the maximal Kauffman state then $M(\mathcal{S}) = T(i)$.*
- (e) *If $\mathcal{S} < \mathcal{S}'$ then $M(\mathcal{S})$ is a submodule of $M(\mathcal{S}')$.*
- (f) *For every submodule M of $T(i)$ there is a unique Kauffman state \mathcal{S} such that $M \cong M(\mathcal{S})$.*

Proof. (a) By definition, $M(\mathcal{S})$ is a representation of Q , so we only need to check that $M(\mathcal{S})$ satisfies the relations given by the cyclic derivatives of the potential W .

Let (p, R) be an arrow in Q , thus $p \in K_0$ and $R \in K_2$ such that the region R is incident to the crossing point p . By definition of the potential, we have

$$\partial_{(p,R)} W = w_p - w_R,$$

where $w_p = (p, R)w'_p$ is the 4-cycle in Q given by the four arrows around the crossing point p and $w_R = (p, R)w'_R$ is the cycle in Q given by the arrows around the region




FIGURE 12. A crossing point p with an adjacent region R in the link diagram on the left and the corresponding paths in the quiver on the right. The cycle w_p is the 4-cycle

R , see Figure 12. We must show that $\partial_{(p,R)}W$ acts trivially on $M(\mathcal{S})$, and for that it suffices to show that the composition of the linear maps in the representation $M(\mathcal{S})$ along the paths w'_p and w'_R are equal.

We will write M_w for the composition of the arrows in $M(\mathcal{S})$ along a path w . According to Remark 6.2, we have $M_{w_p} = J_\ell$ for some $\ell \geq 0$, and thus it suffices to show $M_{w_R} = J_\ell$.

As before, we let \mathfrak{s} be the sequence of counterclockwise transpositions that transforms the minimal Kauffman state into the state \mathcal{S} . Let $\mathfrak{s}(R)$ be the subsequence of \mathfrak{s} consisting of all occurrences of the transpositions at the segments that bound the region R . Let a_0, a_1, \dots, a_t denote these segments in clockwise order around R such that a_0 is the first entry in the sequence $\mathfrak{s}(R)$. By Lemma 6.5, the subsequence $\mathfrak{s}(R)$ is of the form

$$\mathfrak{s}(R) = (a_0 a_1 \dots a_t)^\ell a_0 a_1 \dots a_u, \text{ with } u < t \text{ or}$$

$$\mathfrak{s}(R) = (a_0 a_1 \dots a_t)^\ell$$

for some $\ell \geq 0$. In the first case, the dimension of $M(\mathcal{S})$ is ℓ at vertices a_{u+1}, \dots, a_t and it is $\ell + 1$ at vertices a_0, a_1, \dots, a_u , while in the second case, the dimension is ℓ at all vertices a_0, a_1, \dots, a_t .

Denote the crossing point of the segments a_{j-1}, a_j by p_j . So the arrow (p_j, R) is $a_j \rightarrow a_{j-1}$. Then by definition of $M(\mathcal{S})$ at the crossing point p_j , we have

$$M_{(p_j, R)} = J_{\ell-1} \quad \text{if } j = 0 \text{ and } \mathfrak{s}(R) = (a_0, \dots, a_t)^\ell$$

and otherwise

$$M_{(p_j, R)} = \begin{cases} I_d & \text{if } \dim M_{a_j} = \dim M_{a_{j-1}} = d; \\ V_d & \text{if } \dim M_{a_j} - 1 = \dim M_{a_{j-1}} = d; \\ H_d & \text{if } \dim M_{a_j} = \dim M_{a_{j-1}} - 1 = d. \end{cases}$$

In particular $M_{w_R} = J_\ell$. This shows that $M(\mathcal{S})$ satisfies all relations of the form $\partial_\alpha W$, $\alpha \in Q_1$.

We also have to consider the closure I of the ideal generated by the relations $\partial_\alpha W$. For this we must show that arbitrary long paths act as zero; more precisely, if w is a path such that for all N there exists a path u_n of length $n > N$ such that $w = u_n$ then $M_w = 0$. Suppose w is such a path. Then, for all m there exists n such that there is an $x \in Q_0$ through which u_n passes at least m times. Thus u_n decomposes as

$$u_n = u w_1 w_2 \dots w_m v,$$

where each w_i is an oriented cycle that starts and ends at x and that does not pass through x another time. Take $m > \dim M_x = d$. We shall show below that, on

each cycle w_i , the matrix product M_{w_i} is some power of the matrix J_{d-1} . Therefore $M_{w_1 w_2 \dots w_n} = (J_{d-1})^{m+k}$, which is zero. Thus $M_{u_n} = 0$. Since $u_n = w$, we have $M_w = 0$, as desired.

It remains to show that, for every oriented cycle

$$w = a_0 \xrightarrow{\alpha_1} a_1 \xrightarrow{\alpha_2} \dots \xrightarrow{\alpha_{t-1}} a_t \xrightarrow{\alpha_t} a_0$$


in Q such that $a_i \neq a_j$ if $i \neq j$, the matrix M_w is a power of the matrix J_ℓ , for some ℓ . Let w be such a cycle. By definition of $M(\mathcal{S})$, for every arrow α_j , the matrix M_{α_j} is one of the four matrices I, V, H, J , which satisfy the relations $H_\ell V_\ell = J_\ell$ and $V_\ell H_\ell = J_{\ell-1}$. Hence if M_w is not of the claimed form then $M_w = I_\ell$ and all $M_{\alpha_j} = I_\ell$. Then $\dim M_{a_j} = \ell$ at each vertex a_j in w , which means that the transposition at a_j appears exactly ℓ times in the sequence \mathfrak{s} . Let a_k be the last transposition in the sequence \mathfrak{s} at a vertex in w . Consider the crossing point p where a_{k-1} and a_k meet in the link diagram. By Lemma 6.4, the last transposition at a_k does not increase the rank of the matrix M_{α_k} , since the arrow α_k ends in a_k . Thus $M_{\alpha_k} = J_{\ell-1}$ and hence $M_w \neq I_\ell$, and we are done. This completes the proof of part (a) of the proposition.

(b) Let $\mathcal{S}, \mathcal{S}'$ be two Kauffman states and suppose that $M(\mathcal{S}) = M(\mathcal{S}')$. Let \mathfrak{s} and \mathfrak{s}' be the sequences of transpositions that transform the minimal state into the the state \mathcal{S} and \mathcal{S}' , respectively. Let p be any crossing point and denote by $\mathfrak{s}(p)$ and $\mathfrak{s}'(p)$ the subsequences of \mathfrak{s} and \mathfrak{s}' consisting of all occurrences of transpositions at p . Since $M(\mathcal{S}) \cong M(\mathcal{S}')$, both representations have the same dimension vector and thus $\mathfrak{s}(p)$ and $\mathfrak{s}'(p)$ are equal up to a permutation. In fact, since the minimal state has exactly one marker at the point p , it follows that $\mathfrak{s}(p) = \mathfrak{s}'(p)$. At every crossing point p , the states \mathcal{S} and \mathcal{S}' are determined by the last entry in $\mathfrak{s}(p) = \mathfrak{s}'(p)$, and thus $\mathcal{S} = \mathcal{S}'$.

(c) If \mathcal{S} is the minimal state then its sequence of transpositions \mathfrak{s} is empty. Thus $M(\mathcal{S})$ is the zero module.

(d) Let \mathcal{S} be the maximal state. We have described Kauffman's construction of \mathcal{S} in [14] as a partition of K_1 in section 5.1. The fact that $M(\mathcal{S})$ and $T(i)$ have the same dimension vector follows directly from that. We now show that $M(\mathcal{S})$ and $T(i)$ also have the same linear maps.

Let $\alpha: a \rightarrow d$ be an arrow in Q . It is clear from the definition of $M(\mathcal{S})$ and $T(i)$ and by Lemma 6.3 that the linear maps on α are the same if the dimension at vertex a is different from the dimension at vertex d . Suppose therefore that $\dim M_a = \dim M_d = \ell$. Recall that the arrow α corresponds to a pair (p, R) , where p is a crossing point and R is an adjacent region in K . In the quiver Q , we have two corresponding chordless cycles w_p and w_R that share the arrow α as follows.



The crossing point cycle w_p is the cycle of length 4 on the left. The length of the region cycle w_R is the number of segments that bound R in the link diagram. We denote this length by $t + 1$. Note that in these two cycles the arrow α is the only arrow that ends at d . Consider the sequence of transpositions \mathfrak{s} that transforms the minimal state into the maximal state \mathcal{S} and let $\mathfrak{s}(p)$ and $\mathfrak{s}(R)$ be

the subsequences of all occurrences of transpositions at segments incident to p , respectively at segments bounding R . Suppose first that the first occurrence of a is before the first occurrence of d in $\mathfrak{s}(p)$. Then a must be the first entry in $\mathfrak{s}(p)$, because the direct predecessor of a would have to be d , by Corollary 6.6. Similarly, d must be the last entry in $\mathfrak{s}(p)$, because M_a and M_d have the same dimension ℓ and the direct successor of d in $\mathfrak{s}(p)$ would have to be a . Thus $\mathfrak{s}(p) = (abcd)^\ell$ and this shows that

- (i) $M_\alpha = J_{\ell-1}$, by definition of $M(\mathcal{S})$;
- (ii) The dimension of $M(\mathcal{S})$ is ℓ at each vertex of w_p and thus p is an internal point of $K'(\ell)$ as in Definition 5.3.

Now consider the sequence $\mathfrak{s}(R)$. Since a occurs before d in $\mathfrak{s}(p)$, it also does so in $\mathfrak{s}(R)$, and by the same argument as above, we see that a must be the first entry of $\mathfrak{s}(R)$ and d must be the last. Thus $\mathfrak{s}(R) = (aa_1a_2 \dots a_{t-1}d)^\ell$ and therefore the dimension of $M(\mathcal{S})$ is equal to ℓ at every vertex of w_R . The transposition at a moves two state markers at the endpoints of a counterclockwise. By our convention on the orientation of the quiver, the marker at the endpoint p must lie in the region R . Moreover, the fact that a is the first entry in both sequences $\mathfrak{s}(p)$ and $\mathfrak{s}(R)$ implies that the position (p, R) carries a state marker already in the minimal state. Similarly, since d is the last entry in $\mathfrak{s}(p)$ and $\mathfrak{s}(R)$, the position (p, R) also has a state marker in the maximal state. It follows from the construction of the minimal and maximal states in [14] that the region R is the region $R(p)$ of the internal point p as in Definition 5.3. Now the definition of the maps in the diagram module $T(i)$ implies that $T(i)_\alpha = J_{\ell-1}$. Hence $M(\mathcal{S})_\alpha = T(i)_\alpha$.

It remains the case where the first occurrence of d is before the first occurrence of a in $\mathfrak{s}(p)$. Then Lemma 6.4 implies that each occurrence of a in $\mathfrak{s}(p)$ augments the rank of $M(\mathcal{S})_\alpha$ by one. Thus our assumption $\dim M_a = \dim M_d = \ell$ implies that $M_\alpha = I_\ell$. On the other hand, we also have $T(i)_\alpha = I_\ell$, because the position (p, R) does not carry the state marker of maximal state, and thus R is not the region $R(p)$ of the internal point p . This completes the proof of part (d).

(e) It suffices to show that if the state \mathcal{S}' is obtained from the state \mathcal{S} by a single transposition at some segment a then $M(\mathcal{S})$ is a submodule of $M(\mathcal{S}')$. We use the notation $M(\mathcal{S}) = (M_x, M_\gamma)$, $M(\mathcal{S}') = (M'_x, M'_\gamma)$ and $d_x = \dim M_x$, $d'_x = \dim M'_x$. Define a morphism $f: M(\mathcal{S}) \rightarrow M(\mathcal{S}')$ by

$$f_j = \begin{cases} I_{d_j} & \text{if } j \neq a; \\ H_{d_j} & \text{if } j = a. \end{cases}$$

Clearly f is injective. To show that f is a morphism of B -modules, we need to consider arrows $b \xrightarrow{\alpha} a \xrightarrow{\beta} c$ in Q and show that the following diagram commutes.

$$\begin{array}{ccccc} M_b & \xrightarrow{M_\alpha} & M_a & \xrightarrow{M_\beta} & M_c \\ \downarrow f_b & & \downarrow f_a & & \downarrow f_c \\ M'_b & \xrightarrow{M'_\alpha} & M'_a & \xrightarrow{M'_\beta} & M'_c \end{array}$$

Since \mathcal{S}' is obtained from \mathcal{S} by the transposition at a , Lemmata 6.3 and 6.4 imply that $d_a \in \{d_b, d_b - 1\} = \{d_c, d_c - 1\}$. Moreover $d'_a = d_a + 1$, $d'_b = d_b$ and $d'_c = d_c$, and the maps $M_\alpha, M_\beta, M'_\alpha, M'_\beta$ are uniquely determined by the fact given in Lemma 6.4




FIGURE 13. A local configuration in the link diagram on the left and the corresponding configuration of the quiver on the right.

that $\text{rank}(M'_\alpha) = \text{rank}(M_\alpha)$ and $\text{rank}(M'_\beta) = \text{rank}(M_\beta) + 1$. Thus

$$\begin{aligned} M_\alpha &= I_\ell & \text{and} & & M'_\alpha &= H_\ell & \text{if } d_a = d_b = \ell; \\ M_\alpha &= V_\ell & \text{and} & & M'_\alpha &= J_\ell & \text{if } d_a = d_b - 1 = \ell; \\ M_\beta &= J_{\ell-1} & \text{and} & & M'_\beta &= V_\ell & \text{if } d_a = d_c = \ell; \\ M_\beta &= H_\ell & \text{and} & & M'_\beta &= I_{\ell+1} & \text{if } d_a = d_c - 1 = \ell. \end{aligned}$$

This shows that the diagram commutes and the proof of (e) is complete.

(f) Let \mathcal{S}_{max} denote the maximal Kauffman state. Thus $T(i) = M(\mathcal{S}_{max})$ by part (d). Let $d_a = \dim T(i)_a$. We fix a sequence of transpositions \mathfrak{s} that transforms the minimal state into the maximal state, and we use the basis of $T(i)$ induced by \mathfrak{s} . In particular, we have a basis $\{e_1, e_2, \dots, e_{d_a}\}$ for every vector space $T(i)_a$ with $a \in Q_0$.

Let $M = (M_a, M_\alpha)$ be a submodule of $T(i)$. Each vector space M_a is a subspace of $T(i)_a$, and thus the points in M_a can be expressed as coordinate vectors (x_1, \dots, x_{d_a}) with respect to our basis of $T(i)_a$. For every vertex $a \in Q_0$, let π_a denote the canonical projection from the vector space $M = \bigoplus_{j \in Q_0} M_j$ to the vector space M_a . For any point $x \in M$, we define an integer $m(a, x)$ as follows. If $\pi_a(x) = (x_1, \dots, x_{d_a}) \neq 0$, we let $m(a, x)$ be the unique integer such that $x_{m(a, x)} \neq 0$ and $x_k = 0$, for all $k = m(a, x) + 1, \dots, d_a$. If $\pi_a(x) = 0$, we let $m(a, x) = 0$. We then define a function $m: Q_0 \rightarrow \mathbb{Z}$ by $m(a) = \max_{x \in M} m(a, x)$. We will show that $m(a) = \dim M_a$.

If $m(a) = 0$ then $\pi_a(x) = 0$, for all $x \in M$, and thus M is not supported at a , whence $\dim M_a = 0$. If $m(a) = 1$ then $\pi_a(x) \in \text{span}\{e_1\}$, for all $x \in M$, and hence $\dim M_a = 1$.

Suppose now that $m(a) \geq 2$. Then $\dim T(i)_a \geq 2$ and thus the sequence \mathfrak{s} contains the transposition at a at least twice. Denote the crossing points at the ends of the segment a in K by p and p' , and denote the adjacent segments by b, c, d and b', c', d' in counterclockwise order as shown in the left picture of Figure 13. The sequence \mathfrak{s} must also contain the transpositions at b, c, d, b', c' and d' . The corresponding subquiver of Q is shown on the right in Figure 13. We have chordless cycles $w_p = \delta\gamma\beta\alpha$ and $w_{p'} = \delta'\gamma'\beta'\alpha'$. Because of Remark 6.2, the action of w_p on M_α is as follows. If $\pi_a(x) = (x_1, x_2, \dots, x_{m(a)}, 0, \dots, 0)$ then

$$\begin{aligned} \pi_a(x \cdot w_p) &= (x_2, x_3, \dots, x_{m(a)}, 0, \dots, 0) \\ \pi_a(x \cdot w_p^k) &= (x_{k+1}, x_{k+2}, \dots, x_{m(a)}, 0, \dots, 0) \\ \pi_a(x \cdot w_p^{m(a)-1}) &= (x_{m(a)}, 0, \dots, 0) \end{aligned} \tag{8}$$

Hence, since $x_{m(a)} \neq 0$, the vectors $\pi_a(x), \pi_a(x \cdot w_p), \dots, \pi_a(x \cdot w_p^{m(a)-1})$ are linearly independent vectors in M_a . Thus $M_a = \text{span}\{e_1, e_2, \dots, e_{m(a)}\}$ and hence $\dim M_a = m(a)$ as claimed.

In fact, each subspace $M_a \subset T(i)_a$, and hence the submodule $M \subset T(i)$, is completely determined by $m(a) = \dim M_a$. In particular, the submodules of $T(i)$ are determined by their dimension vector.

We will now show that M corresponds to a Kauffman state using induction on the total dimension $\ell = \sum_{a \in Q_0} \dim M_a$ of M . If $\ell = 0$ then $M = M(\mathcal{S}_{min})$ is the zero module. Suppose now that $\ell \geq 1$. Let $a \in Q_0$ be such that $S(a)$ is a direct summand of $\text{top } M$. Recall that $\text{top } M = M/\text{rad } M = M/(M \cdot \text{rad } B)$ is the largest semisimple quotient of M . In particular, we have a short exact sequence

$$0 \longrightarrow L \xrightarrow{f} M \xrightarrow{g} S(a) \longrightarrow 0, \quad (9)$$

where L is the kernel of g . By induction, we can assume that there exists a state \mathcal{S} such that $L = M(\mathcal{S})$. We will show that $M = M(\mathcal{S}')$, where \mathcal{S}' is the state obtained from \mathcal{S} by the transposition at the segment a .

Since M is determined by its dimension vector, we only need to show that the state \mathcal{S} admits the transposition at a . In other words, the markers at segment a must be in the positions indicated on the left of Figure 13 and corresponding to the arrows $\delta, \delta' \in Q_1$ on the right of the same figure. It suffices to show that the subsequence $\mathfrak{s}(p)$ of \mathfrak{s} ends in the transposition at d and the subsequence $\mathfrak{s}(p')$ ends in the transposition at d' . We will show this for $s(p)$ only, since the other case is symmetric.

By Lemma 6.3, we know that $|\dim L_a - \dim L_d| \leq 1$ and $|\dim M_a - \dim M_d| \leq 1$. Moreover, the short exact sequence (9) implies $\dim M_a = \dim L_a + 1$, and thus $\dim L_a = \dim L_d - 1$ or $\dim L_a = \dim L_d$. In the former case, the sequence $\mathfrak{s}(p)$ must end in d , and we are done.

Suppose now $\dim L_a = \dim L_d$. The morphism g of (9) gives rise to the commutative diagram

$$\begin{array}{ccc} M_b & \xrightarrow{M_\alpha} & M_a \\ \downarrow & & \downarrow g_a \\ 0 & \longrightarrow & S(a)_a \end{array}$$

with $g_a \neq 0$. Thus the commutativity implies that $g_a M_\alpha = 0$, and hence M_α is not surjective. Because of the description of the action of w_p in equation (8), the cokernel of $M_\alpha \circ M_\beta \circ M_\gamma \circ M_\delta$ is of dimension one, and thus the cokernel of M_α is of dimension one. Consequently the exactness in (9) implies that the map L_α in the representation L is surjective, and therefore the sequence $\mathfrak{s}(p)$ does not end in a , by Lemma 6.4(ii).

Next we show that $\mathfrak{s}(p)$ does not end in b or c . Suppose first that $\dim M_b = \dim M_a = \ell + 1$ then $\dim L_b = \dim L_a + 1$ and the sequence $\mathfrak{s}(p)$ starts with b and $L_\alpha = V_\ell$. The morphism f of (9) induces a commutative diagram

$$\begin{array}{ccc}
 L_b & \xrightarrow{L_\alpha=V_\ell} & L_a \\
 \downarrow f_b=I_{\ell+1} & & \downarrow f_a=H_\ell \\
 M_b & \xrightarrow{M_\alpha} & M_a
 \end{array}$$

whence $M_\alpha = J_\ell$.

Then $w_p = \delta\gamma\beta\alpha$ acts like M_α , and we obtain $M_\beta = M_\gamma = M_\delta = I$. Thus $L_\beta = L_\gamma = I$. Again Lemma 6.4(ii) implies that the sequence $\mathfrak{s}(p)$ does not end in b or c . Therefore $\mathfrak{s}(p)$ must end in d , and we are done.

On the other hand, if $\dim M_b \neq \dim M_a$ then $\dim M_b = \dim M_a - 1 = \ell$ and $\dim L_b = \dim L_a = \ell$. Then the definition of $L = M(\mathcal{S})$ implies that the map $L_\alpha: L_b \rightarrow L_a$ is either I_ℓ or $J_{\ell-1}$. In the latter case, the sequence $\mathfrak{s}(p)$ would end in a , a contradiction. Thus $L_\alpha = I_\ell$. In view of the sequence (9), we obtain $M_\alpha = H_\ell$, and from equation (8), we have $M_\alpha \circ M_\beta \circ M_\gamma \circ M_\delta = J_\ell$. In particular, M_β, M_γ and M_δ are surjective, and using (9) again, we see that L_β and L_γ are surjective as well, since β and γ are not incident to the vertex a . Now Lemma 6.4(ii) implies that $\mathfrak{s}(p)$ does not end in b or c . This completes the proof of part (f). \square

Corollary 6.8. (a) For every dimension vector \mathbf{e} the quiver Grassmannian $\text{Gr}_\mathbf{e}(T(i))$ is either empty or a point. In particular, the Euler characteristic

$$\chi(\text{Gr}_\mathbf{e}(T(i))) = 0 \text{ or } 1.$$

(b) The F -polynomial of $T(i)$ is

$$F_{T(i)} = \sum_{L \subset T(i)} \mathbf{y}^{\dim L},$$

where the sum is over all submodules of $T(i)$ and $\mathbf{y}^{\dim L} = \prod_{i=1}^{2n} y_i^{\dim L_i}$.

Proof. In the proof of part (f) of the proposition, we have seen that every submodule of $T(i)$ is determined by its dimension vector. Thus if there is a submodule of dimension vector \mathbf{e} then it is unique, and $\text{Gr}_\mathbf{e}(T(i))$ is a point. Otherwise it is empty. This shows (a), and (b) follows directly. \square

We are ready for the main result of this section.

Theorem 6.9. The map $\mathcal{S} \mapsto M(\mathcal{S})$ is a lattice isomorphism from the lattice of Kauffman states of K relative to the segment i to the lattice of submodules of the module $T(i)$.

Proof. The map is well-defined by parts (a) and (d) of Proposition 6.7, injective by part (b) and surjective by part (f). Part (e) implies that it is order preserving. Moreover the maximum and minimum elements correspond by parts (c) and (d). \square

6.3. Indecomposability.

Proposition 6.10. The B -module $T(i)$ is indecomposable.

Proof. First we show that the support of $T(i)$ induces a connected subquiver of Q . By definition of $T(i)$, the support consists of all segments in $K_1 \setminus K(0)$, which is obtained from K by removing the two regions R_1, R_2 that are incident to the

segment i . If $K \setminus \{R_1, R_2\}$ is disconnected then we can draw a closed curve γ in the plane that separates one connected component from the rest. Putting back R_1 and R_2 , we see that γ crosses exactly two regions in K . This means that K is a connected sum which contradicts our assumption that K is prime. Thus the support of $T(i)$ is connected.

Suppose now $T(i) = M \oplus N$ is the direct sum of two nonzero B -modules. Let $\alpha: j \rightarrow k$ be an arrow in Q such that $T(i)_j$ and $T(i)_k$ are nonzero. By definition, the linear map $T(i)_\alpha$ are nonzero, except possibly if there is a crossing point cycle $j \xrightarrow{\alpha} k \rightarrow l \rightarrow m \rightarrow j$ such that $T(i)$ has dimension one at vertices j, k, l and m , in which case one of the four maps is zero and the others are the identity. Therefore, since the support of $T(i)$ induces a connected subquiver of Q , the supports of the two submodules M and N cannot be disjoint.

Then there exists $j \in Q_0$ such that M_j and N_j are both nonzero. Since j is a segment in K , there exists a crossing point $p \in K_0$ that is incident to j . Let w_p denote the corresponding 4-cycle in Q . Because of Remark ??, the vector space $T(i)_j = M_j \oplus N_j$ has a basis $\{e_1, e_2, \dots, e_d\}$ with respect to which the action of w_p is given by the matrix J_{d-1} .

Now let $m \in M_j, n \in N_j$ be nonzero elements and denote their expansions in the basis as

$$m = \sum_{k=1}^n \mu_k e_k \quad n = \sum_{k=1}^n \nu_k e_k,$$

with $\mu_k, \nu_k \in \mathbb{k}$. Let k_m and k_n be the largest indices such that $\mu_{k_m} \neq 0$ and $\nu_{k_n} \neq 0$. Then $m \cdot w_p^{k_m-1} = \mu_{k_m} e_1 \in M_j$ and $n \cdot w_p^{k_n-1} = \nu_{k_n} e_1 \in N_j$, because M and N are right B -modules. Dividing by the scalars shows that $e_1 \in M_j \cap N_j$, a contradiction to the assumption that the sum $T(i) = M \oplus N$ is direct. \square

7. THE MAIN RESULT

In this section, we prove that the Alexander polynomial of the link is a specialization of the F -polynomial of any indecomposable summand of the link diagram module.

Let K be an oriented link diagram of a prime link and assume that K contains no curls. Let n be the number of crossing points in K , and let i be a segment in K . Let $T(i)$ be the corresponding indecomposable summand of the link diagram module and $F_{T(i)} = \sum_{L \subset T(i)} \mathbf{y}^{\dim L}$ its F -polynomial as in Corollary 6.8(b).

For $f \in \mathbb{Z}[y_1, \dots, y_{2n}]$ we write $f|_t$ for the specialization of f at

$$y_j = \begin{cases} -t & \text{if segment } j \text{ runs from an undercrossing to an overcrossing;} \\ -t^{-1} & \text{if segment } j \text{ runs from an overcrossing to an undercrossing;} \\ -1 & \text{if segment } j \text{ connects two overcrossings or two undercrossings.} \end{cases} \quad (10)$$

Theorem 7.1. *The Alexander polynomial of K is equal to the specialization (10) of the F -polynomial of every indecomposable summand $T(i)$ of the link diagram module T . That is*

$$\Delta = F_{T(i)}|_t.$$

Proof. Let Δ denote the Alexander polynomial of K . Kauffman's theorem says that

$$\Delta \doteq \sum_{\mathcal{S}} \sigma(\mathcal{S}) w(\mathcal{S}), \quad (11)$$

where the sum is over all states, $\sigma(\mathcal{S}) = \pm 1$, and $w(\mathcal{S})$ is a power of t . The symbol \doteq means that the expressions on either side are equal up to sign and up to a power of t . We denote by \mathcal{S}_{min} the minimal state. Normalizing the above identity, we find

$$\Delta \doteq \sum_{\mathcal{S}} \frac{\sigma(\mathcal{S})}{\sigma(\mathcal{S}_{min})} \frac{w(\mathcal{S})}{w(\mathcal{S}_{min})}. \quad (12)$$

Let $\mathfrak{s} = j_1, j_2, \dots, j_t$ be a sequence of transpositions that transforms \mathcal{S}_{min} into \mathcal{S} , and let $M(\mathcal{S})$ denote the state module introduced in Definition 6.1. Then

$$\underline{\dim} M(\mathcal{S}) = \sum_{k=1}^t \mathbf{e}_{j_k}, \quad (13)$$

where $\mathbf{e}_{j_k} \in \mathbb{Z}^{2n}$ is the vector that is 1 at position j_k and 0 elsewhere.

Recall that, if a state \mathcal{S}' is obtained from a state \mathcal{S} by a single transposition at a segment $j \in K_1$ then $w(j) = w(\mathcal{S}')/w(\mathcal{S})$ is independent of the particular states $\mathcal{S}, \mathcal{S}'$ and only depends on the segment j . Moreover, in this situation, Lemma 2.7 of [14] implies that $\sigma(\mathcal{S}') = -\sigma(\mathcal{S})$. Thus

$$\frac{\sigma(\mathcal{S})}{\sigma(\mathcal{S}_{min})} \frac{w(\mathcal{S})}{w(\mathcal{S}_{min})} = (-w(j_1))(-w(j_2)) \dots (-w(j_t)). \quad (14)$$

From our table in Figure 4 we know that $-w(j)$ is equal to the specialization of y_j at (11). Thus

$$-w(j) = y_j|_t = \mathbf{y}^{\mathbf{e}_j}|_t.$$

Therefore the right hand side of equation (14) is equal to $\mathbf{y}^{\underline{\dim} M(\mathcal{S})}|_t$. Applying this result to the formula in (12), we obtain

$$\Delta \doteq \sum_{\mathcal{S}} \mathbf{y}^{\underline{\dim} M(\mathcal{S})}|_t.$$

Now Theorem 6.9 implies

$$\Delta \doteq \sum_{L \subset T(i)} \mathbf{y}^{\underline{\dim} L}|_t = F_{T(i)}|_t,$$


where the sum is over all submodules of $T(i)$, and thus it is the specialized F -polynomial. \square

8. A SPECIAL CASE: 2-BRIDGE LINKS

A special class of links is the family of 2-bridge links $K_{[a_1, a_2, \dots, a_n]}$ which were first studied by Schubert in [28]. These links are parametrized by continued fractions

$$[a_1, a_2, \dots, a_n] = a_1 + \frac{1}{a_2 + \frac{1}{\ddots + \frac{1}{a_n}}} \quad (15)$$

with $a_i \in \mathbb{Z}_{\geq 1}$. The link $K_{[a_1, \dots, a_n]}$ consists of n braids on two strands whose number of crossings is given by the a_i and that are joined together in a linear fashion as shown in Figure 14 and such that the resulting link is alternating. $K_{[a_1, \dots, a_n]}$ is a knot if the numerator of the continued fraction (15) is odd, and it is a link with exactly two components otherwise. For example, $K[2, 1, 2, 3]$ in Figure 14 is a knot, because $[2, 1, 2, 3] = 27/10$.

FIGURE 14. The 2-bridge knot $K[2, 1, 2, 3]$

Let i be the long segment at the bottom of the link diagram that connects the a_1 -braid to the a_n -braid. Then our construction of the link module $T(i)$ produces a Dynkin type \mathbb{A} module whose support is given by one of the quivers

$$1 \leftarrow 2 \leftarrow \dots \leftarrow \ell_1 \rightarrow \dots \rightarrow \ell_2 \leftarrow \dots \leftarrow \ell_{n-1} \rightarrow \dots \rightarrow (\ell_n - 1),$$

$$1 \leftarrow 2 \leftarrow \dots \leftarrow \ell_1 \rightarrow \dots \rightarrow \ell_2 \rightarrow \dots \rightarrow \ell_{n-1} \leftarrow \dots \leftarrow (\ell_n - 1),$$

where $\ell_j = a_1 + a_2 + \dots + a_j$ and direction of the arrows at the right end depend on the parity of n . The module $T(i)$ is of dimension one at each vertex $1, 2, \dots, (\ell_n - 1)$ and its linear maps on the arrows shown above are the identity maps.

In this situation, the F -polynomial of $T(i)$ can be computed as a sum over all perfect matchings of the snake graph $\mathcal{G}_{[a_1, \dots, a_n]}$ associated to the continued fraction in [5]. We have

$$F_{T(i)} = \sum_{P \in \text{Match } \mathcal{G}_{[a_1, \dots, a_n]}} y(P)$$

where $y(P)$ is the height function of the poset of perfect matchings [21].

Remark 8.1. This module $T(i)$ was implicitly used in [17], where the Jones polynomial was realized as the specialization of the F -polynomial of $T(i)$ at $y_1 = t^{-2}$ and $y_j = -t^{-1}$ for all $j = 2, 3, \dots, n$. We do not know how to generalize this specialization to other segments i of this link, or to other type of links.

We obtain two consequences from the above discussion.

8.1. Type \mathbb{A} cluster variables correspond to links. We have the following result.

Theorem 8.2. *Let Q be a quiver of Dynkin type \mathbb{A} and let $\mathcal{A}(Q)$ be its cluster algebra. For every non-initial cluster variable $x \in \mathcal{A}(Q)$ there exists a link diagram K and a segment $i \in K_1$ such that the indecomposable summand $T(i)$ of the link module is mapped to x under the Caldero-Chapoton map.*

Proof. Let CC denote the Caldero-Chapoton map. Since x is non-initial, there exists an indecomposable $\mathbb{k}Q$ -module M such that $\text{CC}(M) = x$, [4, 3]. The support of M defines a connected subquiver of Q which in turn determines a continued fraction [5] and hence a 2-bridge link K . From the discussion above and our main theorem, we have a segment $i \in K_1$ such that $T(i) = M$. \square

8.2. An application to q -deformed rationals. In [20], Morier-Genoud and Ovsienko introduced q -deformed rationals and q -deformed continued fractions. They propose a unimodality conjecture that can be rephrased in terms of the specialized height function as follows.

Let M be an indecomposable type \mathbb{A} module and let h be the linearization of the lattice of submodules of M that maps submodules L of M to their total

dimension. Thus $h(L) = \dim L = \sum_{j \in Q_0} \dim L_j$. Equivalently, we can think of h as a linearization of the lattice of perfect matchings of the associated snake graph \mathcal{G} that maps a perfect matching P to the specialization of the height function setting all y -variables equal to t . Thus $h(P) = y(P)|_{y_j=t}$. In other words, h associates to each lattice element the length of the shortest chain from the element to the minimal element in the lattice.

Conjecture 8.3 (Morier-Genoud–Ovsienko). The function h is unimodal.

Progress towards this conjecture has been made in [19].

Using our main theorem and properties of the Alexander polynomial, we have the following result, which says that the alternating sum of the number of objects on each level of the poset is $-1, 0$, or 1 .

Theorem 8.4. *Let M be a module of Dynkin type \mathbb{A}_n and \mathcal{L} the submodule lattice of M . Then*

$$\sum_{L \in \mathcal{L}} (-1)^{h(L)} = \begin{cases} \pm 1 & \text{if } |\mathcal{L}| \text{ is odd;} \\ 0 & \text{if } |\mathcal{L}| \text{ is even.} \end{cases}$$

Proof. Combining Theorems 8.2 and 7.1, we see that there exists a 2-bridge link K whose Alexander polynomial is the specialized F -polynomial of M . More precisely,

$$\Delta_K \doteq F_M|_t = \sum_{L \in \mathcal{L}} \mathbf{y}^{\dim L}|_t.$$

From the definition of the specialization (10), we see that evaluating the above equation at $t = 1$ gives

$$\Delta_K(1) = \sum_{L \in \mathcal{L}} \mathbf{y}^{\dim L}|_{y_i=-1}. \tag{16}$$

Furthermore

$$\mathbf{y}^{\dim L}|_{y_i=-1} = \prod_i y_i^{\dim L_i}|_{y_i=-1} = (-1)^{\sum_i \dim L_i} = (-1)^{h(L)}.$$

Thus equation (16) becomes

$$\Delta_K(1) = \sum_{L \in \mathcal{L}} (-1)^{h(L)}.$$

Now the result follows from property (ii) of subsection 2.1.1 and the fact that K is a knot if and only if the number of submodules of M is odd. \square

9. EXAMPLES

Example 9.1. Consider the figure-eight knot. We use the same labelling of segments as in Example 3.1. The lattice of the submodules of the module $T(1)$ is shown at Figure 15a.


The lattice isomorphism with the lattice of Kaufman states with regards to the segment 1 is obvious, see Figure 3. The F -polynomial of $T(1)$ is

$$F_{T(1)} = 1 + y_2 + y_8 + y_2y_8 + y_2y_5y_8$$

and its specialization at $y_2 = -t$, $y_5 = -t^{-1}$ and $y_8 = -t$, as given by Equation 10, is

$$F_{T(1)}|_t = 1 - 3 + t^2.$$

Example 9.3. For the Conway knot, the quiver is



The F -polynomial of $T(18)$ has 131 terms. The highest degree term is

$$y_2 y_3 y_5 y_6 y_8 y_9 y_{10} y_{13} y_{14} y_{15} y_{16} y_{17} y_{19} y_{21}$$

The specialization gives $F_{T(18)}|_t = t$, confirming that the Alexander polynomial is trivial.

REFERENCES

- [1] J. W. Alexander, Topological invariants of knots and links, *Trans. Amer. Math. Soc.* **30** (2), (1928) 275–306.
- [2] I. Assem, R. Schiffler, V. Shramchenko, Cluster automorphisms, *Proc. London Math. Soc.* **3** no. 104, 1271–1302 (2012).
- [3] P. Caldero and F. Chapoton, Cluster algebras as Hall algebras of quiver representations. *Comment. Math. Helv.* **81** (2006), no. 3, 595–616.
- [4] P. Caldero, F. Chapoton and R. Schiffler, Quivers with relations arising from clusters (\mathbb{A}_n case). *Trans. Amer. Math. Soc.* **358** (2006), no. 3, 1347–1364 (electronic).
- [5] İ. Çanakçı and R. Schiffler, Cluster algebras and continued fractions, *Compos. Math.*, **54** (3) (2018) 565–593.
- [6] M. Cohen, O. Dasbach and H. Russell, A twisted dimer model for knots. *Fund. Math.* **225** (2014), no. 1, 57–74.

- [7] J. H. Conway, An enumeration of knots and links, and some of their algebraic properties. 1970 Computational Problems in Abstract Algebra (Proc. Conf., Oxford, 1967) pp. 329–358 Pergamon, Oxford.
- [8] H. Derksen, J. Weyman and A. Zelevinsky, Quivers with potentials and their representations. I. Mutations. *Selecta Math.* (N.S.) **14** (2008), no. 1, 59–119.
- [9] H. Derksen, J. Weyman and A. Zelevinsky, Quivers with potentials and their representations II: Applications to cluster algebras. *J. Amer. Math. Soc.* **23** (2010), no. 3, 749–790.
- [10] S. Fomin, L. Williams and A. Zelevinsky, Introduction to Cluster Algebras, Chapter 7, arXiv:2106.02160.
- [11] S. Fomin and A. Zelevinsky, Cluster algebras I: Foundations, *J. Amer. Math. Soc.* **15** (2002), 497–529.
- [12] S. Fomin and A. Zelevinsky, Cluster algebras IV: Coefficients, *Compos. Math.* **143** (2007), 112–164.
- [13] M. H. Freedman, A surgery sequence in dimension four; the relations with knot concordance. *Invent. Math.* **68** (1982), no. 2, 195–226.
- [14] L. Kauffman, Formal Knot Theory, Mathematical Notes, 30. Princeton University Press, Princeton, NJ, 1983.
- [15] B. Keller, Mutation applet.
- [16] K. Lee and R. Schiffler, Positivity for cluster algebras, *Annals of Math.* **182** (1), (2015) 73–125.
- [17] K. Lee and R. Schiffler, Cluster algebras and Jones polynomials, *Sel. Math. New Ser.* (2019) 25: 58.
- [18] W. B. R. Lickorish, An introduction to knot theory. Graduate Texts in Mathematics, 175. Springer-Verlag, New York, 1997.
- [19] T. McConville, B. Sagan and C. Smyth, On a rank-unimodality conjecture of Morier-Genoud and Ovsienko. *Discrete Math.* **344** (2021), no. 8, Paper No. 112483, 13 pp.
- [20] S. Morier-Genoud and V. Ovsienko, q -deformed rationals and q -continued fractions, preprint, arXiv:1812.00170. *Forum Math. Sigma* **8** (2020), Paper No. e13, 55 pp.
- [21] G. Musiker and R. Schiffler, Cluster expansion formulas and perfect matchings, *J. Algebraic Combin.* **32** (2010), no. 2, 187–209.
- [22] W. Nagai and Y. Terashima, Cluster variables, ancestral triangles and Alexander polynomials, *Adv. Math.* **363** (2020), 106965, 37 pp.
- [23] P. Ozsváth, and Z. Szabó, Holomorphic disks and knot invariants. *Adv. Math.* **186** (2004), no. 1, 58–116.
- [24] L. Piccirillo, The Conway knot is not slice, *Ann. of Math.* (2) **191** (2020), no. 2, 581–591.
- [25] J. A. Rasmussen, Floer homology and knot complements. Thesis (Ph.D.)—Harvard University. 2003. 126 pp.
- [26] R. Schiffler, Cluster algebras from surfaces: lecture notes for the CIMPA School Mar del Plata, March 2016. Homological methods, representation theory, and cluster algebras, 65–99, CRM Short Courses, Springer, Cham, 2018.
- [27] R. Schiffler and D. Whiting, Tilting modules arising from knot invariants, preprint, 18 pages, arXiv:2001.04004.
- [28] H. Schubert, Knoten mit zwei Brüchen. (German) *Math. Z.* **65** (1956), 133–170.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269-1009, USA
 Email address: `veronique.bazier-matte@uconn.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269-1009, USA
 Email address: `schiffler@math.uconn.edu`