

Elimination of turbulent modes using a conditional average with asymptotic freedom

W.D. McComb and C. Johnston

Department of Physics, University of Edinburgh, Mayfield Road, Edinburgh, Scotland, EH9 3JZ

(April 12, 1999)

Abstract

The part of the nonlinear term in the Navier-Stokes equation which represents coupling to the small scale modes may be averaged out by introducing a weak conditional average with asymptotic freedom in wavenumber. A residual deterministic part, while important for individual realizations, makes a negligible contribution to the renormalization of the dissipation rate. This is because the full ensemble average, needed to establish the energy balance, relaxes the constraint on the conditional average.

47.27.Ak, 47.27.Eq, 47.27.Gs, 05.20.-y

Typeset using REVTeX

The application of renormalization group to dynamical problems in microscopic physics requires an average over small scales in which large scales are held fixed [1]. Unfortunately, the corresponding procedure for classical nonlinear systems, such as Navier-Stokes turbulence, is impossible *in principle* because of the deterministic nature of such systems. Recently it has been proposed that the chaotic nature of turbulence may justify the use of an approximate conditional average [2]. In the present paper we argue that the conditional elimination of a band of high wavenumber modes may be accomplished in terms of a deterministic part, which has a coherent phase relation with the retained modes, and a random part, which is asymptotically free and may be averaged out with the introduction of an effective viscosity. The reduction of the number of modes takes place at a constant rate of energy dissipation, and it is further argued that the renormalization of this quantity can be adequately represented by the incoherent part only. This is because the full ensemble average, needed for the spectral energy balance, tends to ‘lift’ the constraint on the conditional average.

We consider incompressible fluid turbulence, as governed by the solenoidal Navier-Stokes equation (NSE)

$$\left(\partial_t + \nu_0 k^2\right) u_\alpha(\mathbf{k}, t) = M_{\alpha\beta\gamma}(\mathbf{k}) \int d^3j u_\beta(\mathbf{j}, t) u_\gamma(\mathbf{k} - \mathbf{j}, t), \quad (1)$$

where ν_0 is the kinematic viscosity of the fluid,

$$M_{\alpha\beta\gamma}(\mathbf{k}) = (2i)^{-1} [k_\beta D_{\alpha\gamma}(\mathbf{k}) + k_\gamma D_{\alpha\beta}(\mathbf{k})], \quad (2)$$

and the projector $D_{\alpha\beta}(\mathbf{k})$ is expressed in terms of the Kronecker delta $\delta_{\alpha\beta}$ as

$$D_{\alpha\beta}(\mathbf{k}) = \delta_{\alpha\beta} - k_\alpha k_\beta |\mathbf{k}|^{-2}. \quad (3)$$

In order to pose a specific problem, we restrict our attention to stationary, isotropic, homogeneous turbulence, with dissipation rate ε and zero mean velocity. We also introduce an upper cutoff wavenumber K_{\max} , which is defined through the dissipation integral

$$\varepsilon = \int_0^\infty 2\nu_0 k^2 E(k) dk \simeq \int_0^{K_{\max}} 2\nu_0 k^2 E(k) dk, \quad (4)$$

where $E(k)$ is the energy spectrum, so ensuring that K_{\max} is of the same order of magnitude as the Kolmogorov dissipation wavenumber.

We then filter the velocity field at $|\mathbf{k}| \equiv k = K_c$, where $0 < K_c < K_{\max}$, according to

$$u_\alpha(\mathbf{k}, t) = \begin{cases} u_\alpha^-(\mathbf{k}, t) & \text{for } 0 < k < K_c \\ u_\alpha^+(\mathbf{k}, t) & \text{for } K_c < k < K_{\max}. \end{cases} \quad (5)$$

The NSE may be decomposed using (5), to give

$$(\partial_t + \nu_0 k^2) u_k^- = M_k^- (u_j^- u_{k-j}^- + 2u_j^- u_{k-j}^+ + u_j^+ u_{k-j}^+), \quad (6)$$

$$(\partial_t + \nu_0 k^2) u_k^+ = M_k^+ (u_j^- u_{k-j}^- + 2u_j^- u_{k-j}^+ + u_j^+ u_{k-j}^+), \quad (7)$$

where, for simplicity, all vector indices and independent variables are contracted into a single subscript.

In order to obtain an expression for the average effect of the high wavenumber modes upon a particular low wavenumber mode, we need to average out the u^+ whilst holding the u^- constant. This requires a *conditional average* $\langle \cdot \rangle_c$ such that

$$\langle u_\alpha^-(\mathbf{k}, t) \rangle_c = u_\alpha^-(\mathbf{k}, t). \quad (8)$$

This is the *only* rigorous property we can attribute to the conditional average, and it should also be noted that it is vital to distinguish between this operation and that of a filtered ensemble average.

To establish the statistical properties of $u_\alpha(\mathbf{k}, t)$ we consider an ensemble \mathcal{W} consisting of the set of M time-independent realizations $\{W_\alpha^{(i)}(\mathbf{k})\}$, each realization [3] being labelled by an integer i . Subject to certain weak conditions, the ensemble average is

$$\langle u_\alpha(\mathbf{k}, t) \rangle = \lim_{M \rightarrow \infty} \frac{1}{M} \sum_{i=1}^M W_\alpha^{(i)}(\mathbf{k}) = \bar{U}_\alpha(\mathbf{k}), \quad (9)$$

where $\bar{U}_\alpha(\mathbf{k})$ is the time average of $u_\alpha(\mathbf{k}, t)$. This procedure can then be extended to any well behaved functional, $F[u_\alpha(\mathbf{k}, t)]$, thus:

$$\langle F[u_\alpha(\mathbf{k}, t)] \rangle = \lim_{M \rightarrow \infty} \frac{1}{M} \sum_{i=1}^M F[W_\alpha^{(i)}(\mathbf{k})]. \quad (10)$$

Now we consider how to perform a *conditional* average. To do this, we first select a subensemble, $\mathcal{Y} \equiv \{Y_\alpha^{(i)}(\mathbf{k})\} \subset \mathcal{W}$, and choose the members of this *biased* subensemble to be those N ($N \leq M$) members of \mathcal{W} satisfying the criterion

$$\lim_{\delta \rightarrow 0} \left(\max |\theta^-(k) W_\alpha^{(i)}(\mathbf{k}) - u_\alpha^-(\mathbf{k}, t_1)| \leq \delta \right), \quad (11)$$

where t_1 is some fixed time and $\theta^-(k) = 1$ for $0 < k < K_c$, and zero otherwise. The conditional average is then obtained by generalizing (9) and (10) to the biased subensemble, viz.,

$$\langle u_\alpha(\mathbf{k}, t) \rangle_c = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N Y_\alpha^{(i)}(\mathbf{k}), \quad (12)$$

and

$$\langle F[u_\alpha(\mathbf{k}, t)] \rangle_c = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N F[Y_\alpha^{(i)}(\mathbf{k})]. \quad (13)$$

It follows by construction that (8) holds, since from (11) and (12)

$$\langle u_\alpha^-(\mathbf{k}, t) \rangle_c = \lim_{N \rightarrow \infty} \frac{1}{N} [N u_\alpha^-(\mathbf{k}, t)] = u_\alpha^-(\mathbf{k}, t). \quad (14)$$

The difficulty now facing us lies in the nature of the subensemble, which is an example of deterministic chaos. This can be seen if we consider two extreme scenarios for the behavior of u^+ under the conditional average. Firstly, if we assume that the subensemble is *strictly* deterministic, then in this instance u_k^+ is fully determined by prescribing u_k^- . Accordingly, Eq. (8) implies that $\langle u_j^- u_{k-j}^- \rangle_c = u_j^- u_{k-j}^-$, $\langle u_j^- u_{k-j}^+ \rangle_c = u_j^- u_{k-j}^+$ and $\langle u_j^+ u_{k-j}^+ \rangle_c = u_j^+ u_{k-j}^+$. Thus, the low-pass filtered NSE, Eq. (6), reduces back to itself under the conditional average. Secondly, if we assume that the subensemble is purely random, it follows that in this case, u_k^+ is independent of u_k^- . Hence, applying the conditional average to the low-pass filtered NSE, we find

$$(\partial_t + \nu_0 k^2) u_k^- = M_k^- u_j^- u_{k-j}^-,$$

the $u_j^- u_{k-j}^+$ term being zero since the ensemble average of u^+ is zero, whilst the $u_j^+ u_{k-j}^+$ term is zero due to homogeneity. Thus in this scenario it appears that there is *no* effect of nonlinear coupling.

In reality we are faced with a situation somewhere between these two extremes, and so we replace our criterion for members of the biased subensemble, Eq. (11), which is equivalent to the first of these situations if $\delta = 0$, by the less precise criterion

$$\max |\theta^-(k)W_\alpha^{(i)}(\mathbf{k}) - u_\alpha^-(\mathbf{k}, t_1)| \leq \xi, \quad (15)$$

where, in general, ξ is of the order of the turbulent velocities involved.

To obtain a non-trivial conditional average we must now identify those circumstances in which ξ may be neglected as being, in some sense, small. A measure of the ‘smallness of ξ ’ can be identified by constructing the subensemble as

$$W_\alpha^{(i)}(\mathbf{k}) = u_\alpha^-(\mathbf{k}, t_1) + \phi_\alpha^{(i)}(\mathbf{k}, t_1), \quad (16)$$

where i is any label satisfying (15). If we then further restrict the subensemble to be such that the set $\{\phi_\alpha^{(i)}(\mathbf{k}, t_1)\}$ satisfies (8), we find that

$$\langle u_j^- u_{k-j}^- \rangle_c = u_j^- u_{k-j}^- + \langle \phi_j \phi_{k-j} \rangle_c. \quad (17)$$

Thus in order to maintain form invariance of the NSE under conditional averaging, we require

$$\langle \phi_j \phi_{k-j} \rangle_c \rightarrow 0 \quad (18)$$

in some limit. This is our criterion for the smallness of ξ .

If we further suppose that chaos and unpredictability are local characteristics of turbulence, and there is support for such a view [4,5], then if K_c and K_{\max} are sufficiently far apart we might expect, due to the development of unpredictability as k is increased above K_c , that the effect of the constraint given in Eq. (15) would die away, such that

$$\lim_{k \rightarrow K_{\max}} \langle u_\alpha^+(\mathbf{k}, t) \rangle_c \rightarrow \langle u_\alpha^+(K_{\max}, t) \rangle. \quad (19)$$

We refer to this property as *asymptotic freedom*. In order to extend this concept to higher-order moments, we introduce the following *Hypothesis of Local Chaos*:

“For sufficiently large Reynolds’ number and corresponding K_{\max} , there exists a cut-off wavenumber $K_c < K_{\max}$ such that a mixed conditional moment involving p low wavenumber and r high wavenumber modes takes the limiting form

$$\lim_{\xi \rightarrow 0} \langle u_{\alpha}^{-}(\mathbf{k}_1, t) u_{\beta}^{-}(\mathbf{k}_2, t) \dots u_{\gamma}^{-}(\mathbf{k}_p, t) u_{\delta}^{+}(\mathbf{k}_{p+1}, t) u_{\epsilon}^{+}(\mathbf{k}_{p+2}, t) \dots u_{\sigma}^{+}(\mathbf{k}_{p+r}, t) \rangle_c \rightarrow u_{\alpha}^{-}(\mathbf{k}_1, t) u_{\beta}^{-}(\mathbf{k}_2, t) \dots u_{\gamma}^{-}(\mathbf{k}_p, t) \lim_{\{\cdot\} \rightarrow K_{\max}} \langle u_{\delta}^{+}(\mathbf{k}_{p+1}, t) u_{\epsilon}^{+}(\mathbf{k}_{p+2}, t) \dots u_{\sigma}^{+}(\mathbf{k}_{p+r}, t) \rangle, \quad (20)$$

where $\lim_{\{\cdot\} \rightarrow K_{\max}}$ means take the limit for all wavevector arguments of the u^{+} modes, with the condition of Eq. (18) satisfied as a corollary.”

This provides our definition of an asymptotic conditional average and may be used to evaluate all terms involving mixed products of u^{-} with u^{+} . For example,

$$\lim_{\xi \rightarrow 0} \langle u_j^{-} u_{k-j}^{+} \rangle_c = u_j^{-} \lim_{\{\cdot\} \rightarrow K_{\max}} \langle u_{k-j}^{+} \rangle = 0, \quad (21)$$

since $\langle u_{\alpha}^{+}(\mathbf{k}, t) \rangle = 0$. Note also that the hypothesis as stated is more general than is necessary, since we shall only need to consider products containing at most two u^{-} modes.

If we then take the conditional average of the low-pass filtered NSE, Eq. (6), we obtain

$$(\partial_t + \nu_0 k^2) u_k^{-} = M_k^{-} \{ \langle u_j^{-} u_{k-j}^{-} \rangle_c + 2 \langle u_j^{-} u_{k-j}^{+} \rangle_c + \langle u_j^{+} u_{k-j}^{+} \rangle_c \}, \quad (22)$$

where the conditional average of u_k^{-} on the left hand side has been evaluated using (8). This equation may be further re-written as

$$(\partial_t + \nu_0 k^2) u_k^{-} = M_k^{-} u_j^{-} u_{k-j}^{-} + S^{-}(k|K_c) + M_k^{-} \lim_{\xi \rightarrow 0} \langle u_j^{+} u_{k-j}^{+} \rangle_c, \quad (23)$$

where

$$S^{-}(k|K_c) = M_k^{-} \{ \langle \phi_j^{-} \phi_{k-j}^{-} \rangle_c + 2 \langle u_j^{-} u_{k-j}^{+} \rangle_c + \langle u_j^{+} u_{k-j}^{+} \rangle_c - \lim_{\xi \rightarrow 0} \langle u_j^{+} u_{k-j}^{+} \rangle_c \}. \quad (24)$$

It should also be noted that the hypothesis *must* hold for $K_c \rightarrow 0$, as in this instance Eq. (22) reduces to the Reynolds equation, with $u_{\alpha}(\mathbf{k}, t) \rightarrow \bar{U}_{\alpha}(\mathbf{k})$ as given by (9).

Our hypothesis does not explicitly tell us how evaluate the conditional average in (23), which involves a non-trivial projection of a product of u^{+} modes in the Hilbert space of

the u^- modes, but we may use the high-pass filtered NSE, Eq. (7), to obtain a governing equation for this quantity. To do this, we use (7) to write equations for u_j^+ and u_{k-j}^+ , multiply these equations by u_{k-j}^+ and u_j^+ respectively, add the resulting equations together, and then take the conditional average. After some rearrangement of dummy variables, this gives

$$\begin{aligned} \lim_{\xi \rightarrow 0} (\partial_t + \nu_0 j^2 + \nu_0 |\mathbf{k} - \mathbf{j}|^2) \langle u_j^+ u_{k-j}^+ \rangle_c &= \lim_{\xi \rightarrow 0} 2M_j^+ \\ &\times \{ \langle u_p^- u_{j-p}^- u_{k-j}^+ \rangle_c + 2 \langle u_p^- u_{j-p}^+ u_{k-j}^+ \rangle_c + \langle u_p^+ u_{j-p}^+ u_{k-j}^+ \rangle_c \}. \end{aligned} \quad (25)$$

Applying the hypothesis as give by Eq. (20), it is easily seen that the first term on the right hand side of (25) is zero, since in the limit it involves the ensemble average of u_k^+ , while the second term gives rise to a term linear in u_k^- . The third term may be evaluated by iterating the above procedure to form a dynamical equation for $\langle u_p^+ u_{j-p}^+ u_{k-j}^+ \rangle_c$, which in turn gives rise to higher-order moments.

In general, we can show that a similar pattern occurs for all higher order moments involving only products of u^+ . That is, each such moment gives rise to a moment involving two u^- modes, which in general has to be zero for consistency in its wavevector arguments, a term linear in u_k^- , and a moment involving only u^+ modes of next higher order. Hence we may write the general result

$$M_{\alpha\beta\gamma}^- (\mathbf{k}) \int d^3 j \lim_{\xi \rightarrow 0} \langle u_\beta^+ (\mathbf{j}, t) u_\gamma^+ (\mathbf{k} - \mathbf{j}, t) \rangle_c = \int_{-\infty}^t ds A(\mathbf{k}, t - s) u_\alpha^- (\mathbf{k}, s), \quad (26)$$

where $A(\mathbf{k}, t - s)$ has the form

$$\begin{aligned} A(\mathbf{k}, t - s) &= \int d^3 j \exp[-(\nu_0 j^2 + \nu_0 |\mathbf{k} - \mathbf{j}|^2)(t - s)] \\ &\times \{ 4M_k^- M_j^+ \lim_{\{\cdot\} \rightarrow K_{\max}} \langle u_{j-p}^+ u_{k-j}^+ \rangle \\ &+ 24M_k^- M_j^+ L_{03}^{-1} M_p^+ \lim_{\{\cdot\} \rightarrow K_{\max}} \langle u_{p-q}^+ u_{j-p}^+ u_{k-j}^+ \rangle + \dots \}, \end{aligned} \quad (27)$$

$L_{03} \equiv \partial_t + \nu_0 p^2 + \nu_0 |\mathbf{j} - \mathbf{p}|^2 + \nu_0 |\mathbf{k} - \mathbf{j}|^2$, and where higher order terms are easily found by induction. Thus, in all, Eq. (23) for the low wavenumber modes may be written as

$$(\partial_t + \nu_0 k^2) u_k^- - \int_{-\infty}^t ds A(\mathbf{k}, t - s) u_\alpha^- (\mathbf{k}, s) = M_k^- u_j^- u_{k-j}^- + S(k|K_c). \quad (28)$$

In order to test the hypothesis, we make two approximations. First, we truncate the expansion of $A(\mathbf{k}, t)$ at lowest non-trivial order. This can be justified via the introduction of a *local* Reynolds number based on a length scale K_c^{-1} , the moment expansion being re-expressed as a power series in this parameter. Making the truncation in (26) and (27) leaves us with the expression

$$\begin{aligned} \lim_{\xi \rightarrow 0} M_{\alpha\beta\gamma}^- (\mathbf{k}) \langle u_\beta^+ (\mathbf{j}, t) u_\gamma^+ (\mathbf{k} - \mathbf{j}, t) \rangle_c &= \int_{-\infty}^t ds \exp[-(\nu_0 j^2 + \nu_0 |\mathbf{k} - \mathbf{j}|^2)(t - s)] \\ &\times 4 M_{\alpha\beta\gamma}^- (\mathbf{k}) M_{\beta\delta\epsilon}^+ (\mathbf{j}) \int d^3 p \lim_{\{\cdot\} \rightarrow K_{\max}} \langle u_\epsilon^+ (\mathbf{j} - \mathbf{p}, s) u_\gamma^+ (\mathbf{k} - \mathbf{j}, s) \rangle u_\delta^- (\mathbf{k}, s). \end{aligned} \quad (29)$$

For stationary, homogeneous, and isotropic turbulence we may write

$$\langle u_\epsilon^+ (\mathbf{j} - \mathbf{p}, s) u_\gamma^+ (\mathbf{k} - \mathbf{j}, s) \rangle = Q(|\mathbf{k} - \mathbf{j}|) D_{\epsilon\gamma}(\mathbf{k} - \mathbf{j}) \delta(\mathbf{k} - \mathbf{p}), \quad (30)$$

where $Q(k)$ is the spectral density and δ is the Dirac delta function. This leaves the question of how to perform the time integral

$$\int_{-\infty}^t ds \exp[-(\nu_0 j^2 + \nu_0 |\mathbf{k} - \mathbf{j}|^2)(t - s)] u_\delta^- (\mathbf{k}, s). \quad (31)$$

To do this we change the variable of integration from s to $T = t - s$, expand the resultant $u_\delta^- (\mathbf{k}, t - T)$ as a Taylor series about $T = 0$, and then truncate the expansion at zero order, this approach being based upon the physical idea that the u^- modes are slowly evolving on time scales defined by the inverse of $\nu_0 j^2 + \nu_0 |\mathbf{k} - \mathbf{j}|^2$.

We have investigated the validity of these approximations using results from direct numerical simulations performed on a 256^3 grid, with Taylor-Reynolds number $R_\lambda = 190$. Although these results are not conclusive, since at this resolution the simulations have a very limited inertial range (see [6,7]), they indicate that there is a range of K_c ($K_c \gtrsim 0.5 K_{\max}$) where both approximations give rise to error terms of less than unity, and that the magnitude of these errors will decrease as we increase R_λ to the large values where we may reasonably expect our hypothesis to hold.

With these approximations, the right hand side of (26) is simple to evaluate, and we are left with the final expression for the conditional average on the right hand side of (23)

$$\begin{aligned}
& M_{\alpha\beta\gamma}^-(\mathbf{k}) \lim_{\xi \rightarrow 0} \langle u_{\beta}^+(\mathbf{j}, t) u_{\gamma}^+(\mathbf{k} - \mathbf{j}, t) \rangle_c \\
& = 4M_{\alpha\beta\gamma}^-(\mathbf{k}) M_{\beta\delta\epsilon}^+(\mathbf{j}) \lim_{|\mathbf{k}-\mathbf{j}| \rightarrow K_{\max}} \frac{Q(|\mathbf{k}-\mathbf{j}|) D_{c\gamma}(\mathbf{k}-\mathbf{j})}{\nu_0 j^2 + \nu_0 |\mathbf{k}-\mathbf{j}|^2} u_{\delta}^-(\mathbf{k}, t), \quad (32)
\end{aligned}$$

which is linear in $u_{\mathbf{k}}^-$, meaning that it may be interpreted in terms of an increment to the viscosity.

In order to evaluate the limit, we make a first-order truncation of a Taylor series expansion in wavenumber of Q^+ about K_{\max} . In this way, we re-obtain the results previously obtained using the two-field theory of McComb and Watt [8]. As they showed, a Renormalization Group calculation based upon these equations gives a prediction for the Kolmogorov constant of 1.60 ± 0.01 , in good agreement with experiment, for $0.55K_{\max} \leq K_c \leq 0.75K_{\max}$. This calculation obtained the Kolmogorov exponent and pre-factor by assuming that the effective viscosity and its increment scale in the same way (which is true at the fixed point) and that the rate of energy transfer is renormalized. This latter assumption amounted, in our present terminology, to the neglect of $S(k|K_c)$ in Eq. (23).

A new justification of this step can now be offered as follows. The equation for the energy spectrum is obtained by multiplying the dynamical equation for $u_{\alpha}^-(\mathbf{k}, t)$ by $u_{\alpha}^-(-\mathbf{k}, t)$ and then performing an average over the full ensemble. Thus the effect of $S(k|K_c)$ is just

$$\langle S(k|K_c) u_{\alpha}^-(-\mathbf{k}, t) \rangle.$$

If we consider the form of $S(k|K_c)$ we see that each of the terms in the above expression involves a conditional average. In evaluating such terms we perform a double summation, firstly summing over all members with low wavenumber modes close to a particular member of the ensemble, and then repeating this summation for every member of the ensemble. Now, the initial ensemble was constructed according to the principle of equal *a priori* probabilities but this is no longer necessarily true of the composite ensemble which we are now considering. If it were true, then the terms making up $S(k|K_c)$ would vanish identically for all K_c . However, in view of the results of the Renormalization Group calculations [8], it seems likely that the contribution from $S(k|K_c)$ is small for K_c in the range $0.55K_{\max} \leq K_c \leq 0.75K_{\max}$.

Thus, for this range of cut-off wavenumbers, it would appear that the RG calculation of the effective viscosity [8] is valid in a heuristic sense.

Finally, it should be noted that this work does not suggest that $S(k|K_c)$ can be neglected in Eq. (23), which is the governing equation for a single realization. However it does suggest that, having averaged out the chaotic part to yield an effective viscosity, one should consider modeling the relationship of $S(k|K_c)$ to the $u_{\vec{k}}$ modes as predominantly deterministic. Work along these lines will be the subject of a separate communication.

Both authors acknowledge the support and facilities provided by the Isaac Newton Institute. We also wish to thank Alistair Young for providing results from numerical simulations. C.J. acknowledges the financial support of the Engineering and Physical Sciences Research Council.

REFERENCES

- [1] K.G. Wilson, *Rev. Mod. Phys.* **47**, 773 (1975).
- [2] W.D. McComb, W. Roberts, and A.G. Watt, *Phys. Rev. A* **45**, 3507 (1992).
- [3] Note that this differs from the formulation in Ref. [2], where each realization was time-dependent.
- [4] R.G. Deissler, *Phys. Fluids* **29**, 1453 (1986).
- [5] L. Machiels, *Phys. Rev. Lett.* **79**, 3411 (1997).
- [6] J.G. Brasseur and C.-H. Wei, *Phys. Fluids* **6**, 842 (1994).
- [7] P.K. Yeung and Y. Zhou, *Phys. Rev. E* **56**, 1746 (1997).
- [8] W.D. McComb and A.G. Watt, *Phys. Rev. A* **46**, 4797 (1992).

Recent Newton Institute Preprints

- NI98001-STA **N Linden and S Popescu**
Non-local properties of multi-particle density matrices
- NI98002-AMG **J-L Colliot-Thélène**
Un principe local-global pour les zéro-cycles sur les surfaces fibrés en coniques au-dessus d'une courbe de genre quelconque
- NI98003-AMG **RGE Pinch and HPF Swinnerton-Dyer**
Arithmetic of diagonal quartic surfaces, II
- NI98004-AMG **DR Heath-Brown**
The solubility of diagonal cubic diophantine equations
- NI98005-AMG **B Poonen and M Stoll**
The Cassels-Tate pairing on polarized Abelian varieties
- NI98006-AMG **R Parimala and V Suresh**
Isotropy of quadratic forms over function fields of curves over p -adic fields
- NI98007-AMG **E Peyre**
Application of motivic complexes to negligible classes
- NI98008-AMG **E Peyre**
Torseurs universels et méthode du cercle
- NI98009-RAG **JA Green**
Discrete series characters for $GL(n, q)$
- NI98010-DQC **K Zyczkowski, P Horodecki, A Sanpera et al**
On the volume of the set of mixed entangled states
- NI98011-DQC **K Zyczkowski and W Slomczyński**
Monge distance between quantum states
- NI98012-DAD **JA Sellwood, RW Nelson and S Tremaine**
Resonant thickening of disks by small satellite galaxies
- NI98013-DAD **GI Ogilvie and SH Lubow**
The effect of an isothermal atmosphere on the propagation of three-dimensional waves in a thermally stratified accretion disk
- NI98014-DAD **JA Sellwood and SA Balbus**
Differential rotation and turbulence in extended $H I$ disks
- NI98015-DAD **AM Fridman and OV Khoruzhii**
On nonlinear dynamics of 3D astrophysical disks
- NI98016-DQC **RE Prange, R Narevich and O Zaitsev**
Quasiclassical surface of section perturbation theory
- NI98017-DQC **A Sedrakyan**
Edge excitations of an incompressible fermionic liquid in a disorder magnetic field
- NI98018-DAD **CF Gammie**
Accretion disk turbulence
- NI98019-DAD **R Popham and CF Gammie**
Advection dominated accretion flows in the Kerr metric: II. Steady state global solutions
- NI98020-DAD **CF Gammie, R Narayan and R Blandford**
What is the accretion rate in NGC 4258?
- NI98021-DAD **CF Gammie**
Photon bubbles in accretion disks
- NI98022-DAD **EC Ostriker, CF Gammie and JM Stone**
Kinetic and structural evolution of self-gravitating, magnetized clouds: 2.5-dimensional simulations of decaying turbulence
- NI98023-DAD **JA Sellwood and EM Moore**
On the formation of disk galaxies and massive central objects